

It could be seen from Fig. 1 that changing trend of microbial biomass carbon was divided into two sections, and the demarcation point of water content was about 19.5%. Should water content be higher or lower than the demarcation point, microbial biomass carbon would decline. Briefly, microbial biomass carbon content increased (from 200 to 400 mg kg⁻¹) with the reduction of water content when soil moisture was higher than the demarcation point, while microbial biomass carbon content declined (from 400 to 25 mg kg⁻¹) with the reduction of water content when soil moisture was lower than the demarcation point, which indicated that 19.5% was the optimum water content for microbial biomass carbon in our experimental soil ecosystem.

Changing rate curve of microbial biomass carbon along with mass soil water content was shown in Fig. 2. The changing rate curve was obtained by differentiating the fitting curve in Fig. 1. We can see that the changing rate curve was divided into three sections (section A, B and C) by two demarcation points, which were 19.5 and 14.3% for water content, respectively. The demarcation point of 19.5% was the position that increase or decrease of microbial biomass carbon, while the demarcation point of 14.3% was the position of faster or slower of the decrease rate. In section A microbial biomass content increased with the reduction of water content, which partly attributed to the limit of soil microorganism activity when water content was higher than 19.5%. In section B decrease rate of microbial biomass carbon became faster and faster with the development of drought. In section C decrease rate of microbial biomass carbon became slower and slower as drought stress got more and more serious. In addition, with the development of extreme drought, the change rate of microbial biomass carbon tended to zero.

3.2 Drought effects on proportion of microbial biomass carbon in soil organic carbon

Figure 3 gave dynamic of the proportion of microbial biomass carbon in soil organic carbon during the whole drought stage. It was obvious that microbial biomass carbon comprised only about 1 ~ 4% of soil organic carbon. On the one hand, the changing

9

tendency of proportion of microbial biomass carbon in soil organic carbon was similar with the changing trend of microbial biomass carbon along water content. When water soil moisture was higher than the demarcation point, the proportion of microbial biomass carbon increased from 1.5 to 3.9% with reduction of water content. However, when soil moisture was lower than the demarcation point, the proportion decreased from 3.9 to 1.0% with the reduction of water content. On the other hand, the demarcation point of increase or decrease of the proportion was about 20.5% for water content, which was 1% higher than the demarcation point of microbial biomass carbon changing along with water content (19.5%).

3.3 Rehabilitation of different drought soil ecosystem after rainstorm

The rehabilitation of soil microbial biomass carbon under moderate drought scenario after rainstorm stimulation was shown in Fig. 4a. We could see that microbial biomass carbon recovered at about 16% of mass water content, which was before water content reduced to 14.3%. At this time relative water content was 60% and soil ecosystem was not stressed by drought. On one hand, content of microbial biomass carbon in watered soil was even more than it was before rehydration at the same water content when water content was lower than 16%. On the other hand, when water content was lower than 16%, content of microbial biomass carbon increased before it decreased and water content was 15% at the point that microbial biomass carbon was the most. What more, when soil water was dried to 12%, the content of microbial biomass carbon was as much as it was at the 15% of water content in the soil that was not watered.

The rehabilitation of soil microbial biomass carbon under severe drought scenario after rainstorm stimulation was shown in Fig. 4b. The results showed that content of microbial biomass carbon increased gradually and barely recovered until soil water content reduced to 14% around. When water content was lower than that point, content of microbial biomass carbon in rehydrated soil was almost as much as it was in the soil that was not watered.

2006). Figures 1 and 3 showed that changing trend of proportion of microbial biomass carbon in soil organic carbon lag behind microbial biomass carbon changes along with drought stress, suggesting that microbial biomass carbon responded faster than the proportion of microbial biomass carbon in soil organic carbon to drought stress. On one hand, labile soil organic carbon was closely associated with root productivity (Ros et al., 2009; Rui et al., 2011). In this research when water content reduced to 20.5 %, it was just the time that vegetative growth of above ground plant was vigorous. At the same time root productivity and belowground root biomass increased, resulting in an increase of soil organic carbon. However, microbial biomass carbon had not increased as much as soil organic carbon, so proportion of microbial biomass carbon in soil organic carbon decreased. On the other hand, it had been indicated that, as a response to drought, some soil bacteria are able to synthesize exopolysaccharides (Kohler et al., 2009), resulting in soil organic carbon increase. From the above results, it obtained that change of microbial biomass carbon proportion in soil organic carbon lag behind that of microbial biomass carbon under drought stress.

4.3 Rehabilitation of soil ecosystem under different dry-wet scenarios

From Fig. 4 we can see that rehabilitation of soil ecosystem was positive in moderate and severe drought soil ecosystem and negative in extreme drought soil ecosystem. The results in Fig. 4a showed that soil microbial biomass carbon interfered by moderate drought and then experienced rainstorm could recover before water content reduced to 14.3 %, suggesting that soil ecosystem could recover under this drought-wet scenario. When water content was lower than 16 %, content of microbial biomass was higher than it was before rehydration at the same water content. We can come to the conclusion that some drought tolerant microorganism had already adapted to this drought stress, their tolerance to drought and ratio in all soil microorganisms were both improved. Content of microbial biomass carbon showed a small peak, suggesting that ecological amplitude of drought tolerant microorganism widened under moderate drought stress. As a result, the soil ecosystem was well tolerated to moderate drought stress, its function

13

and activity might be improved also. Results in Fig. 4b showed that content of microbial biomass carbon could barely recovered, suggesting that soil ecosystem influenced by severe drought was at the edge of rehabilitation. The ecosystem could not adapt to the severe drought stress and its tolerance to drought stress was not improved. Results in Fig. 4c showed that content of microbial biomass carbon could not recover within the experimental time, which indicated that soil ecosystem that stressed by extreme drought could not restore within a short time. The eco-hydrological processes were interrupted and ecosystem function and structure were damaged and could not recover though they had experienced a rainstorm. It has been proved that, when water content was less than a certain value, rewetting could lead to microbial stress because its tolerance rapid changes in microorganism osmotic potential, resulting in cell lysis (Van Gestel et al., 1992). As a consequence, microbial biomass carbon went on declining and could not give better resistance to drought. What more, microbial rehabilitation was different in different types of ecosystem and soil terms (David et al., 2013; Chaer et al., 2009; Lacombe et al., 2009; Mader et al., 2002; Van Overbeck et al., 1995).

4.4 An indicator of irrigating – dynamics and demarcation points of microbial biomass carbon along with mass water content of soil

When water content was lower than 14.3 %, microorganism reproduction and substrate utilization in soil was influenced, as well as decomposition of plant and animal residues (Johnson et al., 2003), nutrient cycling (Balsler and Firestone, 2005), soil fertility maintaining and formation of soil aggregates (Gillerke, 1997), which resulted in function and structure weaken in soil ecosystem. On the contrary, high concentration of microbial biomass carbon was characteristic of a sustainable ecosystem. Therefore, the changing tendency and demarcation point of soil microbial biomass carbon along with mass water content of soil could be used to demonstrate alters and degradation of soil ecosystem as well as the irrigation requirement of crops.

It is proved that when soil moisture content was lower than 55 % of field capacity, farmland should be irrigated. Here field capacity of experimental soil was 27 % and the

14

- Chaer, G., Fernandes, M., and Myrold, D.: Comparative resistance and resilience of soil microbial communities and enzyme activities in adjacent native forest and agricultural soils, *Microb. Ecol.*, 58, 414–424, 2009.
- Christ, M. J. and David, M. B.: Temperature and moisture effects the production of dissolved organic carbon in a podosol, *Soil Biol. Biochem.*, 28, 1191–1199, 1996.
- 5 Cong, Z. T., Yao, B. Z., and Ni, G. H.: Water demand forecasting for major crops in China under SRA1B scenario, *Adv. Water Sci.*, 22, 38–43, 2011.
- David, R., Miren, L., and Alain, O.: Soil biochemical properties and microbial resilience in agroforestry systems: effects on wheat growth under controlled drought and flooding conditions, *Sci. Total. Environ.*, 463–464, 51–60, 2013.
- 10 Drenovsky, R. E., Vo, D., and Graham, K. J.: Soil water content and organic carbon availability are major determinants of soil microbial community composition, *Microb. Ecol.*, 48, 424–430, 2004.
- Fierer, N. and Schimel, J. P.: A proposed mechanism for the pulse in carbon dioxide production commonly observed following the rapid rewetting of a dry soil, *Soil Sci. Soc. Am. J.*, 67, 798–805, 2003.
- 15 Fu, H. L., Wang, S. R., and Han, S. J.: Effects of alternate wet and dry conditions on microbial activity and flora in broad leaved *Pinus koraiensis* of Changbai Mountain, *J. Northeast Forest. Univ.*, 37, 80–86, 2009.
- 20 Gestel, M. V., Ladd, J. N., and Amato, M.: Microbial biomass responses to seasonal change and imposed drying regimes at increasing depths of undisturbed topsoil profiles, *Soil Biol. Biochem.*, 24, 103–111, 1992.
- Giller, K. E., Beare, M. H., and Lavell, E. P.: Agricultural intensification soil biodiversity and agro-ecosystem function, *Appl. Soil Ecol.*, 6, 3–16, 1997.
- 25 Houghton, J. T., Ding, Y., and Griggs, D. J.: *Climate change, The Scientific Basis*, Cambridge University Press, New York, 881 pp., 2001.
- Housman, D. C., Yeager, C. M., and Darby, B. J.: Heterogeneity of soil nutrients and subsurface biota in a dry land ecosystem, *Soil Biol. Biochem.*, 39, 2138–2149, 2007.
- Hueso, S., Garcia, C., and Hernandez, T.: Severe drought conditions modify the microbial community structure, size and activity in amended and un-amended soils, *Soil Biol. Biochem.*, 30, 167–173, 2012.
- 30 Insam, H., Mitchell, C. C., and Dormaar, J. F.: Relationship of soil microbial biomass and activity with fertilization practice and crop yield of three soils, *Soil Biol. Biochem.*, 23, 459–564, 1991.

- Jenkinson, D. S. and Powlson, D. S.: Microbial biomass in soil: measurement and turnover, *Soil Biochem.*, 5, 415–471, 1981
- Jensen, K. D. and Beier, C.: Effects of experimental drought on microbial processes in two temperate heathlands at contrasting water conditions, *Appl. Soil Ecol.*, 24, 165–176, 2003.
- 5 Joergensen, R. G.: The fumigation-extraction method to estimate soil microbial biomass: extraction with 0.01 M CaCl_2 , *Agro-Biol. Res.*, 48, 319–324, 1995.
- Johnson, D., Booth, R. E., and Ehiteley, A. S.: Plant community composition affects the biomass, activity and diversity of microorganisms in limestone grassland soil, *Eur. J. Soil Sci.*, 54, 671–677, 2003.
- 10 Johnson, M. J., Lee, K. Y., and Scow, K. M.: DNA fingerprinting reveals links among agricultural crops, soil properties, and the composition of soil microbial communities, *Geoderma*, 114, 279–303, 2003.
- Joshua, P. S., Jay, M. G., and Joy, S. C. C.: Moisture effects on microbial activity and community structure in decomposing birch litter in the Alaskan taiga, *Soil Biol. Biochemical.*, 31, 831–838, 1999.
- 15 Kennedy, A. C. and Smith, K. L.: Soil microbial diversity and the sustainability of agricultural soils, *Plant Soil*, 170, 75–86, 1995.
- Kohler, J., Caravaca, F., and Roldán, A.: Effect of drought on the stability of rhizosphere soil aggregates of *Lactuca sativa* grown in a degraded soil inoculated with PGPR and AM fungi, *Appl. Soil Ecol.*, 42, 160–165, 2009.
- 20 Kong, B., Sun, B., and Zheng, X. Q.: Effects of hydrothermal conditions and fertilization on microbial metabolic characteristic of microbial community in black soil, *Acta Pedol. Sin.*, 46, 100–106, 2009.
- Lacombe, S., Bradley, R. L., and Hamel, C.: Do tree-based intercropping systems increase the diversity and stability of soil microbial communities, *Agr. Ecosyst. Environ.*, 31, 25–31, 2009.
- 25 Lal, S., Bagdi, D. L., Kakralya, B. L., Jat, M. L., and Sharma, P. C.: Role of brassinolide in alleviating the adverse effect of drought stress on physiology, growth and yield of green gram (*vigna radiata*.) genotypes, *Legume Res.*, 36, 359–363, 2013.
- Li, X. Z. and Sarah, P.: Arylsulfatase activity of soil microbial biomass along a Mediterranean arid transport, *Soil Biol. Biochem.*, 35, 925–934, 2003a.
- 30 Li, X. Z. and Sarah, P.: Enzyme activities along a climatic transect in the Judean Desert, *Catena*, 53, 349–363, 2003b.

- Liu, G. S., Xu, D., and Liu, Y.: Influence of spatial scale variation on ET temporal up scaling methods, *J. Hydraul. Eng.*, 43, 999–1003, 2012.
- Liu, X. Y. and Lin, E. D.: Effects of climate change on water demand of major crops in North China, *J. Hydraul. Eng.*, 2, 77–84, 2004.
- 5 Liu, Y. Y., Yao, H. Y., and Huang, C. Y.: Influence of soil moisture regime on microbial community diversity and activity in a paddy soil, *Acta Pedol. Sin.*, 43, 828–834, 2006.
- Liu, Z. F., Fu, B. J., and Zheng, X. X.: Plant biomass, soil water content and soil N:P ratio regulating soil microbial functional diversity in a temperate steppe: a regional scale study, *Soil Biol. Biochem.*, 42, 445–450, 2010.
- 10 Mäder, P., Fließbach, A., and Dubois, D.: Soil fertility and biodiversity in organic farming, *Science*, 296, 1694–1697, 2002.
- Muhammad, S., Evgenia, B., and Abad, C.: Drought effects on microbial biomass and enzyme activities in the rhizosphere of grasses depend on plant community composition, *Appl. Soil Ecol.*, 48, 38–44, 2011.
- 15 Nielsen, N. M., Winding, A., and Binnerup, S.: Microorganisms as indicators of soil health Ministry of the Environment, National Environmental Research Institute, Aarhus, 15–16, 2002.
- Panikwv, N. S.: Understanding and prediction of soil microbial community dynamics under global change, *Appl. Soil Ecol.*, 11, 161–176, 1999.
- 20 Paul, E. A. and Clark, F. E.: *Soil Microbiology and Biochemistry*, Academic Press, San Diego, 245–264, 1996.
- Rattan, L.: Food security in a changing climate, *Ecohydrol. Hydrobiol.*, 13, 8–21, 2013.
- Rietz, D. N. and Haynes, R. J.: Effects of irrigation induced salinity and sodicity on soil microbial activity, *Soil Biol. Biochem.*, 35, 845–854, 2003.
- Singh, K., Singh, B., and Singh, R. R.: Effect of land rehabilitation on physicochemical and microbial properties of a sodic soil, *Catena*, 109, 49–57, 2013.
- 25 SL 424-2008: Standard of classification for drought severity [S], China's Water Conservancy and Hydropower Press, Beijing, 2009.
- Somova, L. A. and Pechurkin, N. S.: Functional, regulatory and indicator features of microorganisms in manmade ecosystems, *Adv. Space Res.*, 27, 1563–1570, 2001.
- 30 Sowerby, A., Bridget, E., and Claus, B.: Microbial community changes in health and soil communities along a geographical gradient: interaction with climate change manipulations, *Soil Biol. Biochem.*, 37, 1806–1813, 2005.

- Tejada, M., García, C., Gonzalez, J. L., and Hernandez, M. T.: Use of organic amendments as a strategy for saline soil remediation: influence on the physical, chemical and biological properties of soil, *Soil Biol. Biochem.*, 38, 1413–1421, 2006.
- 5 Van, M. J. M., Tietema, A., and Van, L. E. E.: Microbial dynamics and litter decomposition under a changed climate in a Dutch health land, *Appl. Soil Ecol.*, 38, 119–127, 2008.
- Wang, L. C., Tamai, R., and Nagata, M.: Effects of water and salty to activity of soil microbial, *Reclaim. Rice Cultiv.*, 3, 40–42, 1998.
- Wilkinson, S. C., Anderson, J. M., and Scardelis, S. P.: PLFA profiles of microbial communities in decomposing conifer litters subject to moisture stress, *Soil Biol. Biochemical*, 34, 189–200, 2002.
- 10 Wu, J. S., Lin, Q. M., and Huang, Q. Y.: *Methods and Applications of Soil Microbial Detection*, Meteorological Press, Beijing, China, 54–57, 2006.
- Yao, H. Y. and Huang, C. Y.: *Soil Microbial Ecology and its Experimental Techniques*, Science Press, Beijing, 139–140, 2006.
- 15 Yuan, B. C., Li, Z. Z., Liu, H., Gao, M., and Zhang, Y. Y.: Microbial biomass and activity in salt affected soils under arid conditions, *Appl. Soil Ecol.*, 35, 319–328, 2007.
- Zak, D. R., Holmes, W. E., and White, D. C.: Plant diversity, microbial communities, and ecosystem function: are there any links, *Ecology*, 84, 2042–2050, 2003.
- 20 Zelles, L.: Fatty acid patterns of phospholipids and lipopolysaccharides in the characterization of microbial communities in soil: a review, *Biol. Fert. Soils*, 29, 111–129, 1999.

Table 1. Soil physical, chemical and biological properties.

| Soil parameters | Value | Units |
|----------------------|--------------|---------------------|
| pH | 7.8 | – |
| organic matter | 20 ~ 153 | % |
| total nitrogen | 0.075 | % |
| available nitrogen | 0.121 | mg kg ⁻¹ |
| total phosphorus | 1.912 | % |
| available phosphorus | 38.04 | mg kg ⁻¹ |
| total potassium | 58.41 | % |
| available potassium | 134.62 | mg kg ⁻¹ |
| microbial carbon | 18.2 ~ 373.9 | mg kg ⁻¹ |
| microbial nitrogen | 49.45 | mg kg ⁻¹ |
| unit weight of soil | 2.78 | g cm ⁻³ |
| clay | 12.83 | % |
| silt | 28.92 | % |
| sand | 58.25 | % |
| soil field capacity | 27 | % |

Table 2. Standard of classification for drought severity.

| Drought severity | Relative soil water content |
|------------------|-----------------------------|
| not drought | $R > 60\%$ |
| mild drought | $60\% \geq R > 50\%$ |
| moderate drought | $50\% \geq R > 40\%$ |
| severe drought | $40\% \geq R > 30\%$ |
| extreme drought | $30\% \geq R$ |




Fig. 1. Dynamics and fitting curve of microbial biomass carbon along with mass water content of soil (■ is measured data of soil microbial biomass carbon and correspond mass water content; red curve is the fitting curve).




Fig. 2. Chang rate curve of microbial biomass carbon along with mass water content of soil.




Fig. 3. Dynamics of proportion of microbial biomass carbon in soil organic carbon along with mass water content (■ is measured data of proportion of microbial biomass carbon in soil organic carbon correspond to mass water content).




Fig. 4. Rehabilitation of microbial biomass carbon after rainstorm stimulation in moderate drought scenario (A), severe drought scenario (B) and extreme drought scenario (C); ■ is measured data of microbial biomass carbon correspond to soil mass water content before rehydration under drought scenarios; ▲ is measured data of microbial biomass carbon after rehydration; Red curves are the fitting curves between microbial biomass carbon and mass water content of soil.




Fig. 5. Comparison of rehabilitation of microbial biomass carbon content after rainstorm stimulation under moderate drought scenario (■), severe drought scenario (●) and extreme drought scenario (▲).