

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez^{1,2,3}, O. Morales¹, F. Cisneros¹, W. Bauwens², and G. Wyseure³

¹Programa para el manejo del Agua y del Suelo (PROMAS), Universidad de Cuenca, Cuenca, Ecuador

²Department of Hydrology and Hydraulic Engineering, Earth System Sciences Group, Vrije Universiteit Brussel (VUB), Brussels, Belgium

³Department of Earth and Environmental Science, KU Leuven, Leuven, Belgium

Received: 29 September 2015 – Accepted: 21 October 2015 – Published: 3 November 2015

Correspondence to: V. Iñiguez (vicente.iniguez@ucuenca.edu.ec)

Published by Copernicus Publications on behalf of the European Geosciences Union.

HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The neotropical Andean grasslands above 3500 m.a.s.l. known as “páramo” offer remarkable ecological services for the Andean region. Most important is the water supply – of excellent quality – to many cities and villages established in the lowlands of the inter-Andean valleys and to the coast. However, the páramo ecosystem is under constant and increased threat by human activities and climate change. In this paper we study the resilience of its soils for drought periods during the period 2007–2013. In addition, field measurements and hydrological conceptual modelling at the catchment-scale are comparing two contrasting catchments in the southern Ecuadorian Andes. Both were intensively monitored during two and a half years (2010–2012) in order to analyse the temporal variability of the soil moisture storage. A typical catchment on the páramo at 3500 m.a.s.l. was compared to a lower grassland one at 2600 m.a.s.l. The main aim was to estimate the resilience capacity of the soils during a drought period and the recovery during a subsequent wet period. Local soil water content measurements in the top soil (first 30 cm) through TDR were used as a proxy for the catchment’s average soil moisture storage. The local measurements were compared to the average soil water storage as estimated by the probabilistic soil moisture (PDM) model. This conceptual hydrological model with 5 parameters was calibrated and validated for both catchments. The study reveals the extraordinary resilience capacity of this type of shallow organic soils during the droughts in 2009 and 2010. During these droughts, the soil water content dropped from a normal value of about 0.80 to $\sim 0.60 \text{ cm}^3 \text{ cm}^{-3}$, while the recovery time was only two to three months.

1 Introduction

In the northern Andean landscape, between ca. 3500 and 4500 m.a.s.l., an “alpine” neotropical grassland ecosystem – locally known as “páramo” – covers the mountains. Their major ecological characteristics has been documented by several authors (e.g.

HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

increase in the mean annual precipitation in the region that is of interest to our study. On the other hand, the carbon storage and the water yield could be reduced by the higher temperatures and the higher climate variability. However, the uncertainties on the potential impact of the climate change remain high (Buytaert and De Bièvre, 2012; Buytaert et al., 2010).

Another important factor for the region are the El Niño Southern Oscillation events. The Amazon river basin, with its headwaters in the Andes, has indeed faced severe droughts in 2005 and 2010 (Lewis et al., 2011; Phillips et al., 2009). These dry periods have been attributed to the severe El Niño Southern Oscillation events and north-west displacement of the intertropical convergence zone (ITCZ) (Marengo et al., 2008, 2011).

The El Niño Southern Oscillation events not only have an impact on the ecohydrology of the forest area of the Amazon River basin but also in its headwater or páramos. Indeed, the droughts in the páramo have been observed during the months with less rainfall (from September to December), which coincide with the displacement of the ITCZ and by the Pacific and Atlantic anomalies (Buytaert et al., 2006b; Vuille et al., 2000). Thus, the climate variability in the mountains is exacerbated by these climate global events.

Since 2004 droughts in the páramo took place in 2005, 2009, 2010, 2011 and 2012, of which the first three years have been classified as “el Niño” years. The drought periods in the páramo had a negative impact on the water supply and on the economy of the whole region that depends on water supply from the Andes. For instance, the water levels in the reservoirs of the main hydropower projects reached their lowest values as a consequence of the drought between December 2009 and February 2010. This caused several, intermittent, power cuts in many regions of Ecuador.

The hydrological capacity of the páramo resides on its soils. Therefore the present study investigates the response of páramo’ soils to drought and compares with other soils on grasslands at lower altitude in the same region. The drought analysed is a soil moisture drought as defined by Van Loon (2015). The resilience or resistance

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

to drought is defined here as the time needed to recover to its pre-drought state of water content once that rainfall has started in a continuous way to exceed the vegetation water demand. The pre-drought state of soil water content is estimated from longer term periods whereby rainfall exceeds the vegetation requirements. The observation period includes the droughts of 2009, 2010, 2011 and 2012 together with intermediate wet periods. The analysis is done with special focus on the 2010 drought.

The hydrological drought is compared and related to this soil water drought. For this purpose, the water balance components of two experimental catchments – one with and one without páramo – were investigated by means of experimental measurements and by means of a hydrological model. The experimental work included the measurement of rainfall, flow and soil moisture. For the modelling, a parsimonious conceptual hydrological model – using the Probability Distributed Moisture simulator (PDM) was calibrated and validated for each experimental catchment. The PDM model allows to analyse the spatial variability of the soil water content as well as the maximum storage capacity at the catchment scale. Therefore, the representativeness of the point measurements of soil water content can be assessed by means of this model.

Our main hypothesis is that experimental monitoring combined with mathematical models enables the quantification of the sensitivity – or resilience – to drought of the land cover and soil systems in the Andes above 2600 m a.s.l.

2 Materials

2.1 The study area

The catchments under study are located in the southwest highlands of the Paute river basin, which drains to the Amazon River (Fig. 1). These highlands form part of the Western Cordillera in the Ecuadorian Andes with a maximum altitude of 4420 m a.s.l. The study area comprises a mountain range from 2647 until 3882 m a.s.l. Two catchments have been selected from this region: Calluancay and Cumbe.

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Several types of soils can be identified in Cumbe and Calluancay, which are mainly conditioned by the topography. Dercon et al. (1998, 2007) have described the more common toposequences in the southern Ecuadorian Andes according to the WRB classification (FAO/ISRIC/ISSS, 1998). Cumbe has a toposequence of soils from Vertic Cambisols, located in the alluvial area, surrounded by Dystric Cambisols at the hillslopes in the lower and middle part of the catchment. Eutric Cambisols or Humic Umbrisols extend underneath the forest patches between 3000 and 3300 m a.s.l. The highest part of the catchment – from 3330 until 3467 m a.s.l. – is covered by Humic Umbrisols or Andosols.

In contrast, Calluancay hosts only two groups of organic soils under páramo: Andosols (in the higher and steeper parts) and Histosols (in the lower and gentler parts of the catchment). The soils were formed as the geology of Calluancay is characterized by igneous rocks such as andesitic lava and pyroclastic igneous rock (mainly the Quimsacocha and Tarqui formations, dating from the Miocene and Pleistocene respectively), forming an impermeable bedrock underneath the catchment. In the Cumbe catchment, the highlands and some areas of the middle part (about 55 % of the area) are characterized by pyroclastic igneous rocks (mainly the Tarqui formation). The valley area (37 % of the basin) is covered by sedimentary rocks like mudstones and sandstones (mainly the Yunguilla formation, dating from the upper Cretaceous). Only 8 % of the Cumbe catchment comprises alluvial and colluvial deposits, which date from the Holocene (Hungerbühler et al., 2002).

2.2 The monitoring data

An intensive monitoring with a high time resolution was carried out in the study area during 28 months.

The gauging station at the outlet of Cumbe consists of a concrete trapezoidal supercritical-flow flume (Kilpatrick and Schneider, 1983) and a water level sensor (WL16 – Global Water). Logging occurs at a 15 min time interval. Regular field measurements of the discharge were carried out to cross-check the rating curve.

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

The measurements at Calluancay are part of a larger hydrological monitoring network maintained by PROMAS. Water levels are logged every 15 min at two gauging stations, which consist of a concrete V-shaped weir with sharp metal edges and a water level sensor (WL16- Global Water). The first station is installed at the outlet of the catchment. The second gauging station monitors an irrigation canal to which water is diverted from the main river. The gauging station was installed where the canal passes the water divide of the catchment. So, the total discharge can be evaluated.

For the Calluancay, rainfall is measured by a tipping bucket rain gauge (RG3M-Onset HOBO Data Loggers) located inside the catchment and with a resolution of 0.2 mm.

Three similar rain gauges were installed in the larger Cumbe catchment and located at the high, middle and lower part of the catchment. The areal rainfall for Cumbe was calculated with the inverse distance weighing (IDW) method, using the R implementation of GSTAT (Pebesma, 2004).

In each experimental catchment an automatic weather station measured the meteorological variables such as air temperature, relative humidity, solar radiation and wind speed at a 15 min time interval. These stations were used to estimate the potential reference evapotranspiration according to the FAO–Penman–Monteith equation.

2.3 The physical characteristics of the soil

In both catchments, the soil moisture content of the top soil layer was measured by means of time domain reflectometry (TDR) probes at representative sites in the vicinity of the weather stations. In both catchments, 6 TDR probes were installed in each plot. The TDRs were installed vertically from the soil surface with a length of 30 cm and logged at 15 min time intervals. In Calluancay, every fortnight soil water content was also measured by sampling from November 2007 until November 2008. In this catchment the TDR time series was from May 2009 until November 2013. In Cumbe, the TDR-time series extends from July 2010 until November 2012.

For Cumbe and Calluancay, the TDR measurements were calibrated based on soil samples ($R^2 = 0.79$ and 0.80 respectively). In addition, the curves were regularly cross-validated by undisturbed soil samples during the monitoring period.

The soil water retention curves were determined based on undisturbed and disturbed soil samples collected near to the TDR probes. In the laboratory, pressure chambers in combination with a multi-step approach allowed to define pairs of values for moisture (θ) and matric potential (h). The soil water retention curve model proposed by van Genuchten (1980) was fitted on the data.

3 Methods

3.1 The water balance based on the experimental data

The soil water balance of the root zone in each catchment over a selected time interval is estimated by the following equation:

$$\frac{dS_r}{dt} = P - E_a - Q_o - Q_l + C_r - D_p \quad (1)$$

where dS_r = the storage variation in the root zone during the time interval [mm], dt = the length of the time interval [days], P = the precipitation intensity during the time interval [mm day^{-1}], E_a = the actual evapotranspiration rate during the time interval [mm day^{-1}], Q_o = the overland flow during the time interval measured at the outlet of the catchment [mm day^{-1}], Q_l = the lateral flow during the time interval [mm day^{-1}], C_r = the capillary rise from a water table during the time interval [mm day^{-1}] and D_p = the deep percolation rate during the time interval [mm day^{-1}].

In the Cumbe catchment some water is diverted from the river for irrigation. As a result the flow at the outlet is reduced by the amount of irrigation. This irrigation is mainly concentrated in the valley and is rather informal by small farmer constructed

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



offtakes without major hydraulic structures. In addition there are no irrigation associations present and therefore an estimation of this withdrawal is very difficult.

Based on geological data, in Calluancay the deep percolation and capillary rise are considered to be negligible since the soils overlay bedrock consisting of igneous rocks with limited permeability. In páramos, the saturation overland flow is the dominant flow processes of runoff generation (Buytaert and Beven, 2011). The stream discharge (Q) at the outlet of the catchment thus comprises mainly overland flow and lateral flow.

In Cumbe, a surface-based electrical resistivity tomography test (Koch et al., 2009; Romano, 2014; Schneider et al., 2011) revealed no significant shallow groundwater for the alluvial area. In addition, the flat alluvial area near the catchment outlet is very small (2.7 % of the catchment area). Therefore, the terms D_p and C_r are also regarded to be negligible.

Based on the soil texture in Cumbe (clay) it is inferred that the infiltration overland flow is the dominant flow process of runoff generation. As a result, the stream discharge in Cumbe will be constituted, as in Calluancay, by two kinds of flows: overland and lateral flow.

Considering that the overland flow (Q_o) and the lateral flow constitute the observed river flow, Q , the water balance in our two catchments can thus be written as:

$$\frac{dS_c}{dt} = P - E_a - Q \quad (1a)$$

where dS_c = the average storage variation in the soil of the catchment during the time interval [mm] and Q = the total runoff leaving the catchment during the time interval [mm day⁻¹].

If we consider that P and Q are measured and we assume that the change of storage can be estimated based on the TDR measurements in our sampling points (as we will show in the results section), the actual evapotranspiration is the only unknown in this equation.

HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



During wet periods the water content in the root zone is constant at field capacity. The continuity equation may then be reduced to:

$$E_a = P - Q \quad (1b)$$

3.1.1 The potential evapotranspiration

The FAO–Penman–Monteith approach (Allen et al., 1998) is used to estimate the potential evapotranspiration of a reference crop (grass):

$$E_p = \frac{0.408\Delta(R_n - G_n) + \gamma \frac{900}{T+273} u_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34u_2)} \quad (2)$$

where E_p = the potential reference evapotranspiration [mm day^{-1}], R_n = the net radiation at the crop surface [$\text{MJ m}^{-2} \text{day}^{-1}$], G_n = the soil heat flux density [$\text{MJ m}^{-2} \text{day}^{-1}$], T = the mean daily air temperature at 2 m height [$^{\circ}\text{C}$], u_2 = the wind speed at 2 m height [m s^{-1}], e_s = the saturation vapour pressure [kPa], e_a = the actual vapour pressure [kPa], $e_s - e_a$ = the saturation vapour pressure deficit [kPa], Δ = the slope of the vapour pressure curve [$\text{kPa } ^{\circ}\text{C}^{-1}$] and γ = the psychrometric constant [$\text{kPa } ^{\circ}\text{C}^{-1}$].

The suitability of the FAO–Penman–Monteith approach for high altitudinal areas has been evaluated by Garcia et al. (2004). They found that the FAO-approach gives the smallest bias (-0.2 mm day^{-1}) as compared to lysimetric measurements.

The measurements of the solar radiation in our experimental catchments were not consistent and appeared to be unreliable. Therefore, the FAO–Penman–Monteith estimation for E_p was used with the solar radiation estimated by means of the Hargreaves–Samani equation (Hargreaves and Samani, 1985) using the daily maximum and minimum air temperature:

$$R_s = R_a c (T_{\max} - T_{\min})^{0.5} \quad (3)$$

where R_s = the solar radiation [$\text{MJ m}^{-2} \text{day}^{-1}$], R_a = the extra-terrestrial solar radiation [$\text{MJ m}^{-2} \text{day}^{-1}$], c = an empirical coefficient [–] and T_{\max} , T_{\min} = the daily maximum

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and minimum air temperature respectively [$^{\circ}\text{C}$]. According to Hargreaves and Samani (1985) “ c ” has a value of 0.17 for inland areas.

3.1.2 The actual evapotranspiration

The FAO–Penman–Monteith approach is used to calculate the potential evapotranspiration of a reference crop (normally grass) under stress free conditions without water limitation (E_p). This reference crop evapotranspiration can be converted to the evapotranspiration of another vegetation type by means of a vegetation coefficient k_v . During dry periods, with water stress, the vegetation extracts less water as compared to the vegetation requirement. The relative reduction of the evapotranspiration due to this may be expressed by a water stress coefficient k_s .

The actual evapotranspiration, E_a , can thus be calculated as:

$$E_a = k_s \cdot k_v \cdot E_p \quad (4)$$

In general, k_v is time-dependent, as it is linked to the growth cycle of the vegetation and thus to the season. For the páramo, this seasonality may be neglected as the grasses are slow-growing and perennial. In our study we therefore calculate a constant k_v by considering wet periods (during which $k_s = 1$) and using Eq. (4), in combination with the equations 1b (for calculating E_a) and 2 (for calculating E_p). Hereby, the wet periods were identified based on the TDR measurements of the soil water content.

Below the critical water content, E_a becomes less than the vegetation requirement and the soil water stress coefficient may be calculated as (Seneviratne et al., 2010):

$$k_s = 1 - \frac{\theta_{\text{crit}} - \theta_{\text{act}}}{\theta_{\text{crit}} - \theta_{\text{wp}}} \quad (5)$$

where θ_{crit} is the critical soil water content, θ_{act} is the actual soil water content and θ_{wp} is the permanent wilting point of the soil.

With k_v derived during wet periods as described above, k_s can now be estimated as a function of the actual soil moisture content by considering the (daily) water balance

HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



during dry periods. Hereto, we combine the Eqs. (1a), (4), (2) and (5). If we consider that the permanent wilting point can be derived from the soil water retention curve, based on the soil and vegetation characteristics, the critical soil water content is the only parameter that needs to be determined.

3.2 The actual evapotranspiration estimated by hydrological modelling

The actual evapotranspiration estimation based on local soil water measurements is compared to the actual evapotranspiration at catchment-scale calculated by the PDM model (Moore and Clarke, 1981; Moore, 1985). This hydrological model will be used to assess the impact of the soil moisture on the evapotranspiration. The PDM is a lumped rainfall–runoff model and its structure consists of two modules. The first is a soil moisture accounting (SMA) module which is based on a distribution of soil moisture storages with different capacities used to account for heterogeneity in the catchment. The probability distribution used is the Pareto distribution. The second part of the model structure is a routing module which consist of two linear reservoirs in parallel in order to consider the fast and slow flow pathways respectively. As in our study we consider small basins at a daily time step, the routing component is not so critical. The PDM model has been implemented within a MATLAB toolbox with the option of calculating the actual evapotranspiration E_a as a function of the potential evaporation rate E_p , and the soil moisture deficit (Wagener et al., 2001):

$$E_a = \left\{ 1 - \left[\frac{(S_{\max} - S(t))}{S_{\max}} \right] \right\} \cdot E_p \quad (6)$$

where, S_{\max} is the maximum storage and $S(t)$ is the actual storage at the beginning of the interval. A description of the model parameters is provided in Table 2.

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Results and discussion

4.1 The potential evapotranspiration derived from soil water observations

4.1.1 The potential evapotranspiration

The potential reference evapotranspiration (E_p) for the period from 16 July 2010 until 15 November 2012 was calculated by the FAO–Penman–Monteith approach with the solar radiation estimated by Hargreaves–Samani. The daily average of E_p for Calluancay and Cumbe was 2.35 and 3.04 mm day⁻¹ respectively. The temporal variation of E_p is depicted in Fig. 2. It reveals a sinusoidal pattern with higher atmospheric evaporative demand during the drier months (from August to March) and a lesser demand during the subsequent wet periods (from April to July). E_p ranged between 0.76 and 4.17 mm day⁻¹ for Calluancay and between 1.56 and 4.62 mm day⁻¹ for Cumbe. The difference can be attributed to the altitude difference between both catchments, with 900 m difference in elevation. The daily average minimum and maximum temperatures in Calluancay were 3.0 and 10.2 °C respectively. While, in Cumbe they were 7.8 and 17.4 °C. In addition, the wind speed is different in both catchments. Calluancay is very exposed to prevailing winds while Cumbe is relatively sheltered. The daily average wind speed for Calluancay and Cumbe are 4.2 (max: 11.9) and 0.9 (max: 2.6) m s⁻¹ respectively.

4.1.2 The vegetation coefficient

The time series during a wet period ranging between November 2007 and November 2008 (about a year) was used to estimate the vegetation coefficient, k_v for Calluancay. During this time period, the water content shows values greater or equal to field capacity (0.835 cm³ cm⁻³) (Table 1). So, this period is water stress-free and k_s was set to 1. For this wet period, k_v was estimated as 0.63. Similar but somewhat lower values in the range of 0.42 to 0.58 have been reported in the literature (Buytaert

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



determined in the laboratory (Fig. 3b). Critical soil moisture values between ca. 50–80 % of field capacity are reported in the literature (Seneviratne et al., 2010). However, most of them correspond to mineral soils in the context of crop water requirements and therefore those values cannot be applied for the present study. High critical soil moisture could be partially explained by the plant physiology. It is important in páramo because the tussock grasses (mainly *Calamagrostis* spp. and *Stipa* spp.) are characterized by specific adaptations to extreme conditions. For instance, the plants have scleromorphic leaves which are essential to resist intense solar radiation (Ramsay and Oxley, 1997). In addition, the plants are surrounded by dead leaves that protect the plant and reduce the water uptake. As well as in Cumbe the grazing pastures are characterized by plants of type C3 (*Pennisetum clandestinum*) which are highly resistant to drought. Therefore, the water uptake is mainly regulated by the plants during dry periods. This is clearly observed in the TDR data (Fig. 3). The time series of soil water content reveal a constant rate of water uptake during dry periods.

4.2 The actual vegetation evapotranspiration derived from soil water observations

For Calluancay, the lowest value of k_s that was observed amounts to 0.62 (11 November 2010). For the same day, in Cumbe the k_s was 0.50. In other words, the water uptake by the roots was reduced by 38 % in the páramo and by 50 % for the pasture in Cumbe. This is a clear indication of the magnitude of the 2010 drought. A similar reduction could be expected for the drought event in 2009 as the soil moisture content was even a bit lower to the registered value in 2010 (Fig. 3a). Nevertheless, in these climate conditions the vegetation was affected significantly. In addition, the probability of human-caused fires in páramo is higher during dry periods. This could affect the resilience of the soils especially during the subsequent wet period, since the vegetation is the main factor influencing the infiltration of the water.

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[⏴](#)

[⏵](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

So, for the whole period, which includes the drought of 2010, the daily average of E_a , for Calluancay and Cumbe is 1.42 (range 0.48 to 2.37) and 2.09 (range 0.99 to 3.04) mm day^{-1} respectively.

The water balance in the two experimental catchments and its components expressed as cumulative volume over the period from July 2010 until February 2012 are given in Table 1 and Fig. 4. From this analysis it is clear that the páramo vegetation and soil are more resilient to drought as compared to the lower grass vegetation and soil. Especially the recovery after the drought period is much shorter. The lower potential evapotranspiration is an important reason. Both the reference crop evapotranspiration and the vegetation coefficient are lower, so that páramo consumes less water. According to our experimental results, during wet periods the proportion of the stream discharge that can be regarded as potential water use is 54 %. In addition the rainfall is slightly higher, so the runoff coefficient in páramo during the wet period remains at 0.68 of the rainfall and is still as high as 0.50 during the dry period. For the lower Cumbe catchment these coefficients are much lower: 0.21 and 0.23 for the wet and dry periods, respectively. Although the soil characteristics are very important for sustaining the base flow the different vegetation requirements are the crucial factors that explain these differences. Furthermore, the evaporative demand is higher in Cumbe as compared with Calluancay, due to the altitude difference. As a consequence, the length of recovery period for the drought event in 2010 was three months for the organic soils while in the mineral soils it took eight months (Fig. 3).

4.3 Actual catchment evapotranspiration estimated by hydrological modelling

An initial inspection of the discharge and rainfall data revealed that the drought period of 2010 was followed by a wet period induced by a flood event on 10 April 2011. On the other hand, 2012 was relatively normal with the classic bimodal pattern (wet and dry period) (Celleri et al., 2007).

Therefore, the time periods from 16 April 2011 until 16 January 2012 and from 17 January 2012 until 16 October 2012 are used as calibration and validation period re-

spectively. These periods do not include the aforementioned extreme events. The selected periods for calibration and validation resemble the average climatic conditions of the southern Ecuadorian Andes (Buytaert et al., 2006b; Celleri et al., 2007).

The Table 3 and Fig. 5 depict the results of the implementation of the PDM model.

The performance of the model for the calibration period is good (average Nash–Sutcliffe efficiency 0.74). But, during the evaluation period the bias is high and the number of samples inside of the uncertainty bound is only 64 % for both catchments. Important is that the point measurements of the soil moisture content are in line with the catchment's average soil moisture storage calculated by the conceptual model (Fig. 5).

The calibrated maximum storage capacity in Calluancay is two times higher as compared to the value for Cumbe (Table 2). This confirms the water holding capacity of the Andosols, despite the fact that the soils are shallow. The spatial variability of the topsoil moisture storage is also high in páramo, which is congruent with the field observations carried out during soil surveys. In Cumbe the spatial variability of the topsoil moisture storage is lower. The values of the parameter b for both catchments, that reflect the spatial variability of the storage capacity, also reflect this. The lower spatial variability is probably also the reason why the simulated and observed soil moisture storage agrees better in Cumbe (Fig. 5b). These results are in line with the literature (Brocca et al., 2012). The point measurements of soil moisture can thus be confirmed as representative for the catchment's average soil moisture storage or general wetness condition.

The daily average values of E_a , as estimated by the PDM models for Calluancay and Cumbe, are 1.34 (range 0.17 to 2.79) and 1.77 (range 0.34 to 3.50) mm day^{-1} respectively. The mean values are similar to those obtained by the water balance equation and soil moisture observations. However, the range of variation is different for both methods. More variation is observed in the time series of E_a estimated by the PDM model. This is more evident in the Cumbe catchment (see Fig. 6). The PDM model does not regard a critical soil moisture value and therefore there are no constraints on

HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

The estimated coefficients k_v during the wet periods where the potential vegetation evapotranspiration is observed for both páramo's vegetation and grazing pastures were 0.63 and 0.82 respectively. These data are consistent with the literature. The k_v value is slightly higher than reported in previous studies in the case of páramo's vegetation, but obviously frequent burning and human intervention on this páramo generate more vigorous vegetation and so more demand for water. The critical soil water content for Andosols and Dystric Cambisols were 0.81 and 0.50 $\text{cm}^3 \text{cm}^{-3}$. The daily average actual evapotranspiration in páramo is low, 1.42 mm day^{-1} . While, for grazing pastures the E_a is 2.09 mm day^{-1} . From the water balance, the proportion of potential water use in the páramo of Calluancay is 54 %.

During the drought events in 2009 and 2010, the soil water content in páramo reached values never seen before. And so it was possible to establish the amount of water of the topsoil which is active during these dry periods. The reservoir can deliver a water volume equivalent to 0.24 $\text{cm}^3 \text{cm}^{-3}$ (this represents the maximum soil water content change) during extreme climate conditions such as the droughts in 2009 and 2010. In normal conditions the maximum change observed in the soil water content is no more than 0.05 $\text{cm}^3 \text{cm}^{-3}$. As consequence the real evapotranspiration can be reduced up to 38 % of its potential by stress conditions.

Thus, despite having suffered an extreme drought in 2009, the páramo soils recovered of another drought in 2010. During last period an extreme drought event was recorded in the entire Amazon River basin. In the páramo, three months of precipitations were enough to recover its normal moisture conditions. This did not occur at lower altitudes where mineral soils (Cumbe) needed about eight months in total to achieve this recovery. The combination of two factors explains the recovery of the páramo ecosystem, a big soil water storage capacity of Andosols and a low atmospheric evaporative demand due to altitude and the typical vegetation. These factors play a pivotal role in the resilience capacity to droughts especially in páramos with seasonal patterns as in Calluancay. Therefore, the páramo ecosystems have a high resilience to droughts.

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

In this context, point measurements of soil moisture in the topsoil (30 cm) were in line with the catchment's average soil moisture storage as estimated by the PDM model. The storage parameters of the PDM are also in line with field observations and literature. The E_{a^*} is however overestimated by the PDM model as compared to the water balance based on soil water measurements.

Finally, the present research has shown the value of soil water measurements at representative sites as they correspond well to the soil storage as estimated in a conceptual model. A more realistic estimation of the actual evapotranspiration can be done on the basis of the soil water content measurements. As continuous soil water data logging by TDR requires large investments the locations have to be selected with great care so that this point measurements can be considered a reliable proxy for the catchment's average soil moisture storage.

The Supplement related to this article is available online at doi:10.5194/hessd-12-11449-2015-supplement.

Acknowledgements. We thank the VLIR-IUC programme and IFS for its financial support during this research.

References

- Allen, R., Pereira, L. S., Raes, D., and Smith, M.: Crop Evapotranspiration. Guidelines for Computing Crop Water Requirements, FAO Irrigation Drainage Paper 56, FAO, Rome, 1998.
- Beven, K. and Binley, A.: The future of distributed models: model calibration and uncertainty prediction, *Hydrol. Process.*, 6, 279–298, doi:10.1002/hyp.3360060305, 1992.
- Brocca, L., Tullo, T., Melone, F., Moramarco, T., and Morbidelli, R.: Catchment scale soil moisture spatial-temporal variability, *J. Hydrol.*, 422–423, 63–75, doi:10.1016/j.jhydrol.2011.12.039, 2012.

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Buytaert, W. and Beven, K.: Models as multiple working hypotheses: hydrological simulation of tropical alpine wetlands, *Hydrol. Process.*, 25, 1784–1799, doi:10.1002/hyp.7936, 2011.
- Buytaert, W. and De Bièvre, B.: Water for cities: the impact of climate change and demographic growth in the tropical Andes, *Water Resour. Res.*, 48, W08503, doi:10.1029/2011WR011755, 2012.
- Buytaert, W., Wyseure, G., De Bièvre, B., and Deckers, J.: The effect of land-use changes on the hydrological behaviour of Histic Andosols in south Ecuador, *Hydrol. Process.*, 19, 3985–3997, doi:10.1002/hyp.5867, 2005.
- Buytaert, W., Célleri, R., De Bièvre, B., Cisneros, F., Wyseure, G., Deckers, J., and Hofstede, R.: Human impact on the hydrology of the Andean páramos, *Earth-Sci. Rev.*, 79, 53–72, doi:10.1016/j.earscirev.2006.06.002, 2006a.
- Buytaert, W., Celleri, R., Willems, P., Bièvre, B. De and Wyseure, G.: Spatial and temporal rainfall variability in mountainous areas: A case study from the south Ecuadorian Andes, *J. Hydrol.*, 329, 413–421, doi:10.1016/j.jhydrol.2006.02.031, 2006b.
- Buytaert, W., Iñiguez, V., Celleri, R., De Bièvre, B., Wyseure, G., and Deckers, J.: Analysis of the water balance of small páramo catchments in south Ecuador, in: *Environmental Role of Wetlands in Headwaters SE-24*, vol. 63, edited by: Krecek, J. and Haigh, M., Springer Netherlands, Dordrecht, the Netherlands, 271–281, 2006c.
- Buytaert, W., Deckers, J., and Wyseure, G.: Regional variability of volcanic ash soils in south Ecuador: the relation with parent material, climate and land use, *Catena*, 70, 143–154, doi:10.1016/j.catena.2006.08.003, 2007a.
- Buytaert, W., Iñiguez, V., and De Bièvre, B.: The effects of afforestation and cultivation on water yield in the Andean páramo, *Forest Ecol. Manage.*, 251, 22–30, doi:10.1016/j.foreco.2007.06.035, 2007b.
- Buytaert, W., Vuille, M., Dewulf, A., Urrutia, R., Karmalkar, A., and Célleri, R.: Uncertainties in climate change projections and regional downscaling in the tropical Andes: implications for water resources management, *Hydrol. Earth Syst. Sci.*, 14, 1247–1258, doi:10.5194/hess-14-1247-2010, 2010.
- Buytaert, W., Cuesta-Camacho, F., and Tobón, C.: Potential impacts of climate change on the environmental services of humid tropical alpine regions, *Global Ecol. Biogeogr.*, 20, 19–33, doi:10.1111/j.1466-8238.2010.00585.x, 2011.
- Celleri, R., Willems, P., Buytaert, W., and Feyen, J.: Space–time rainfall variability in the Paute basin, Ecuadorian Andes, *Hydrol. Process.*, 21, 3316–3327, doi:10.1002/hyp.6575, 2007.

HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Dercon, G., Bossuyt, B., De Bièvre, B., Cisneros, F., and Deckers, J.: Zonificación agroecológica del Austro Ecuatoriano, U Ediciones, Cuenca, Ecuador, 1998.
- Dercon, G., Govers, G., Poesen, J., Sánchez, H., Rombaut, K., Vandembroeck, E., Loaiza, G., and Deckers, J.: Animal-powered tillage erosion assessment in the southern Andes region of Ecuador, *Geomorphology*, 87, 4–15, doi:10.1016/j.geomorph.2006.06.045, 2007.
- Duan, Q., Sorooshian, S., and Gupta, V.: Effective and efficient global optimization for conceptual rainfall–runoff models, *Water Resour. Res.*, 28, 1015–1031, doi:10.1029/91WR02985, 1992.
- FAO/ISRIC/ISSS: World Reference Base for Soil Resources, No. 84 in World Soil Resources Reports, FAO, Rome, 1998.
- Farley, K. A., Kelly, E. F., and Hofstede, R. G. M.: Soil organic carbon and water retention after conversion of grasslands to pine plantations in the Ecuadorian Andes, *Ecosystems*, 7, 729–739, doi:10.1007/s10021-004-0047-5, 2004.
- Garcia, M., Raes, D., Allen, R., and Herbas, C.: Dynamics of reference evapotranspiration in the Bolivian highlands (Altiplano), *Agr. Forest Meteorol.*, 125, 67–82, doi:10.1016/j.agrformet.2004.03.005, 2004.
- Hargreaves, G. H. and Samani, Z. A.: Reference crop evapotranspiration from temperature, *Appl. Eng. Agric.*, 1, 96–99, 1985.
- Hofstede, R. G. M., Groenendijk, J. P., Coppus, R., Fehse, J. C., and Sevink, J.: Impact of pine plantations on soils and vegetation in the Ecuadorian High Andes, *Mt. Res. Dev.*, 22, 159–167, doi:10.1659/0276-4741(2002)022[0159:IOPPOS]2.0.CO;2, 2002.
- Hofstede, R., Segarra, P., and Mena, P.: Los páramos del mundo, Global Peatland Initiative/NC-IUCN/EcoCiencia, Quito, 2003.
- Hungerbühler, D., Steinmann, M., Winkler, W., Seward, D., Egüez, A., Peterson, D. E., Helg, U., and Hammer, C.: Neogene stratigraphy and Andean geodynamics of southern Ecuador, *Earth-Sci. Rev.*, 57, 75–124, doi:10.1016/S0012-8252(01)00071-X, 2002.
- Kilpatrick, F. and Schneider, V.: Use of Flumes in Measuring Discharge, US Geological Survey Techniques of Water Resources Investigations, US Geological Survey, Washington, USA, 1983.
- Klemeš, V.: Operational testing of hydrological simulation models, *Hydrolog. Sci. J.*, 31, 13–24, doi:10.1080/02626668609491024, 1986.
- Koch, K., Wenninger, J., Uhlenbrook, S., and Bonell, M.: Joint interpretation of hydrological and geophysical data: electrical resistivity tomography results from a process hydrological

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

research site in the Black Forest Mountains, Germany, *Hydrol. Process.*, 23, 1501–1513, doi:10.1002/hyp.7275, 2009.

Lewis, S. L., Brando, P. M., Phillips, O. L., van der Heijden, G. M. F., and Nepstad, D.: The 2010 Amazon Drought, *Science*, 331, 554, doi:10.1126/science.1200807, 2011.

Luteyn, J. L.: *Páramos: a Checklist of Plant Diversity, Geographical Distribution, and Botanical Literature*, The New York Botanical Garden Press, New York, 1999.

Marengo, J. A., Nobre, C. A., Tomasella, J., Oyama, M. D., de Oliveira, G. S., de Oliveira, R., Camargo, H., Alves, L. M., and Brown, I. F.: The drought of Amazonia in 2005, *J. Climate*, 21, 495–516, doi:10.1175/2007JCLI1600.1, 2008.

Marengo, J. A., Tomasella, J., Alves, L. M., Soares, W. R., and Rodriguez, D. A.: The drought of 2010 in the context of historical droughts in the Amazon region, *Geophys. Res. Lett.*, 38, 1–5, doi:10.1029/2011GL047436, 2011.

Moore, R. J.: The probability-distributed principle and runoff production at point and basin scales, *Hydrolog. Sci. J.*, 30, 273–297, doi:10.1080/02626668509490989, 1985.

Moore, R. J. and Clarke, R. T.: A distribution function approach to rainfall runoff modeling, *Water Resour. Res.*, 17, 1367–1382, doi:10.1029/WR017i005p01367, 1981.

Mora, D. E., Campozano, L., Cisneros, F., Wyseure, G., and Willems, P.: Climate changes of hydrometeorological and hydrological extremes in the Paute basin, Ecuadorian Andes, *Hydrol. Earth Syst. Sci.*, 18, 631–648, doi:10.5194/hess-18-631-2014, 2014.

Pebesma, E. J.: Multivariable geostatistics in S: the gstat package, *Comput. Geosci.*, 30, 683–691, doi:10.1016/j.cageo.2004.03.012, 2004.

Phillips, O. L., Aragão, L. E. O. C., Lewis, S. L., Fisher, J. B., Lloyd, J., López-González, G., Malhi, Y., Monteagudo, A., Peacock, J., Quesada, C. A., van der Heijden, G., Almeida, S., Amaral, I., Arroyo, L., Aymard, G., Baker, T. R., Bánki, O., Blanc, L., Bonal, D., Brando, P., Chave, J., de Oliveira, A. C. A., Cardozo, N. D., Czimczik, C. I., Feldpausch, T. R., Freitas, M. A., Gloor, E., Higuchi, N., Jiménez, E., Lloyd, G., Meir, P., Mendoza, C., Morel, A., Neill, D. A., Nepstad, D., Patiño, S., Peñuela, M. C., Prieto, A., Ramírez, F., Schwarz, M., Silva, J., Silveira, M., Thomas, A. S., Ter Steege, H., Stropp, J., Vásquez, R., Zelazowski, P., Alvarez Dávila, E., Andelman, S., Andrade, A., Chao, K.-J., Erwin, T., Di Fiore, A., Honorio, C. E., Keeling, H., Killeen, T. J., Laurance, W. F., Peña Cruz, A., Pitman, N. C. A., Núñez Vargas, P., Ramírez-Angulo, H., Rudas, A., Salamão, R., Silva, N., Terborgh, J., and Torres-Lezama, A.: Drought sensitivity of the Amazon rainforest, *Science*, 323, 1344–1347, doi:10.1126/science.1164033, 2009.

HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Podwojewski, P., Poulénard, J., Zambrana, T., and Hofstede, R.: Overgrazing effects on vegetation cover and properties of volcanic ash soil in the páramo of Llangahua and La Esperanza (Tungurahua, Ecuador), *Management*, 18, 45–55, doi:10.1111/j.1475-2743.2002.tb00049.x, 2002.

Ramsay, P. M. and Oxley, E. R. B.: The growth form composition of plant communities in the ecuadorian páramos, *Plant Ecol.*, 131, 173–192, doi:10.1023/A:1009796224479, 1997.

Romano, N.: Soil moisture at local scale: measurements and simulations, *J. Hydrol.*, 516, 6–20, doi:10.1016/j.jhydrol.2014.01.026, 2014.

Schneider, P., Vogt, T., Schirmer, M., Doetsch, J., Linde, N., Pasquale, N., Perona, P., and Cirpka, O. A.: Towards improved instrumentation for assessing river-groundwater interactions in a restored river corridor, *Hydrol. Earth Syst. Sci.*, 15, 2531–2549, doi:10.5194/hess-15-2531-2011, 2011.

Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B., and Teuling, A. J.: Investigating soil moisture-climate interactions in a changing climate: a review, *Earth-Sci. Rev.*, 99, 125–161, doi:10.1016/j.earscirev.2010.02.004, 2010.

Stocki, R.: A method to improve design reliability using optimal Latin hypercube sampling, *Comput. Assist. Mech. Eng. Sci.*, 12, 393–411, 2005.

Vanacker, V., Molina, A., Govers, G., Poesen, J., and Deckers, J.: Spatial variation of suspended sediment concentrations in a tropical Andean river system: the Paute River, southern Ecuador, *Geomorphology*, 87, 53–67, doi:10.1016/j.geomorph.2006.06.042, 2007.

van Genuchten, M. T.: A closed-form equation for predicting the hydraulic conductivity of unsaturated soils, *Soil Sci. Soc. Am. J.*, 44, 892–898, doi:10.2136/sssaj1980.03615995004400050002x, 1980.

Van Loon, A. F.: Hydrological drought explained, *Wiley Interdisciplinary Reviews: Water*, 2, 359–392, doi:10.1002/wat2.1085, 2015.

Viviroli, D., Archer, D. R., Buytaert, W., Fowler, H. J., Greenwood, G. B., Hamlet, A. F., Huang, Y., Koboltschnig, G., Litaor, M. I., López-Moreno, J. I., Lorentz, S., Schädler, B., Schreier, H., Schwaiger, K., Vuille, M., and Woods, R.: Climate change and mountain water resources: overview and recommendations for research, management and policy, *Hydrol. Earth Syst. Sci.*, 15, 471–504, doi:10.5194/hess-15-471-2011, 2011.

Vuille, M., Bradley, R. S., and Keimig, F.: Climate variability in the Andes of Ecuador and its relation to Tropical Pacific and Atlantic Sea surface temperature anomalies, *J. Climate*, 13, 2520–2535, doi:10.1175/1520-0442(2000)013<2520:CVITAO>2.0.CO;2, 2000.

Wagener, T., Boyle, D. P., Lees, M. J., Wheeler, H. S., Gupta, H. V., and Sorooshian, S.: A framework for development and application of hydrological models, *Hydrol. Earth Syst. Sci.*, 5, 13–26, doi:10.5194/hess-5-13-2001, 2001.

HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

Table 2. The calibrated parameters of the PDM model.

Parameters	Description	Feasible range	Calluancay	Cumbe
C_{\max}	Maximum storage capacity	30–120 [mm]	101.2	44.2
b	Degree of spatial variability of the storage capacity	0.1–2.0 [–]	1.82	0.21
f_{rt}	Fast routing store residence time	1–2 [days]	1.7	1.1
s_{rt}	Slow routing store residence time	10–50 [days]	14.3	32.5
$\%(q)$	Percentage flow through fast flow	0.25–0.75 [–]	0.58	0.42

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.

Table 3. The Nash and Sutcliffe efficiencies and the bias for the PDM models*.

Catchment	Total period		Period 1		Period 2		Split sample		
	NS (–)	Bias (%)	NS (–)	Bias (%)	NS (–)	Bias (%)	NS (–)	Bias (%)	AC (%)
Calluancay	0.73	–9.6	0.67	–10.0	0.83	–19.2	0.74	–20.9	65
Cumbe	0.81	–3.8	0.82	–3.0	0.78	–0.3	0.66	–17.5	64

* Total period: 16 Jul 2010–15 Nov 2012, Period 1: 16 Apr 2011–16 Jan 2012, Period 2: 17 Jan 2012–16 Oct 2012. The parameter set calibrated for period 1 is used for validation in the period 2 (split sample test). NS is the Nash–Sutcliffe efficiency and AC is the percentage of data (discharge) that falls inside the uncertainty bands.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.




Figure 1. The study area.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

HESSD

12, 11449–11484, 2015

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.




Figure 2. The potential evapotranspiration E_p for Calluancay (black) and Cumbe (grey).

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.




Figure 3. The soil water content data for Calluancay (a) and Cumbe (b). The drought periods are shaded in grey. The blue lines show the soil water content simulated by means of the soil water balance in the root zone for each catchment during the period from on 16 July 2010 up to on 1 February 2012. In Supplement there are dotted plots with the parameters optimized during the water balance.

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.




Figure 4. Water balance components for **(a)** Calluancay and **(b)** Cumbe. The curves in Calluancay show a non-linear behaviour and so suggest a seasonality for this páramo area. This climate pattern is enhanced by the occurrence of drought events. A high evapotranspiration is revealed in Cumbe. Most of this E_a can be attributed to the irrigation systems which are operational during dry periods.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.




Figure 5. Representative sample of rainfall (top), runoff (middle) and soil moisture (bottom) time series. Uncertainty bounds are also included in the graphs. The scaled soil moisture storage in the root zone is shown in the bottom inset in the plot in solid and dashed black lines for measured and modelled respectively.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Analysis of the drought resilience of Andosols on southern Ecuadorian Andean páramos

V. Iñiguez et al.




Figure 6. The actual evapotranspiration (E_a) using the water balance approach (WBA) and the PDM model. **(a)** Calluancay and **(b)** Cumbe.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)