

Evaluation of a multi-satellite soil moisture product and the Community Land Model 4.5 simulation in China

Binghao Jia¹, Jianguo Liu^{1,2} and Zhenghui Xie¹

[1] {State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics (LASG), Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China}

[2] {High Performance Computing Center, Department of Mathematics and Applied Mathematics, Huaihua University, Huaihua, Hunan, China}

Correspondence to: Jianguo Liu (jgliu@mail.iap.ac.cn)

Submitted to *Hydrology and Earth System Sciences*

5 May 2015 – Revised: 21 August 2015

Abstract

Twenty years of in situ soil moisture data from 306 stations located in China were used to perform an evaluation of two surface soil moisture datasets: (1) a microwave-based multi-satellite product (ESA CCI SM) and (2) soil moisture estimations from the Community Land Model 4.5 (CLM4.5), forced by observation-based atmospheric forcing data. Both soil moisture products generally showed a good agreement with in situ observations, with unbiased root mean square differences (ubRMSD) of $0.05 \text{ m}^3 \text{ m}^{-3}$. The average Spearman rank correlation coefficient (R_{sp}) between the ESA CCI SM product and all in situ observations was 0.37. In contrast, the CLM4.5 model produced better temporal variation of surface soil moisture ($R_{sp} = 0.42$) than the ESA CCI SM product, but showed larger ubRMSD in southwestern China, which may have been related to inaccurate precipitation data. The ESA CCI SM product is more likely to be superior to the CLM4.5 model in semi-arid regions, mainly because of the accurate data retrievals and high observation density, but inferior over areas covered by dense vegetation. Furthermore, the ESA CCI SM product showed a stable to

binghaojia 15/7/29 8:33 PM

已删除: more than 300

Binghao Jia 15/8/20 6:12 PM

已删除: ar

binghaojia 15/6/24 3:09 PM

已删除: ECV-SM

binghaojia 15/6/24 2:41 PM

已删除: the land surface model simulation

Binghao Jia 15/8/20 6:13 PM

已删除: a land surface model,

Binghao Jia 15/8/20 6:13 PM

已删除: an

Binghao Jia 15/8/21 8:34 AM

已删除: values

Binghao Jia 15/8/21 8:34 AM

已删除: -

binghaojia 15/6/29 2:23 PM

已删除: ECV-S

Binghao Jia 15/8/21 8:34 AM

已删除: i

binghaojia 15/6/29 2:23 PM

已删除: M product has a low bias, with a root mean square difference (RMSD) of $0.075 \text{ m}^3 \text{ m}^{-3}$, but shows a weak correlation with in situ observations ($R = 0.41$)

binghaojia 15/7/29 8:52 PM

已删除: simulation, forced by an ... (1)

Binghao Jia 15/8/21 8:35 AM

已删除: s

binghaojia 15/7/29 8:52 PM

已删除: 52

Binghao Jia 15/8/21 8:35 AM

已删除: s

binghaojia 15/7/29 8:53 PM

已删除: a clear overestimation (bias ... (2)

Binghao Jia 15/8/21 8:36 AM

已删除: ($0.09 \text{ m}^3 \text{ m}^{-3}$)

binghaojia 15/7/29 8:54 PM

已删除: , especially

Binghao Jia 15/8/21 8:36 AM

已删除:

binghaojia 15/7/29 8:54 PM

已删除: eastern

binghaojia 15/7/29 8:55 PM

已删除: caused by

binghaojia 15/7/29 8:54 PM

已删除: descriptions of soil characteristics

binghaojia 15/6/29 2:23 PM

已删除: ECV-SM

Binghao Jia 15/8/21 8:38 AM

已删除: it

Binghao Jia 15/8/21 8:38 AM

已删除: s

60 | slightly better performance in China over time, except for a decline in performance during
61 | 2007–2010, when different data for the satellite product were blended. Results from this
62 | study can provide comprehensive insight into the performances of the two soil moisture
63 | datasets in China, which will facilitate improvements in merging algorithms or model
64 | simulations and for applications in soil moisture data assimilation.

65 | **Key words:** soil moisture, remote sensing, CLM4.5, essential climate variable,

66 | **1. Introduction**

67 | Soil moisture is a key variable in hydrological, climatological, biological and ecological
68 | processes. It is central to land–atmosphere interactions because of its control on the
69 | partitioning of water and energy fluxes at the Earth's surface (Dai et al., 2004). Soil moisture
70 | also affects the seasonal and inter-annual dynamics of vegetation, which is an essential
71 | component of the coupled hydrological and carbon cycles (Ciais et al., 2005). Many studies
72 | have been conducted to obtain estimates of soil moisture using remote sensing techniques
73 | (Njoku et al., 2003; Owe et al., 2008; Kerr et al., 2012). Land surface modeling (Dirmeyer et
74 | al., 2006; Wang et al., 2011; Liu and Xie, 2013) or a combination of both through a land data
75 | assimilation system (e.g., Dharssi et al., 2011, de Rosnay et al., 2013).

76 | Spaceborne microwave instruments can provide quantitative information about surface
77 | soil water content (Schmugge, 1983), particularly in the low-frequency microwave region
78 | from 1 to 10 GHz (Albergel et al., 2012). Several microwave-based soil moisture datasets
79 | have been generated using satellite data retrievals from active microwave sensors, e.g. the
80 | European Remote Sensing Satellites 1 and 2 Active Microwave Instrument Wind
81 | Scatterometer (AMI-WS; Scipal et al., 2002) and Advanced SCATterometer (ASCAT)
82 | onboard the Meteorological Operational satellite program (MetOp; Bartalis et al., 2007), and
83 | from passive microwave sensors, including the Scanning Multichannel Microwave
84 | Radiometer (SMMR; Owe et al., 2008), the Special Sensor Microwave Imager (SSM/I) of the

Binghao Jia 15/8/21 8:40 AM

已删除: increasing

Binghao Jia 15/8/21 8:41 AM

已删除: decrease

Binghao Jia 15/8/21 8:42 AM

已删除: the

Binghao Jia 15/8/21 8:43 AM

已删除: blending period

Binghao Jia 15/8/21 8:44 AM

已删除: be useful for their

Binghao Jia 15/8/21 8:45 AM

已删除: , in situ measurements

Binghao Jia 15/8/21 8:45 AM

已删除: A significant number of

Binghao Jia 15/7/1 10:34 AM

已删除: and

Binghao Jia 15/8/21 8:46 AM

已删除: are able to

94 Defense Meteorological Satellite Program (DMSP; Owe et al., 2008), the Tropical Rainfall
95 Measuring Mission microwave imager (TMI; Jackson and Hsu, 2001; Gao et al., 2006; Owe
96 et al., 2008), and more recently the Advanced Microwave Scanning Radiometer–Earth
97 observing system (AMSR-E) onboard the Aqua satellite (Njoku et al., 2003; Owe et al.,
98 2008). The AMSR-E radiometer was switched off in October 2011 because of rotation
99 problems with its antenna; however, the AMSR2, launched in May 2012 onboard the Global
100 Change Observation Mission 1–Water (GCOM–W1), was intended to extend [its](#) valuable
101 heritage [\(Parinussa et al., 2015\)](#). Even though none of these sensors were specifically
102 designed for measuring soil moisture, good correspondences have been found between the
103 individual datasets and ground-based observations taken over a large variety of
104 environmental conditions (Albergel et al., 2009, 2012; Draper et al., 2009; Gruhier et al.,
105 2010; Brocca et al., 2011).

106 However, none of the individual microwave products cover the decadal timescales
107 required to be considered for [use in](#) climate applications. Recently, a multi-satellite [soil](#)
108 moisture dataset [\(ESA CCI SM\)](#) [spanning](#) over thirty years was constructed by merging two
109 active and six passive microwave products (Liu et al., 2011, 2012). The [combined](#) product,
110 which was initially developed under the European Space Agency (ESA) Water Cycle
111 Multi-Mission Observation Strategy (WACMOS) project, is now being extended and
112 improved within the Climate Change Initiative (CCI) (<http://www.esa-soilmoisture-cci.org/>;
113 Wagner et al., 2012). A few studies have [evaluated](#) the [ESA CCI SM product](#) using in situ
114 observations. Liu et al. (2011, 2012) indicated that the merged dataset had a similar accuracy
115 to that of the best input product but with an increased temporal sampling density. Albergel et
116 al. (2013a) found that the [ESA CCI SM dataset](#) agreed well with in situ measurements
117 between 2007 and 2010 for 196 sites from five networks across the world, but that its
118 performance over most networks remained poorer than that of the [most recent](#) reanalysis

Binghao Jia 15/7/1 10:42 AM

已删除: the

Binghao Jia 15/7/1 10:43 AM

已删除: of AMSR-E and provide improved spatial resolution because of its larger reflector

Binghao Jia 15/8/21 8:49 AM

已删除: the

binghaojia 15/7/20 10:11 PM

已删除: Essential Climate Variable

binghaojia 15/6/24 3:50 PM

已删除: ECV-SM

Binghao Jia 15/8/21 8:50 AM

已删除: with

Binghao Jia 15/8/21 8:50 AM

已删除: is

Binghao Jia 15/8/21 8:52 AM

已删除: considered the evaluation of

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

binghaojia 15/6/29 2:25 PM

已删除: ECV-SM

130 | products. Dorigo et al. (2015) provided a more in-depth evaluation; they used 596 stations
131 | from 28 historical and active monitoring networks worldwide. Whilst the performance of the
132 | ESA CCI SM dataset appeared to be relatively stable over time, large discrepancies were
133 | observed among different networks. In addition, the ESA CCI SM can also capture long-term
134 | systematic changes in trends of ground-based observations (Dorigo et al., 2012; Albergel et
135 | al., 2013b), which suggests that it has a large potential for climate trend assessments (Loew et
136 | al., 2013).

137 | China has the third largest land area and diverse climates and biomes. Previous studies
138 | have used only 34 sites for the period 1981–2000 across China and only 20 sites for the
139 | period 2008–2010 from the Maqu network in northwest China to investigate the performance
140 | of the ESA CCI SM product (Albergel et al., 2013a; Dorigo et al., 2015). Such sparse
141 | observations clearly affect the evaluation results, leading to large uncertainties. Recently,
142 | Chinese soil moisture observations for 306 sites from 1993 were updated by the China
143 | Meteorological Administration (CMA) National Meteorological Information Center (NMIC).
144 | These data have been extensively used to investigate the variations of soil moisture and
145 | evaluate the land surface modeling (Li et al, 2005; Wang and Zeng, 2011; Liu and Xie,
146 | 2013).

147 | Land surface modeling is another strategy to produce large-scale surface and root zone
148 | soil moisture estimations. When forced by high quality atmospheric forcing data, this strategy
149 | has proved to be an effective tool to complement the commonly use of in situ measurements
150 | and for the evaluation of satellite retrievals at both regional and global scales (Albergel et al.,
151 | 2010, 2012). The ESA CCI SM product has also demonstrated a potential for evaluating
152 | climate model performances (Loew et al, 2013; Szczypa et al., 2014). As one of the
153 | state-of-art land surface models, the Community Land Model version 4.5 (CLM4.5) from the
154 | National Center for Atmospheric Research (NCAR) was released in 2013 (Oleson et al.,

Binghao Jia 15/8/21 8:54 AM
已删除: soil moisture
binghaojia 15/6/29 4:26 PM
已删除: 2014
binghaojia 15/6/29 2:26 PM
已删除: ECV-SM
binghaojia 15/6/29 2:26 PM
已删除: ECV-SM

binghaojia 15/6/29 2:26 PM
已删除: ECV-SM
binghaojia 15/6/29 4:23 PM
已删除: 2014
Binghao Jia 15/8/21 8:55 AM
已删除: will
binghaojia 15/8/11 9:59 AM
已删除: 778

Binghao Jia 15/8/21 8:57 AM
已删除: with
Binghao Jia 15/7/1 10:55 AM
已删除: d
Binghao Jia 15/8/21 9:00 AM
已删除: the
binghaojia 15/6/29 2:26 PM
已删除: ECV-SM
Binghao Jia 15/8/21 9:01 AM
已删除: shown a

168 2013). To our knowledge, few studies [have focused](#) on the performance of CLM4.5 soil
169 moisture simulations in China. [Lai et al. \(2015\) validated the temporal variation of soil](#)
170 [moisture simulated from the CLM4.0, a previous version of CLM4.5, but used only 30 sites](#)
171 [to cover China for the period 1981–1999; they also compared the spatial variation of soil](#)
172 [moisture in China using the ESA CCI SM product. However, the performance of the ESA](#)
173 [CCI SM product was not discussed in their work.](#)

174 In this study, we [conducted](#) an in-depth evaluation of the [ESA CCI SM](#) product and
175 CLM4.5 simulation in China using ground-based observations from [306](#) sites. We
176 [investigated](#) their performances over [various](#) sub-regions under different climate conditions.
177 This in-depth evaluation [provided](#) a better understanding of the quality of both soil moisture
178 products and their potential problems [than was hereafter available, and](#) can be used to
179 improve their accuracy. The soil moisture datasets used in this study and the methodology for
180 their evaluation are described in Section 2. The results and discussion are then presented in
181 Section 3 and Section 4, respectively. Finally, [our conclusions are](#) given in Section 5.

182 2. Material and methods

183 2.1 Community Land Model (CLM4.5)

184 The NCAR/CLM is a community-developed model for simulating land surface
185 processes, such as water, energy, and carbon fluxes. [The CLM4.5 version](#) is the latest of the
186 CLM family of models (Oleson et al., 2013). It contains several notable improvements over
187 previous releases, including the decrease of biases associated with the modeled terrestrial
188 carbon cycle and modifications of canopy and hydrology processes (Oleson et al. 2013). In
189 CLM4.5, spatial land surface heterogeneity is represented as a nested subgrid hierarchy in
190 which grid cells are composed of multiple land units, snow/soil columns, and plant functional
191 types (PFTs). [The model](#) has one vegetation layer, fifteen layers for soil and up to five layers
192 for snow, depending on snow depth. The soil depths for [the uppermost](#) five layers are 1.75,

Binghao Jia 15/8/21 9:04 AM

已删除: will provide

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

Binghao Jia 15/8/6 10:27 AM

已删除: 778

Binghao Jia 15/8/21 9:05 AM

已删除: will also

Binghao Jia 15/8/21 9:05 AM

已删除: the

Binghao Jia 15/8/6 10:28 AM

已删除: , including bias characteristics and temporal variations

Binghao Jia 15/8/21 9:06 AM

已删除: is expected to

Binghao Jia 15/8/21 9:07 AM

已删除: which

Binghao Jia 15/8/21 9:07 AM

已删除: the

Binghao Jia 15/8/21 9:07 AM

已删除: is

Binghao Jia 15/8/21 9:08 AM

已删除: version

Binghao Jia 15/8/21 9:09 AM

已删除: l

Binghao Jia 15/8/21 9:10 AM

已删除: CLM4.5

Binghao Jia 15/8/21 9:11 AM

已删除: top

Binghao Jia 15/8/21 9:11 AM

已删除: cm

209 4.51, 9.06, 16.55, and 28.91 cm. For soil points, temperature calculations are performed over
210 all layers, but hydrology calculations are performed over the top 10 layers only; the bottom
211 five layers are classified as bedrock. A detailed description of the physical processes included
212 within CLM4.5 can be found in Oleson et al. (2013). It should be noted that although
213 CLM4.5 includes the option to be operated with a dynamic vegetation or prognostic
214 carbon-nitrogen model, in our study, CLM4.5 was used with prescribed satellite-based
215 phenology, taken from the Moderate Resolution Imaging Spectroradiometer (MODIS;
216 Lawrence and Chase, 2007).

217 In this study, CLM4.5 was forced by a 34-yr (1979–2012) observation-based
218 atmospheric forcing dataset from the Institute of Tibetan Plateau Research, Chinese Academy
219 of Sciences (hereafter ITP), with a spatial resolution of $0.1^{\circ} \times 0.1^{\circ}$ at a three-hourly temporal
220 resolution over China (15–55 °N, 70–140 °E). This dataset was constructed by merging the
221 observations from 740 operational stations of the CMA with the corresponding
222 meteorological forcing dataset from the Global Land Data Assimilation System (Rodell et al.,
223 2004) to produce near-surface air temperature, pressure, wind speed and specific humidity
224 fields. It combined three precipitation datasets, including ground-based observations and two
225 satellite retrieval products (Chen et al., 2011), to determine the precipitation field. It also
226 corrected the Global Energy and Water Cycle Experiment–Surface Radiation Budget (Pinker
227 and Laszlo, 1992) with reference to radiation estimates (Yang et al., 2010) to ascertain the
228 incident shortwave radiation fields. Chen et al. (2011) demonstrated that simulations driven
229 by the ITP forcing data improve land surface temperature modeling for dry land in China.
230 The soil moisture simulations of CLM3.5, an old version of CLM4.5, forced by four different
231 atmospheric forcing datasets were compared against a common set of in situ observations and
232 results showed that, over most regions of China, the soil moisture estimations forced by the

Binghao Jia 15/8/21 9:11 AM
已删除: cm
Binghao Jia 15/8/21 9:11 AM
已删除: cm
Binghao Jia 15/8/21 9:11 AM
已删除: cm
Binghao Jia 15/8/21 9:12 AM
已删除: while
Binghao Jia 15/8/21 9:12 AM
已删除: ten
Binghao Jia 15/8/21 9:12 AM
已删除: being
Binghao Jia 15/8/21 9:13 AM
已删除: run
Binghao Jia 15/8/21 9:12 AM
已删除: al
Binghao Jia 15/8/21 9:13 AM
已删除: run using
Binghao Jia 15/7/1 11:07 AM
已删除: (Lawrence and Chase, 2007)
Binghao Jia 15/8/21 9:14 AM
已删除: year
Binghao Jia 15/8/21 9:14 AM
已删除:
Binghao Jia 15/8/21 9:14 AM
已删除:

Binghao Jia 15/8/5 11:35 AM
已删除: Liu and Xie (2013)
Binghao Jia 15/8/5 11:34 AM
已删除: found
Binghao Jia 15/8/5 11:26 AM
已删除: of the CLM3.5

249 ITP forcing dataset had closer correlations with ground-based observations than did the three
250 other simulations (Liu and Xie, 2013).

Binghao Jia 15/8/5 11:34 AM
已删除: ve...closer correlations with ... [3]

251 2.2 ESA CCI SM data

binghaojia 15/7/20 10:08 PM
已删除: ECV ...SA CCI SMsoil mc ... [4]

252 In response to the Global Climate Observing System endorsement of soil moisture as an
253 essential climate variable, the ESA-WACMOS and CCI projects have supported the
254 generation of the ESA CCI SM product by merging multiple microwave-based soil moisture
255 products (Wagner et al., 2012), including passive data derived from SMMR, SSM/I, TMI,
256 and AMSR-E and active data from the ERS and ASCAT (Liu et al. 2011, 2012). The ESA
257 CCI SM version 2.0 (v2.0), released by the Vienna University of Technology in July 2014,
258 was used in this study. Compared to previous versions, the merging schemes and procedures
259 for ESA CCI SM v2.0 has been improved. Furthermore, the dataset was extended to the year
260 2013 by including the WindSat and AMSR2 data (Parinussa et al., 2012, 2015). Initially, the

binghaojia 15/7/20 10:10 PM
已删除: ECV ... the ESA-WACMOS ... [5]

261 ESA CCI SM data were separated into two homogenized products (one for active and one for
262 passive data); and then merged into a single active-passive product according to their relative
263 sensitivity to vegetation density (Liu et al, 2011, 2012; Wagner et al., 2012). The ESA CCI
264 SM has a spatial resolution of $0.25^\circ \times 0.25^\circ$ (unit: $\text{m}^3 \text{m}^{-3}$) and a daily time-step centered at
265 0:00 UTC, although the actual observation time corresponds to that of the input products at a
266 specific time (Liu et al., 2012). Quality “flags” of the input products were transferred to ESA
267 CCI SM to mask pixels affected by snow coverage, temperature below 0°C , dense vegetation,
268 and pixels where the retrieval of soil moisture data failed (Dorigo et al., 2015).

Binghao Jia 15/8/21 9:18 AM
已删除: it

binghaojia 15/6/29 2:26 PM
已删除: ECV-SM

Binghao Jia 15/8/21 9:18 AM
已删除: ... (one for active and one f ... [6]

binghaojia 15/6/29 2:26 PM
已删除: ECV-SM

Binghao Jia 15/8/21 9:19 AM
已删除: ... $\times 0.25$... (unit: $\text{m}^3 \text{m}^{-3}$) ... [7]

binghaojia 15/8/11 10:47 AM
已删除: These data are in volumetric ($\text{m}^3 \text{m}^{-3}$) units and q...uality “flags” of th ... [8]

269 2.3 In situ measurements in China

270 This study made use of in situ soil moisture measurements from agricultural
271 meteorological stations across mainland China, collected by the CMA-NMIC, to evaluate the
272 ESA CCI SM and CLM4.5. The original data for 20 years (1993–2012) from 778 stations
273 were obtained every 10 days (i.e., on the 8th, 18th and 28th day of every month) at soil

Binghao Jia 15/8/21 9:21 AM
已删除: kes

binghaojia 15/6/29 2:26 PM
已删除: ECV-SM

316 | depths of 0–10, 10–20, 20–50, 50–70, and 70–100 cm, respectively. No measurements were
317 | recorded in frozen soil. Soil water content was measured using the gravimetric technique and
318 | originally recorded as mass percentage. It was converted to volumetric soil moisture using
319 | field capacity and soil bulk density observations (Liu and Xie, 2013). This soil moisture
320 | observation dataset has been widely used to study temporal variations in soil moisture and
321 | evaluate land surface model simulations in China (Li et al., 2005; Wang and Zeng, 2011; Liu
322 | and Xie, 2013), and is constantly updated. However, not all datasets are suitable for the
323 | evaluation of remote sensing products and model simulations. In this study, we used a simple
324 | quality control procedure (Wang and Zeng, 2011; Liu and Xie, 2013) on the updated soil
325 | moisture observations in terms of observation frequency; specifically, the ratio of valid
326 | measurements during the period from March to October (1993–2012) was required to be
327 | greater than 50%. It is noted that if more than one in situ station remained in a single 0.25°
328 | grid box, only one of them was included. The correlation check method was adopted to
329 | remove the redundant sites (Dorigo et al., 2015); in this procedure, the correlation between
330 | the in situ measurements and ESA CCI SM product and that between the in situ
331 | measurements and CLM4.5 were first calculated, respectively, and their average value was
332 | compared for each site, after which the site with the highest average correlation was selected.
333 | Finally, the monthly soil water content values at depths of 0–10 cm at 306 stations were used
334 | in this study to evaluate the remotely sensed and simulated values. These stations were
335 | grouped into eight sub-regions on the basis of the spatial patterns of the centers of dryness
336 | and wetness throughout China, based on Zhu (2003) and Liu and Xie (2013), and are defined
337 | in Table 1. Figure 1 shows the eight sub-regions and the location of all 306 in situ
338 | measurement sites, 289 of which were located in the eight sub-regions in this study.

339 | **2.4 Evaluation strategy**

340 | For the CLM4.5 simulation, we first spun up 300 years by repeating 30-yr (1979–2009)

Binghao Jia 15/8/21 9:21 AM
已删除: cm
Binghao Jia 15/8/21 9:22 AM
已删除: cm
Binghao Jia 15/8/21 9:22 AM
已删除: cm
Binghao Jia 15/8/21 9:22 AM
已删除: cm

Binghao Jia 15/8/21 9:22 AM
已删除: Here
Binghao Jia 15/8/21 9:23 AM
已删除: selected
Binghao Jia 15/8/21 9:25 AM
已删除: :

Binghao Jia 15/8/21 9:26 AM
已删除: is over

binghaojia 15/7/10 5:59 PM
已删除: 308
binghaojia 15/7/13 11:18 AM
已删除: at those depths
binghaojia 15/7/10 5:59 PM
已删除: 308
binghaojia 15/7/13 7:10 PM
已删除: 291
binghaojia 15/8/11 2:20 PM
已设置格式: 正常, 缩进: 首行缩进: 2 字符

Binghao Jia 15/8/21 9:42 AM
已删除: i
Binghao Jia 15/8/21 9:42 AM
已删除: -
Binghao Jia 15/8/21 9:42 AM
已删除: year

356 ITP meteorological forcing data to achieve an equilibrium state. The 1979–2012 atmospheric
357 forcing data were then used to force CLM4.5 and the simulated results were used in this
358 study. To be consistent with the ESA CCI SM dataset, these simulations were all made at
359 spatial resolution of $0.25^\circ \times 0.25^\circ$ at 30-min time-steps. It should be noted that the data
360 affected by snow cover and temperatures below 0 °C for both the in situ measurements and
361 CLM4.5 predictions between March and October were masked using the quality flags of the
362 ECV CCI SM product. Given the low temporal frequency of the in situ datasets, the
363 validation of the ESA CCI SM and CLM4.5 was conducted at the monthly timescale to
364 reduce the effect caused by the mismatch between actual observation and model time (Wang
365 and Zeng, 2011; Liu and Xie, 2013). Only daily data of the ESA CCI SM product and
366 CLM4.5 simulation on the days of each month when in situ observations were available were
367 used to compute their monthly mean values.

368 The ‘nearest neighbor’ approach was retained to match the grid point location from the
369 ESA CCI SM product or CLM4.5 simulation with that of the in situ measurements. Since soil
370 layer thickness of CLM4.5 model did not match that of in situ observations (0–10 cm), the
371 weighted average was computed based on top four soil layer thicknesses (1.75, 2.76, 4.55,
372 0.94 cm, respectively).

373 Previous studies (Loew et al., 2013; Dorigo et al., 2015) pointed out that the statistical
374 metrics, e.g., the bias and the root mean square difference (RMSD), principally reflected the
375 differences between the in-situ data and the GLDAS-Noah model dataset that was used as a
376 reference for scaling, and thus were scientifically not meaningful. Besides, at the level of the
377 individual stations little is known about the accuracy of the measurements themselves and the
378 ability of the sites to represent absolute soil moisture levels over the coarse satellite footprint
379 scale (Gruber et al., 2013). Therefore, the unbiased RMSD (ubRMSD) was used to express
380 differences in soil moisture levels in this study, and can be calculated using Eq. (1):

Binghao Jia 15/8/21 9:42 AM

已删除: and initial conditions

Binghao Jia 15/8/21 9:44 AM

已删除: will be

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

Binghao Jia 15/8/21 9:44 AM

已删除: run

binghaojia 15/7/10 11:30 AM

已删除:

binghaojia 15/7/10 11:30 AM

已删除:

Binghao Jia 15/8/21 9:44 AM

已删除:

Binghao Jia 15/8/21 9:45 AM

已删除: ute

Binghao Jia 15/8/21 9:45 AM

已删除:

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

$$\text{ubRMSD} = \sqrt{\frac{1}{n} \sum_{i=1}^n [(S_i - \bar{S}) - (O_i - \bar{O})]^2} \quad (1)$$

in which S represents soil moisture from either the ESA CCI SM dataset or CLM4.5 simulation, O is the in situ measurement and \bar{S} and \bar{O} are the corresponding mean values related to S and O . It is noted that, prior to computing the ubRMSD, both the ESA CCI SM dataset and CLM4.5 simulations were scaled into the dynamic range of the in situ data using a linear rescaling method based on the mean and standard deviation (Brocca et al., 2013):

$$S_{\text{res}} = \frac{S - \bar{S}}{\sigma_S} \times \sigma_O + \bar{O} \quad (2)$$

where S_{res} is the linearly rescaled soil moisture dataset, and σ_S and σ_O are the corresponding standard deviations related to S and O , respectively. In addition, the Spearman rank correlation (R_{sp}) was calculated to describe the temporal agreement between the in situ data and the ESA CCI SM dataset and CLM4.5 simulation (Dorigo et al., 2015).

Because the different products used to develop ESA CCI SM vary over space and time, the differences in the microwave observation channels and sampling densities are expected to influence the quality of the different periods for evaluating the new merged satellite product (Liu et al., 2012; Dorigo et al., 2015). The following describes the eight sub-periods used to construct the ESA CCI SM dataset (Albergel et al., 2013a; Dorigo et al., 2015):

- blend 1: January 1979–August 1987, based on SMMR observations only;
- blend 2: September 1987–June 1991, based on SSM/I only;
- blend 3: July 1991–December 1997, based on a combination of SSM/I and ERS AMI;
- blend 4: January 1998–June 2002, based on a combination of TMI and AMI between 40 °N and 40 °S, and a combination of SSM/I and ERS AMI elsewhere;
- blend 5: July 2002–December 2006, based on a combination of AMSR-E and ERS AMI;

binghaojia 15/7/10 5:58 PM

已删除: Taylor diagrams are used to represent these three statistics using two-dimensional plots (Taylor, 2001).

binghaojia 15/7/20 9:06 PM

已删除: following five statistical indicators (Eqs. 1–5) are

binghaojia 15/7/20 9:07 PM

已删除: used to

binghaojia 15/7/20 9:07 PM

已删除: quantitatively evaluate the accuracy of

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/7/20 9:12 PM

已删除: product

binghaojia 15/7/20 9:12 PM

已删除: simulation: mean bias (BIAS), root-mean-squared difference (RMSD), correlation coefficient (R), the normalized standard deviation (SDV), and the centered normalized RMSD (E). They are defined as follows:

Taylor diagrams are used to represent these three statistics using two-dimensional plots (Taylor, 2001).

binghaojia 15/7/20 9:39 PM

已删除: Taylor diagrams are used to represent these three statistics using two-dimen...

Binghao Jia 15/8/21 10:17 AM

已删除: In the following section, we...

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

Binghao Jia 15/8/21 10:18 AM

已删除: ing

binghaojia 15/6/29 4:27 PM

已删除: 2014

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/6/29 4:27 PM

已删除: 2014

Binghao Jia 15/8/21 10:19 AM

已删除:

Binghao Jia 15/8/21 10:19 AM

已删除:

Binghao Jia 15/8/21 10:19 AM

已删除:

Binghao Jia 15/8/21 10:18 AM

已删除:

Binghao Jia 15/8/21 10:19 AM

已删除:

Binghao Jia 15/8/21 10:19 AM

已删除:

Binghao Jia 15/8/21 10:19 AM

已删除:

- 457 • blend 6: January 2007–September 2011, based on a combination of AMSR-E and ASCAT;
- 458 • blend 7: October 2011–June 2012, based on a combination of WindSat and ASCAT;
- 459 • blend 8: July 2012–December 2013, based on a combination of AMSR2 and ASCAT.

460 To illustrate the potential effects of these developmental stages on the quality of the ESA CCI
461 SM dataset, the product evaluation was repeated for the last five time periods (during which
462 in situ observations were available).

463 3. Results

464 3.1 Availability of the ESA CCI SM data in China

465 Previous studies (Loew et al., 2013; Dorigo et al., 2015) showed that the performance of
466 the ESA CCI SM product may be strongly affected by data gaps and the period of
467 observation. Therefore, we first analyzed the data availability of the ESA CCI SM product in
468 China for the entire period (1979–2013) as well as for the eight individual time periods
469 (described in Section 2.4). A clear increase of observation density can be observed over time
470 (Fig. 2), which is mainly due to the growing number of satellites available for soil moisture
471 data retrieval (Liu et al., 2012; Dorigo et al., 2015). The improved instrument design and
472 sensor performance also contributes to this effect. Better sensors have led to a convergence of
473 active and passive remote sensing products, especially over areas with intermediate
474 vegetation cover (Liu et al., 2012), and an increase over time in the number of areas where
475 both products are used in a synergistic way.

476 For example, for the second period (1 September 1987–30 June 1991), the ESA CCI SM
477 product was generated using the SSM/I Ku-band (19.3 GHz) data alone, which was strongly
478 attenuated by the vegetation canopy; thus, large areas were masked due to moderate to dense
479 vegetation, such as northeast and southern China (Fig. 2b). The introduction of the ERS AMI
480 C-band (5.3 GHz) scatterometer during the subsequent period (1 July 1991–31 December
481 1997) was able to partly fill these gaps (Fig. 2c), e.g. there was a clear increase in the number

Binghao Jia 15/8/21 10:19 AM

已删除:

Binghao Jia 15/8/21 10:19 AM

已删除:

Binghao Jia 15/8/21 10:19 AM

已删除:

Binghao Jia 15/8/21 10:19 AM

已删除:

Binghao Jia 15/8/21 10:19 AM

已删除:

Binghao Jia 15/8/21 10:20 AM

已删除:

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

Binghao Jia 15/8/21 10:20 AM

已删除:

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/6/29 4:27 PM

已删除: 2014

binghaojia 15/6/29 4:27 PM

已删除: Loew, 2014

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

Binghao Jia 15/8/21 10:21 AM

已删除: any

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/6/29 4:27 PM

已删除: 2014

Binghao Jia 15/8/21 10:22 AM

已删除: They

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

Binghao Jia 15/8/21 10:24 AM

已删除: and it then led to

Binghao Jia 15/8/21 10:24 AM

已删除: masked

Binghao Jia 15/8/21 10:24 AM

已删除: with

Binghao Jia 15/8/21 10:25 AM

已删除: of

of observations over southern China. With the introduction of the high quality AMSR-E C-band (6.9 GHz) data in July 2002, a large increase of the observation density can be found over most areas of China, except the Tibetan Plateau (Fig. 2e). As a result of the unavailability of the AMSR-E data in October 2011, the observation density decreased over large parts of northwest China and Inner Mongolia, which are mainly arid or semiarid areas, despite the inclusion of the high quality WindSat data for this period (Fig. 2g). In addition, the introduction of the AMSR2 product clearly increases the fractions of valid observations over most areas of China, except central and southern China (Fig. 2h).

3.2 Evaluation using in situ data

Figure 3 presents the Spearman rank correlation coefficients (R_{sp}) of the ESA CCI SM product and CLM4.5 simulation against in situ measurements, in which the values from both techniques are ranked using the same intervals. The data gaps for the ESA CCI SM product (Fig. 2) lead to only 299 stations available; over 61% of these stations had correlation values between 0.2 and 0.6 (shown in green and purple, Fig. 3a), with a mean of 0.27 (Table 2). In contrast, the CLM4.5 model captured surface soil moisture temporal variability better (averaged R value is 0.35, Table 2) than the ESA CCI SM product; about 10% of the stations had R_{sp} values greater than 0.6 (shown as orange and blue, Fig. 3b), whereas only 5% stations for the ESA CCI SM achieved such correlations (Fig. 3a). When only the configurations associated to significant correlation values ($p \leq 0.05$) are considered (cycles), there were 212 and 253 stations available for the ESA CCI SM and CLM4.5, respectively. In the current study, the averaged R_{sp} value of the ESA CCI SM is 0.37, which is slightly higher than that ($R_{sp} = 0.32$) from 34 sites over China in Dorigo et al. (2015); the averaged R_{sp} is 0.42 for the CLM4.5 model.

As described in several studies (Liu et al., 2011, 2012; Dorigo et al., 2015), the mean and dynamic range of the ESA CCI SM time series represent those of the GLDAS-Noah

- Binghao Jia 15/8/21 10:25 AM
已删除: .
- binghaojia 15/7/21 7:32 PM
已删除: biases and RMSDs...of the ... [11]
- Binghao Jia 15/8/21 10:26 AM
已删除: with...in situ measurement ... [12]
- binghaojia 15/6/29 2:26 PM
已删除: ECV-SM...SA CCI SM prc ... [13]
- Binghao Jia 15/8/21 10:27 AM
已删除: have
- binghaojia 15/7/21 7:40 PM
已删除: 4a...a), with a mean of 0.35 ... [14]
- Binghao Jia 15/8/21 10:29 AM
已删除: s
- binghaojia 15/7/21 9:31 PM
已删除: ing...surface soil moisture ... [15]
- Binghao Jia 15/8/21 10:29 AM
已删除: ve
- binghaojia 15/7/21 7:47 PM
已删除: circles... Fig. 4b ... [16]
- Binghao Jia 15/8/21 10:31 AM
已删除: but
- binghaojia 15/7/21 10:20 PM
已删除: 10...% stations for the ... [17]
- Binghao Jia 15/8/21 10:32 AM
已删除: =...0.05) are considered (c ... [18]
- binghaojia 15/6/29 2:26 PM
已删除: ECV-SM...SA CCI SM an ... [19]
- Binghao Jia 15/8/21 10:35 AM
已删除: ...while it ... [20]
- binghaojia 15/7/22 11:07 AM
已删除: and ...0.....52...2 for the C ... [21]

surface soil moisture product. Therefore, we used a linear rescaling method (Eq. 2) to rescale both the ESA CCI SM dataset and CLM4.5 soil moisture predictions prior to computing their ubRMSDs (Brocca et al., 2013) against in situ measurements (Fig. 4). For the ESA CCI SM product, there were 214 stations (72% of total) with the ubRMSD values less than $0.06 \text{ m}^3 \text{ m}^{-3}$. The CLM4.5 showed a slightly improved ubRMSD, of which about 29% of the stations had a value less than $0.04 \text{ m}^3 \text{ m}^{-3}$ (Fig. 4b), compared to 22% using the ESA CCI SM product (Fig. 4a). Table 2 shows that averaged ubRMSD values are 0.053 and $0.050 \text{ m}^3 \text{ m}^{-3}$ for the ESA CCI SM product and CLM4.5 simulation, respectively, which suggests that the CLM4.5 simulation is closer to in situ measurements in China than is the ESA CCI SM product. When only sites with a significant ($p < 0.05$) positive Spearman's correlation were considered (cycles in Figs. 3 and 4), the averaged ubRMSD values were 0.050 and $0.048 \text{ m}^3 \text{ m}^{-3}$ for the ESA CCI SM product and CLM4.5 simulation, respectively. It should be noted that because the ubRMSD is only an indication of the mismatch between in situ data and remote sensing or model simulated datasets, the errors in each dataset are also included. The triple collocation technique has proven to be an independent technique for estimating the random error component of soil moisture datasets and been applied to the direct comparison with the ubRMSD metric (Dorigo et al., 2010; Gruber et al., 2013; Dorigo et al., 2015). Therefore, future work is expected to use this method to reduce the effect of random errors in each dataset on the comparison.

3.3 Performances over eight sub-regions

Since soil moisture has a close relationship with the status of climate dryness and wetness, eight sub-regions were defined (Table 1 and Fig. 1) in this study based on Zhu (2003), which used a time series for dryness and wetness encompassing the last 530 years to analyze the spatial patterns of the centers of dryness and wetness throughout China by applying the rotated empirical orthogonal function method. To evaluate the performance of

624 the satellite-based product and process-based model simulation over eight sub-regions, two
625 evaluation strategies were adopted in our study: (1) we computed the statistical metrics for
626 the stations individually (e.g., Figs. 3 and 4) and then averaged their values over each region
627 (hereinafter referred to as "S1"); (2), we generated the time series of soil moisture by
628 averaging available observations at all stations of each region, only considering those grid
629 cells closest to the relevant observation stations for the ESA CCI SM product and CLM4.5
630 simulation, and then calculated their statistical metrics against in situ measurements
631 (hereinafter referred to as "S2"). It should be noted that only sites with a significant ($p < 0.05$)
632 positive Spearman's correlation (cycles in Figs. 3 and 4) for both ESA CCI SM product and
633 CLM4.5 simulation were considered in the following analysis whatever the evaluation
634 strategy. After screening using the criteria above, 200 stations remained available for the
635 following evaluation, 189 of which were located in the eight sub-regions; the number of valid
636 stations for each sub-region is shown in Table 2 (in the brackets). The statistical metrics of
637 the ESA CCI SM product and CLM4.5 simulation against in situ measurements for both
638 evaluation strategies are presented in Fig. 5.

639 Over most regions (except for China V), the ESA CCI SM product, of which average
640 ubRMSD values ranged between 0.044 and 0.059 $\text{m}^3 \text{m}^{-3}$, had slightly higher ubRMSD
641 values than CLM4.5 simulations (0.042–0.056 $\text{m}^3 \text{m}^{-3}$) for the first evaluation strategy (Fig.
642 5a), which is consistent with the results from Fig. 4 and Table 2. Figure 5b shows that the
643 average Spearman correlation coefficients of the ESA CCI SM product ranged between 0.27
644 and 0.47, while higher correlations in situ measurements were achieved by CLM4.5
645 simulations (0.34–0.54), except in China V. Good performance over China V, which is a
646 semi-arid region, for the ESA CCI SM product was also found by Albergel et al. (2013b) and
647 Dorigo et al., (2015) using in situ observations from the Maqu network, which is located over
648 the southern part of this sub-region. Our finding suggests that the ESA CCI SM product has a

binghaojia 15/7/22 4:23 PM

已删除: of China (Table 1 and Fig. 1), their statistical metrics with in situ measurements are presented in Fig. 5. It should be noted that we first generated

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/7/22 4:55 PM

已删除: ;

binghaojia 15/7/22 4:55 PM

已删除: used them to

binghaojia 15/7/22 4:48 PM

已删除: performances

binghaojia 15/7/22 4:50 PM

已删除: .

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/7/23 4:33 PM

已删除: underestimates soil moisture (negative values) except for southwest China (China VII and VIII, Fig. 5a). Conversely,

binghaojia 15/7/23 4:14 PM

已删除: the

binghaojia 15/7/23 4:14 PM

已删除: simulation shows clearly positive biases over all sub-regions except for China VI and VIII, which suggests a systematic overestimation for model simulation. Larger RMSDs for the CLM4.5 can be observed in eastern China (China I–IV), which may be due to the deficiencies in CLM4.5; for example, scaling of canopy interception (Lawrence et al., 2007), soil texture and model structure (Liu and Xie, 2013). The ECV-SM product shows lower RMSD values than the CLM4.5 simulation, with the exception of China VI (Fig. 5b)

binghaojia 15/7/23 4:40 PM

已删除: 2

677 good potential for application in areas where models cannot realistically represent soil
678 moisture (Albergel et al., 2013b).

679 Compared to S1, the other strategy (S2) showed large variations in ubRMSD (Fig. 5a)
680 and R_{sp} (Fig. 5b) between different sub-regions. However, the two strategies produced similar
681 results although CLM4.5 simulations showed a slightly higher R_{sp} and lower ubRMSD than
682 the ESA CCI SM product. In addition, compared with performance in other regions, both
683 remotely sensed and modeled soil moisture datasets showed larger ubRMSD values over
684 southwest China (China VII and VIII), which is mainly covered with moderate to dense
685 vegetation, including mixed evergreen coniferous and broadleaf deciduous forests, and
686 broadleaf deciduous forests. This result is mainly due to strong attenuation by the vegetation
687 canopy of the microwave retrievals for the ESA CCI SM product (Liu et al., 2012; Dorigo et
688 al., 2015). For the CLM4.5 simulation, the precipitation data in the ITP atmospheric forcing
689 data were generated by merging ground-based observations and two satellite retrieval
690 products, because few in situ precipitation measurements were available over these areas
691 (Chen et al., 2011).

692 Figure 6 compares the annual soil moisture cycle (March–October) of the ESA CCI SM
693 product and CLM4.5 simulation against in situ measurements, averaged over the eight
694 sub-regions using the S2 evaluation strategy (without the linear scaling). Both the ESA CCI
695 SM product and CLM4.5 simulation generally captured the annual cycle well in most
696 sub-regions. However, soil moisture simulated by the CLM4.5 model was usually greater
697 than the observations, showing systematic overestimation in all sub-regions except for China
698 VI and VIII; this is consistent with the research of Liu and Xie (2013), which was based on
699 simulations with the CLM3.5 model using the same atmospheric forcing data as we used. The
700 comparable findings suggest that land surface model simulation may reproduce temporal
701 variations well, but fail to simulate the mean soil moisture (Entin et al., 2000; Gao and

Binghao Jia 15/8/21 2:20 PM

已删除: while

Binghao Jia 15/8/21 2:20 PM

已删除: ing

Binghao Jia 15/8/21 2:22 PM

已删除: ere

Binghao Jia 15/8/21 2:22 PM

已删除: are

Binghao Jia 15/8/21 2:23 PM

已删除: It

Binghao Jia 15/8/21 2:23 PM

已删除: on

Binghao Jia 15/8/21 2:24 PM

已删除: while

binghaojia 15/7/23 5:22 PM

已删除: However, Fig. 5c shows that the CLM4.5 simulation correlates better with in situ data ranging from 0.51 (China VI) to 0.81 (China III), compared to the ECV-SM, which ranges from 0.34 (China VI) to 0.71 (China VIII). A negative R value of the ECV-SM for China II is also observed in Fig. 5c. This is because the ECV-SM product overestimates soil moisture over northern parts of China II while the main underestimations occur over southern parts; and the averaging at all stations of this region leads to a weak temporal variation with in situ measurements.

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/7/24 9:53 AM

已删除: , but

Binghao Jia 15/8/20 4:44 PM

已删除: clearly

Binghao Jia 15/8/21 2:30 PM

已删除: over

binghaojia 15/7/24 9:54 AM

已删除: .

728 Dirmeyer, 2006; Liu and Xie, 2013). In contrast, the ESA CCI SM product was closer to
729 matching in situ measurements than the CLM4.5 model over some areas (e.g., China V in this
730 study, a semi-arid region), although it represents the dynamic range of the GLDAS-Noah
731 surface soil moisture product (Liu et al., 2012; Dorigo et al., 2015).

732 3.4 Comparison of the anomaly time series

733 The results presented above are based on comparisons of the multi-year observation
734 dataset and the ESA CCI SM or CLM4.5 datasets on a monthly timescale, which may be
735 somewhat affected by the seasonal cycle of soil moisture. The time series of the soil moisture
736 anomaly from the three datasets (ESA CCI SM, CLM4.5, and in situ observations) for the
737 eight sub-regions, using the S2 strategy (Section 3.3), computed by removing the multi-year
738 annual cycle (only March–October available), are presented in Fig. 7. In general, both the
739 ESA CCI SM product and the CLM4.5 simulation captured the temporal evolution of the
740 observed soil moisture anomalies reasonably well for most of the regions, except a slight
741 overestimation in the amplitude of fluctuations over China VII and VIII by the ESA CCI SM
742 product, which was consistent with the results shown in Fig. 5.

743 As a further quantitative illustration of the effects of seasonal cycle on the evaluation
744 results, Fig. 8 shows the Spearman correlation coefficients (R'_{sp}) for the anomaly data of both
745 the ESA CCI SM product and CLM4.5 simulation over the eight sub-regions. In general, R'_{sp}
746 values (Fig. 8) were lower than R_{sp} (Fig. 5) for most sub-regions, especially over China VI
747 and VIII. This differences suggested that the seasonal cycle in soil moisture has a large effect
748 on the comparison of Spearman correlation values over the areas with pronounced wet and
749 dry periods (Liu et al., 2012; Albergel et al., 2013a; Dorigo et al., 2015).

750 3.5 Evolution of the ESA CCI SM over time

751 The microwave observation channels and sampling densities were expected to affect the
752 quality of the dataset for different periods because the different products that were used to

binghaojia 15/7/24 10:09 AM

已删除: However, compared to the ECV-SM, CLM4.5 correlates better with the observations. The ECV-SM is drier than the observations in spring over most parts of China, except China VII.

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/7/24 3:09 PM

已删除:

binghaojia 15/7/24 3:09 PM

已删除: (Table 1)

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

binghaojia 15/7/24 3:17 PM

已删除: . However, the ECV-SM product (red line) slightly overestimates the amplitude of fluctuations over China I, VII, and VIII. In contrast, CLM4.5 simulation tends to reproduce the temporal variation of surface soil moisture anomalies better because it is sensitive to precipitation (Gao and Dirmeyer, 2006; Liu and Xie, 2013). The precipitation data in the ITP atmospheric forcing data were generated by merging ground-based observations and two satellite retrieval products (Chen et al., 2011). Due to strong attenuation by the vegetation canopy on the microwave retrievals, the soil moisture anomalies of the ECV-SM product tends to be considerably stronger than in situ observations and generates spurious temporal variations over northeast (China I) and southwest China (China VII and VIII). These areas are mainly covered with moderate to dense vegetation, including mixed evergreen coniferous ... [22]

binghaojia 15/7/27 5:56 PM

已删除:

binghaojia 15/7/27 6:21 PM

已删除: two Taylor diagrams

binghaojia 15/7/27 6:21 PM

已删除: original and

binghaojia 15/7/27 6:25 PM

已删除: comparing the two datasets with in situ measurements

binghaojia 15/7/27 6:25 PM

已删除: R

binghaojia 15/7/28 9:21 AM

已删除: over China VI and VIII for both the ECV-SM and CLM4.5

binghaojia 15/7/28 10:26 AM

已删除: It is also clear that both the ECV-SM and CLM4.5 have very we ... [23]

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

Binghao Jia 15/8/21 2:44 PM

已删除: Since

generate the ESA CCI SM dataset vary over space and time (Liu et al., 2012; Albergel et al., 2013a). Figure 9 presents the averaged ubRMSDs and R_{sp} of the ESA CCI SM product for the last five individual time periods defined in Section 2.4. Due to limited time length and sparse samples for the last two blended periods (blend 7 and blend 8), they were combined into one period. Considering the sites that have significant R_{sp} values ($p = 0.05$) for all periods (denoted by red in Fig. 9), averaged correlations ranged from 0.42 to 0.62 while ubRMSD values ranged from 0.024 m³ m⁻³ to 0.045 m³ m⁻³. A similar pattern was obtained when considering the sites for which the correlation was significant for each blended period with averaged R_{sp} values ranging from 0.56 to 0.93.

For the fourth period (January 1998–June 2002), a higher correlation was observed (Fig. 9b) compared to the previous period (January 1993–December 1997), which may be related to the introduction of the circular non-polar orbiting TMI data in 1998 (Doirgo et al., 2015). In July 2002, a high quality soil moisture dataset from C-band AMSR-E retrievals was introduced instead of the SSM/I and TMI, which increased the correlations of the ESA CCI SM dataset with in situ measurements and reduced its ubRMSD values for the fifth period (July 2002–December 2006). These observations were consistent with the results of the comparison between the ESA CCI SM and the ERA-Land, an update of the land surface component of the ERA-Interim reanalysis from the European Centre for Medium-Range Weather Forecasts (Albergel et al. 2013a), and that between the ESA CCI SM product and in situ measurements from other soil moisture networks (Dorigo et al., 2015). Unexpectedly, there was a decrease in quality in the sixth period (January 2007–September 2011), with weaker correlations and higher ubRMSD (Albergel et al., 2013a; Dorigo et al., 2015). The cause of this degradation is still not entirely clear, but it may be related to the resampling and scaling strategy used to incorporate a new active input product from ASCAT (Dorigo et al., 2015). The best performance for all metrics was obtained for the last two periods (October

binghaojia 15/6/29 2:26 PM

已删除: ECV-SM

Binghao Jia 15/8/21 2:45 PM

已删除: , the microwave observation channels and sampling densities are expected to affect the quality for different periods... (L... [24]

binghaojia 15/7/28 10:39 AM

已删除: correlation coefficients...sp... [25]

Binghao Jia 15/8/21 2:46 PM

已删除: which

binghaojia 15/7/27 6:33 PM

已删除: 46 ...2 to 0.69 ...2 while ut... [26]

Binghao Jia 15/8/21 2:47 PM

已删除: is...obtained when consider... [27]

binghaojia 15/7/27 6:33 PM

已删除: 54 ...6 to 0.91 [28]

Binghao Jia 15/8/21 2:48 PM

已删除: is

binghaojia 15/7/27 6:35 PM

已删除: 2014...015). In July 2002, ... [29]

Binghao Jia 15/8/21 2:49 PM

已删除: is is

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM...SA CCI SM an... [30]

Binghao Jia 15/8/21 2:52 PM

已删除: we see a drop...in quality ic... [31]

binghaojia 15/6/24 2:19 PM

已删除: 07

Binghao Jia 15/8/21 2:52 PM

已删除: , also found by ...lbergel et... [32]

binghaojia 15/6/29 4:27 PM

已删除: 2014...015). The cause of t... [33]

936 2011–December 2012), which was expected, given the higher quality satellite
937 instrumentation (AMSR2) and improved retrieval algorithms. However, the shorter length of
938 time within the two periods may have had a slight effect on the statistical metrics and
939 therefore more in situ measurements would be needed to confirm our results.

940 4. Discussion

941 During the past twenty years, huge efforts have been made to make more in situ soil
942 moisture observations available in China. These measurements are important for the
943 evaluation of both remotely sensed and modeled soil moisture. In this study, we evaluated the
944 performances of the ESA CCI SM product and CLM4.5 simulation in China using 20 years
945 of in situ observations from 306 sites. However, some limitations to the evaluation should be
946 noted. The spatial representativity, temporal mismatch, instrumental errors, and installation
947 depth of in situ observations may have had a negative effect on the evaluation results. Such
948 limitations have been extensively reported in previous studies (Brocca et al., 2010; Crow et
949 al., 2012). Therefore, future work is expected to reduce these uncertainties through the
950 enhancement of ground-based measurements and the improvement of the evaluation strategy.

951 In addition, soil moisture was measured using the gravimetric technique. This destructive
952 sampling method may affect the evaluation results (Dorigo et al., 2015). According to the
953 user guide of in situ soil moisture measurements from agricultural meteorological stations of
954 CMA (in Chinese, <http://cdc.nmic.cn>), soil moisture observation technique can be
955 summarized as follows: (1) the observation field of each station is divided to four parts and
956 four soil samples will be collected each time; (2) their soil moisture contents in dry weight
957 basis (the ratio of water mass to dried soil's weight) are determined by drying the soil,
958 respectively; (3) the average value from the four samples are recorded as mass percentage for
959 this station. It is noted that the horizontal distance between two successive samples at the
960 same part of each station is no more than 2 meters, which leads to almost the same

binghaojia 15/7/28 11:08 AM

已删除: 2011

Binghao Jia 15/8/21 2:53 PM

已删除: is

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

binghaojia 15/8/18 2:03 PM

已删除: more than 300

Binghao Jia 15/8/21 3:01 PM

已删除: lead to

966 meteorological and soil conditions. Moreover, to reduce the effect of soil moisture
967 heterogeneity, the station was usually chosen to be over flat surface. However, the original
968 observations from the four samples at each station were not available; thus, we could not
969 investigate the effect of destructive sampling on the comparison results in this study. Previous
970 studies (Brocca et al., 2014a, b) showed that soil moisture had enormous variability even at
971 local scales, in particular in absolute terms. These methods will be used in our future work to
972 discuss the impacts of spatial variability and destructive sampling method.

973 Land surface models can capture the temporal dynamics of soil moisture well when
974 forced by high quality atmospheric forcing data (Albergel et al., 2010, 2012) and are usually
975 employed to upscale in situ surface soil moisture observations or complete the evaluation of
976 satellite derived products (Albergel et al., 2013a). In this study, the CLM4.5 was forced by an
977 atmospheric forcing dataset generated using many ground-based observations, and showed
978 better correlation with in situ soil moisture observations than did the ESA CCI SM product.
979 However, many precipitation measurements were located in the same 0.25° grid box as the in
980 situ soil moisture observations; although the two types of data were obtained from different
981 sources and were not collected at the same locations, it was necessary to investigate the effect
982 of in situ precipitation measurements on the comparison results. The averaged statistical
983 metrics over the sites with ("Rain") and without ("No Rain") in situ precipitation
984 measurements are presented in Fig. 10. It is noted that only the sites (200 of 306) with a
985 significant ($p < 0.05$) positive Spearman's correlation for both the ESA CCI SM product and
986 CLM4.5 simulation were considered, as described in Section 3.2. Figure 10 shows that, for
987 the CLM4.5 simulation, in situ precipitation measurements (101 of 200) led to a slightly
988 higher correlation ($R_{sp} = 0.46$) with in situ soil moisture measurements than was obtained
989 without the precipitation measurements ($R_{sp} = 0.44$, 99 of 200). This may be related to the
990 issues of spatial resolution. The observations from 740 operational stations of the CMA were

Binghao Jia 15/8/21 3:09 PM

已删除: with

Binghao Jia 15/8/21 3:09 PM

已删除: used

Binghao Jia 15/8/21 3:10 PM

已删除: lots of

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

merged with other meteorological forcing datasets to generate the ITP forcing data with a spatial resolution of $0.1^{\circ} \times 0.1^{\circ}$, while the CLM4.5 was run at $0.25^{\circ} \times 0.25^{\circ}$ resolution in this study. If the CLM4.5 had been run at $0.1^{\circ} \times 0.1^{\circ}$ resolution, the effect of ground-based precipitation on the comparison results may have been clearer (not shown in this study).

The in situ observations used in this study are average soil moisture values of surface soil of 0–10 cm; to be consistent with the in situ observations, the weighted average of CLM4.5 at 0–10 cm was computed based on the uppermost four soil layer thicknesses (Section 2.4). However, the ESA CCI SM data represent the upper 0–2 cm of the soil (Liu et al., 2011, 2012; Dorigo et al., 2015). As in situ soil moisture observations of 0–2 cm are not available, it is difficult to investigate the effect of the mismatch between soil depths on the statistical results. Instead, we compared the in situ observations and ESA CCI SM data with average soil moisture values of CLM4.5 at different soil depths: first layer (“L1”, 0–1.75 cm), uppermost two layers (“L1–2”, 0–1.75 cm, 1.75–4.51 cm), uppermost three layers (“L1–2–3”, 0–1.75 cm, 1.75–4.51 cm, 4.51–9.06 cm), 0–2 cm, 0–5 cm, and 0–10 cm. Their statistical metrics, including ubRMSD and R_{sp} , are presented in Fig. 11. Results showed that the choice of soil depth for CLM4.5 had a significant effect on the statistics. The average R_{sp} against in situ observations ranged between 0.417 and 0.425 at different soil depths while the ESA CCI SM had higher R_{sp} values and a larger range, which was between 0.507 and 0.546. CLM4.5 agreed better with in situ observations at 0–10 cm than those at other soil depths, confirming that it is better to evaluate the CLM4.5 using the weighted average values at 0–10 cm. In contrast, the best agreement between the CLM4.5 simulation and ESA CCI SM product was found at the first layer (0–1.75 cm). These findings suggest that the mismatch in soil depths between the ESA CCI SM and in situ observations may have had a large effect on the statistical metrics, which is one of the reasons for its higher ubRMSD and lower R_{sp} .

In addition, although the CLM4.5 model had better performance against in situ

binghaojia 15/7/29 12:01 PM

已删除: However

Binghao Jia 15/8/20 4:40 PM

已删除: In addition,

1022 | observations than the ESA CCI SM product over most regions in China, there were still many
 1023 | unrealistic representations in model parameterizations (soil water and temperature, for
 1024 | example) and large uncertainties in land surface datasets (such as soil texture and PFTs) for
 1025 | the CLM4.5 model, which affected its accuracy in soil moisture simulation. To improve soil
 1026 | moisture estimations, many studies have incorporated remotely sensed brightness
 1027 | temperatures (Jia et al., 2013) or soil moisture retrievals (Draper et al., 2012) into land
 1028 | surface models using various modern land data assimilation systems. Our study provides an
 1029 | in-depth evaluation of both the ESA CCI SM product and CLM4.5 model in China, including
 1030 | their performances over different sub-regions. This evaluation is expected to be useful for the
 1031 | assimilation of the ESA CCI SM product into land surface models, especially the CLM4.5,
 1032 | because the observation and model errors have been estimated approximately.

1033 | For the ESA CCI SM product, only soil moisture retrievals acquired by night time
 1034 | overpasses (occurring between 19:00 and 08:00 local time) were used (Liu et al., 2012)
 1035 | because all input products had varying local observation times (Dorigo et al., 2015). However,
 1036 | the original in situ soil moisture observations were available every 10 days (three times each
 1037 | month) and the time step of CLM4.5 was 30 min. To reduce the effect caused by the
 1038 | mismatch between actual observation and model time, we compared the three datasets at a
 1039 | coarser time scale using their monthly values, in a similar way as previous studies (Li et al.,
 1040 | 2005; Wang and Zeng, 2011; Liu and Xie, 2013). However, it should be noted that the effect
 1041 | of the temporal mismatch among different soil moisture datasets was not considered in this
 1042 | study.

1043 | 5. Conclusions

1044 | In this study, the performances of a microwave-based merged satellite product (ESA
 1045 | CCI SM) and the CLM4.5 simulation were investigated using 20 years of in situ observations
 1046 | from 306 stations in China. In general, both soil moisture products represent in situ

Binghao Jia 15/8/21 3:34 PM

已删除: are...still many unrealistic [34]

binghaojia 15/7/29 1:42 PM

已删除: However, data assimilation is beyond the scope of the present study. Nonetheless, o

Binghao Jia 15/8/21 3:37 PM

已删除: This

binghaojia 15/7/29 1:48 PM

已删除: ur...study provides an in-d [35]

Binghao Jia 15/8/21 3:41 PM

已删除: ,

binghaojia 15/7/29 1:48 PM

已删除: . This

Binghao Jia 15/8/21 3:42 PM

已删除: which

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

Binghao Jia 15/8/21 3:42 PM

已删除: since

binghaojia 15/7/29 12:00 PM

已设置格式

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

Binghao Jia 15/8/21 3:43 PM

已删除: since

binghaojia 15/6/29 4:28 PM

已删除: 2014

Binghao Jia 15/8/21 3:43 PM

已删除: are ...ere available every 10 [36]

Binghao Jia 15/8/21 3:48 PM

已删除: ing

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

Binghao Jia 15/8/21 3:48 PM

已删除: in China

binghaojia 15/7/13 9:15 PM

已删除: 308

Binghao Jia 15/8/21 3:49 PM

已删除: two ...oil moisture product [37]

1103 observations well, with ubRMSD values of about $0.05 \text{ m}^3 \text{ m}^{-3}$. The ESA CCI SM product has
1104 slightly weaker Spearman correlations ($R_{sp} = 0.37$) than the CLM4.5 model, but the CLM4.5
1105 simulation produces better temporal variation of surface soil moisture ($R_{sp} = 0.42$).

1106 Both the satellite product and model simulation show large discrepancies over eight
1107 sub-regions in China. The CLM4.5 model shows larger ubRMSDs in southwestern China
1108 (China VII and VIII) than other regions, which may be related to the scarcity of in situ
1109 precipitation measurements available in these areas. The ESA CCI SM product is likely to be
1110 best-suited for applicaiton in semi-arid regions (such as China V), even better than the
1111 CLM4.5 model, mainly because of accurate data retrievals and high observation density; in
1112 contrast, the ESA CCI SM product is not well suited for areas covered by dense vegetation
1113 (such as China VII and VIII). In addition, the statistical scores of the ESA CCI SM product in
1114 China corroborate the findings of Albergel et al. (2013a) and Dorigo et al. (2015) of a stable
1115 to slightly improving performance over time in China, with the exception of a decrease
1116 during the 2007–2010 blending period, which may be related to the resampling and scaling
1117 strategy used to incorporate the ASCAT product into the ESA CC SM dataset.

1118 In response to the sparse observation temporal frequency (three times each month), all
1119 analyses were performed on a monthly timescale. Future efforts are expected to make more in
1120 situ soil moisture observations available in China for the evaluation of either remote sensing
1121 retrievals or modeled simulation. Furthermore, data assimilation has been an effective tool to
1122 incorporate remotely sensed soil moisture into land surface models to improve soil moisture
1123 simulation (Draper et al., 2012). The in-depth evaluation of the ESA CCI SM product and
1124 CLM4.5 model in China provided by this study will facilitate soil moisture data assimilation,
1125 which is expected to provide further information regarding observation and model errors.

1126
1127 **Acknowledgements**

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

Binghao Jia 15/8/21 3:50 PM

已删除: a

binghaojia 15/7/29 3:21 PM

已删除: lower bias, with RMSD = $0.075 \text{ m}^3 \text{ m}^{-3}$ but with

binghaojia 15/7/29 3:21 PM

已删除: 41

Binghao Jia 15/8/21 3:50 PM

已删除: . In contrast,

binghaojia 15/7/29 3:21 PM

已删除: 52

binghaojia 15/7/29 3:19 PM

已删除: , but shows clear overestim... [38]

binghaojia 15/7/29 8:09 PM

已删除: simulation is mainly affectc... [39]

binghaojia 15/7/29 3:33 PM

已删除: eastern

binghaojia 15/7/29 8:09 PM

已删除:

Binghao Jia 15/8/21 3:53 PM

已删除: (Chen et al., 2011)

binghaojia 15/7/29 8:03 PM

已删除: caused by inaccurate descri... [40]

Binghao Jia 15/8/21 3:54 PM

已删除: performance of the

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

binghaojia 15/7/29 8:03 PM

已删除: III and

Binghao Jia 15/8/21 3:55 PM

已删除: (Albergel et al., 2013a),

Binghao Jia 15/8/21 3:56 PM

已删除: and worst over

binghaojia 15/7/29 8:03 PM

已删除: IV,

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

Binghao Jia 15/8/21 3:58 PM

已删除: confirm

binghaojia 15/6/29 4:28 PM

已删除: 2014

binghaojia 15/7/29 3:30 PM

已删除: the incorporation of

binghaojia 15/7/29 3:36 PM

已删除: the differences in observati... [41]

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

Binghao Jia 15/8/21 3:59 PM

已删除: in

Binghao Jia 15/8/21 3:59 PM

已删除: be useful for its application in

1170 This research was supported by the National Natural Science Foundation of China (Grant
1171 Numbers, 41305066, 41405083 and 91125016), the Special Funds for Public Welfare of
1172 China (Grant Number, GYHY201306045) and the Natural Science Foundation of Hunan
1173 Province (Grant Number, 2015JJ3098). The ESA soil moisture dataset was downloaded from
1174 <http://www.esa-soilmoisture-cci.org>; the in situ soil moisture observations were downloaded
1175 from <http://cdc.nmic.cn/>. The forcing dataset used in this study was developed by the Data
1176 Assimilation and Modeling Center for Tibetan Multi-spheres, Institute of Tibetan Plateau
1177 Research, Chinese Academy of Sciences. We also thank the editor, Prof. Alexander Loew,
1178 the reviewer, Dr. Wouter Dorigo, and another anonymous reviewer for their constructive
1179 comments and suggestions, which have significantly improved our paper.

Binghao Jia 15/8/21 4:00 PM

已删除: o

Binghao Jia 15/8/21 4:01 PM

已删除: .

Binghao Jia 15/8/21 4:00 PM

已删除: ,

Binghao Jia 15/8/21 4:00 PM

已删除: o.

Binghao Jia 15/8/21 4:01 PM

已删除: o.

binghaojia 15/7/29 10:45 AM

已删除: We thank Prof. Kun Yang for providing the ITP meteorological forcing data

Binghao Jia 15/8/21 4:01 PM

已删除: s

1188 **References**

- 1189 Albergel, C., de Rosnay, P., Gruhier, C., Muñoz-Sabater, J., Hasenauer, S., Isaksen, L., Kerr,
1190 Y., and Wagner, W.: Evaluation of remotely sensed and modelled soil moisture products
1191 using global ground-based in situ observations, *Remote Sens. Environ.*, 118, 215–226,
1192 2012.
- 1193 Albergel, C., Dorigo, W., Balsamo, G., Muñoz-Sabater, J., de Rosnay, P., Isaksen, L., Brocca,
1194 L., de Jeu, R., and Wagner, W.: Monitoring multi-decadal satellite earth observation of
1195 soil moisture products through land surface reanalyses, *Remote Sens. Environ.*, 138,
1196 77–89, 2013a.
- 1197 Albergel, C., Dorigo, W., Reichle, R., Balsamo, G., de Rosnay, P., Muñoz-Sabater, J.,
1198 Isaksen, L., and Wagner, W.: Skill and global trend analysis of soilmoisture
1199 fromreanalyses andmicrowave remote sensing, *J. Hydrometeor.*, 14, 1259–1277, doi:
1200 10.1175/JHM-D-12-0161.1, 2013b.
- 1201 Albergel, C., Rudiger, C., Carrer, D., Calvet, J. C., Fritz, N., Naeimi, V., Bartalis, Z., and
1202 Hasenauer, S.: An evaluation of ASCAT surface soil moisture products with in-situ
1203 observations in Southwestern France, *Hydrol. Earth Syst. Sci.*, 13, 115–124, 2009.
- 1204 Bartalis, Z., Wagner, W., Naeimi, V., Hasenauer, S., Scipal, K., Bonekamp, H., Figa, J., and
1205 Anderson, C.: Initial soil moisture retrievals from the METOP-A advanced scatterometer
1206 (ASCAT), *Geophys. Res. Lett.*, 34, L20401, doi:10.1029/2007GL031088, 2007.
- 1207 Brocca, L., Ciabatta, L., Massari, C., Moramarco, T., Hahn, S., Hasenauer, S., Kidd, R.,
1208 Dorigo, W., Wagner, W., and Levizzani, V. : Soil as a natural rain gauge: Estimating
1209 global rainfall from satellite soil moisture data. *J. Geophys. Res. Atmos.*, 119, 5128–5141,
1210 doi:10.1002/2014JD021489, 2014a.
- 1211 Brocca, L., Hasenauer, S., Lacava, T., Melone, F., Moramarco, T., Wagner, W., Dorigo, W.,
1212 Matgen, P., Martínez-Fernández, J., Llorens, P., Latron, J., Martin, C., and Bittelli, M.:

1213 Soil moisture estimation through ASCAT and AMSR-E: An intercomparison and
 1214 validation study across Europe, *Remote Sens. Environ.*, 115(12), 3390–3408, doi:
 1215 10.1016/j.rse.2011.08.003, 2011.

1216 Brocca, L., Melone, F., Moramarco, T., and Morbidelli, R.: Spatial–temporal variability of
 1217 soil moisture and its estimation across scales, *Water Resour. Res.*, 46, W02516,
 1218 doi:10.1029/2009WR008016, 2010.

1219 [Brocca, L., Zucco, G., Mittelbach, H., Moramarco, T., and Seneviratne, S. I.: Absolute versus](#)
 1220 [temporal anomaly and percent of saturation soil moisture spatial variability for six](#)
 1221 [networks worldwide, *Water Resour. Res.*, 50, 5560–5576, doi:10.1002/2014WR015684,](#)
 1222 [2014b.](#)

1223 [Brocca, L., Melone, F., Moramarco, T., Wagner, W., and Albergel, C.: Chapter 17: Scaling](#)
 1224 [and filtering approaches for the use of satellite soil moisture observations. In G. P.](#)
 1225 [Petropoulos \(Ed.\), *Remote sensing of land surface turbulent fluxes and soil surface*](#)
 1226 [moisture content: State of the art \(pp. 562\), Taylor & Francis, 2013.](#)

1227 Chen, Y., Yang, K., He, J., Qin, J., Shi, J., Du, J., and He, Q.: Improving land surface
 1228 temperature modeling for dryland of China, *J. Geophys. Res.*, 116, D20104,
 1229 doi:10.1029/2011JD015921, 2011.

1230 Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogee, J., Allard, V., Aubinet, M.,
 1231 Buchmann, N., Bernhofer, C., Carrara, A., Chevallier, F., De Noblet, N., Friend, A. D.,
 1232 Friedlingstein, P., Grunwald, T., Heinesch, B., Keronen, P., Knohl, A., Krinner, G.,
 1233 Loustau, D., Manca, G., Matteucci, G., Miglietta, F., Ourcival, J. M., Papale, D.,
 1234 Pilegaard, K., Rambal, S., Seufert, G., Soussana, J. F., Sanz, M. J., Schulze, E. D., Vesala,
 1235 T., and Valentini, R.: Europe-wide reduction in primary productivity caused by the heat
 1236 and drought in 2003, *Nature*, 437, 529–533, doi:10.1038/nature03972, 2005.

1237 Crow, W. T., Berg, A. A., Cosh, M. H., Loew, A., Mohanty, B. P., Panciera, R., de Rosnay,

1238 P., Ryu, D., and Walker, J. P.: Upscaling sparse ground-based soil moisture observations
1239 for the validation of coarse-resolution satellite soil moisture products, *Rev. Geophys.*, 50,
1240 RG2002, doi:10.1029/2011RG000372, 2012.

1241 [de Rosnay P., Drusch, M., Vasiljevic, D., Balsamo, G., Albergel, C., and Isaksen, L.: A](#)
1242 [simplified Extended Kalman Filter for the global operational soil moisture analysis at](#)
1243 [ECMWF, Q. J. R. Meteorol. Soc., 139\(674\), 1199–1213, doi: 10.1002/qj.2023, 2013.](#)

1244 Dai, A., Trenberth, K. E., and Qian, T. T.: A global dataset of Palmer Drought Severity Index
1245 for 1870–2002: Relationship with soil moisture and effects of surface warming, *J.*
1246 *Hydrometeor.*, 5, 1117–1130, doi:10.1175/JHM-386.1, 2004.

1247 [Dharssi, I., Bovis, K. J., Macpherson, B., and Jones, C. P.: Operational assimilation of](#)
1248 [ASCAT surface soil wetness at the Met Office, Hydrol. Earth Syst. Sci., 15, 2729–2746,](#)
1249 [2011.](#)

1250 Dirmeyer, P. A., Gao, X., Zhao, M., Gao, Z., Oki, T., and Hanasaki, N.: GSWP-2:
1251 Multimodel analysis and implications for our perception of the land surface, *Bull. Amer.*
1252 *Meteor. Soc.*, 87, 1381–1397, 2006.

1253 Dorigo, W., de Jeu, R., Chung, D., Parinussa, R., Liu, Y., Wagner, W., and Fernandez, D.:
1254 Evaluating global trends (1988–2010) in homogenized remotely sensed surface soil
1255 moisture, *Geophys. Res. Lett.*, 39, L18405, doi: 10.1029/2012GL052988, 2012.

1256 Dorigo, W. A., Gruber, A., ~~De~~ Jeu, R. A. M., Wagner, W., Stacke, T., Loew, A., Albergel, C.,
1257 Brocca, L., Chung, D., Parinussa, R. ~~M.~~, and Kidd, R.: Evaluation of the ESA CCI soil
1258 moisture product using ground-based observations, *Remote Sens. Environ.*, ~~162~~, 380–395,
1259 ~~2015~~.

1260 [Dorigo, W. A., Scipal, K., Parinussa, R. M., Liu, Y. Y., Wagner, W., de Jeu, R. A.M., and](#)
1261 [Naeimi, V.: Error characterisation of global active and passive microwave soil moisture](#)
1262 [data sets, Hydrol. Earth Syst. Sci., 14, 2605–2616, 2010.](#)

binghaojia 15/6/29 4:24 PM

已删除: de

binghaojia 15/6/29 4:25 PM

已删除: doi:10.1016/j.rse.2014.07.023

binghaojia 15/6/29 4:25 PM

已删除: 2014

1266 | Draper, C. S., Walker, J. P., Steinle, P. J., de Jeu, R. A. M., and Holmes, T. R. H.: An
1267 | evaluation of AMSR-E derived soil moisture over Australia, Remote Sens. Environ., 113,
1268 | 703–710, 2009.

1269 | [Entin, J. K., Robock, A., Vinnikov, K. Y., Hollinger, S., Liu, E. S., and Namkhai, A.:
1270 | Temporal and spatial scales of observed soil moisture variations in the extratropics, J.
1271 | Geophys. Res., 105, 11865–11877, 2000.](#)

1272 | Gao, H., Wood, E. F., Jackson, T. J., Drusch, M., and Bindlish, R.: Using TRMM/TMI to
1273 | retrieve surface soil moisture over the southern United States from 1998 to 2002, J.
1274 | Hydrometeor., 7, 23–38, 2006.

1275 | Gao, X., and Dirmeyer, P. A.: A multimodel analysis, validation and transferability study of
1276 | global soil wetness products, J. Hydrometeor., 7, 1218–1236, 2006.

1277 | [Gruber, A., Dorigo, W. A., Zwieback, S., Xaver, A. and Wagner, W.: Characterizing
1278 | coarse-scale representativeness of in-situ soil moisture measurements from the
1279 | International Soil Moisture Network, Vadose Zone J., 12 \(2\), doi:10.2136/vzj2012.0170,
1280 | 2013.](#)

1281 | Gruhier, C., de Rosnay, P., Hasenauer, S., Holmes, T., de Jeu, R., Kerr, Y., Mougin, E.,
1282 | Njoku, E., Timouk, F., Wagner, W., and Zribi, M.: Soil moisture active and passive
1283 | microwave products: Intercomparison and evaluation over a Sahelian site, Hydrol. Earth
1284 | Syst. Sci., 14, 141–156, 2010.

1285 | Jackson, T. J., and Hsu, A. Y.: Soil moisture and TRMM microwave imager relationships in
1286 | the Southern Great Plains 1999 (SGP99) experiment, IEEE Trans. Geosci. Remote Sens.,
1287 | 39, 1632–1642, 2001.

1288 | Jia, B., Tian, X., Xie, Z., Liu, J., and Shi, C.: Assimilation of microwave brightness
1289 | temperature in a land data assimilation system with multi-observation operators, J.
1290 | Geophys. Res., 118, doi:10.1002/jgrd.50377, 2013.

binghaojia 15/7/28 8:53 AM
已设置格式: 字体:(中文) 宋体

1291 Kerr, Y. H., Waldteufel, P., Richaume, P., Wigneron, J. P., Ferrazzoli, P., Mahmoodi, A., Al
 1292 Bitar, A., Cabot, F., Gruhier, C., Juglea, S. E., Leroux, D., Mialon, A., and Delwart, S.:
 1293 The SMOS soil moisture retrieval algorithm, *IEEE Trans. Geosci. Remote Sens.*, 50,
 1294 1384–1403, 2012.
 1295 [Lai, X., Wen, J., Cen, S., Huang, X., Tian, H., and Shi, X.: Spatial and temporal soil moisture](#)
 1296 [variations over China from simulations and observations, *Adv. Meteorol.*, 2015, 529825,](#)
 1297 [2015.](#)
 1298 Lawrence, P. J., and Chase, T. N.: Representing a new MODIS consistent land surface in the
 1299 Community Land Model (CLM 3.0), *J. Geophys. Res.*, 112, G01023,
 1300 doi:10.1029/2006JG000168, 2007.
 1301 Li, H. B., Robock, A., Liu, S. X., Mo, X. G., and Viterbo, P.: Evaluation of reanalysis soil
 1302 moisture simulations using updated Chinese soil moisture observations, *J. Hydrometeor.*,
 1303 6, 180–193, 2005.
 1304 Liu, J. G., and Xie, Z. H.: Improving simulation of soil moisture in China using a multiple
 1305 meteorological forcing ensemble approach, *Hydrol. Earth Syst. Sci.*, 17, 3355–3369,
 1306 doi:10.5194/hess-17-3355-2013, 2013.
 1307 Liu, Y. Y., Dorigo, W. A., Parinussa, R., de Jeu, R. A. M., Wagner, W., McCabe, M. F.,
 1308 Evans, J. P., and van Dijk, A. I. J. M.: Trend-preserving blending of passive and active
 1309 microwave soil moisture retrievals, *Remote Sens. Environ.*, 123, 280–297, 2012.
 1310 Liu, Y. Y., Parinussa, R., Dorigo, W. A., De Jeu, R. A.M., Wagner, W., van Dijk, A. I. J. M.,
 1311 McCabe, M. F., and Evans, J. P.: Developing an improved soil moisture dataset by
 1312 blending passive and active microwave satellite-based retrievals, *Hydrol. Earth Syst. Sci.*,
 1313 15, 425–436, doi:10.5194/hess-15-425-2011, 2011.
 1314 Loew, A., Stacke, T., Dorigo, W., de Jeu, R., and Hagemann, S.: Potential and limitations of
 1315 multidecadal satellite soil moisture observations for selected climate model evaluation

1316 studies, *Hydrol. Earth Syst. Sci.*, 17, 3523–3542, 2013.

1317 Njoku, E. G., Jackson, T. L., Lakshmi, V., Chan, T., and Nghiem, S. V.: Soil moisture
 1318 retrieval from AMSR-E, *IEEE Trans. Geosci. Remote Sens.*, 41, 215–229, 2003.

1319 Oleson, K. W., Lawrence, D. M., Bonan, G. B., Drewniak, B., Huang, M., Koven, C. D.,
 1320 Levis, S., Li, F., Riley, W. J., Subin, Z. M., Swenson, S. C., Thornton, P. E., Bozbiyik, A.,
 1321 Fisher, R., Kluzek, E., Lamarque, J.-F., Lawrence, P. J., Leung, L. R., Lipscomb, W.,
 1322 Muszala, S., Ricciuto, D. M., Sacks, W., Sun, Y., Tang, J., and Yang, Z.-L.: Technical
 1323 description of version 4.5 of the community land model (CLM), NCAR Technical Note
 1324 NCAR/TN-503+STR, National Center for Atmospheric Research, Boulder, CO, 420pp,
 1325 2013.

1326 Owe, M., de Jeu, R., and Holmes, T.: Multisensor historical climatology of satellite-derived
 1327 global land surface moisture, *J. Geophys. Res.*, 113, F01002, doi:10.1029/2007JF000769,
 1328 2008.

1329 Parinussa, R. M., Holmes, T. R. H., and de Jeu, R. A. M.: Soil moisture retrievals from the
 1330 WindSat polarimetric microwave radiometer. *IEEE T. Geosci. Remote Sens.*, 50,
 1331 2683–2694, doi:10.1109/TGRS.2011.2174643, 2012.

1332 Parinussa, R., Holmes, T., Wanders, N., Dorigo, W., and de Jeu, R.: A preliminary study
 1333 toward consistent soil moisture from AMSR2, *J. Hydrometeor.*, 16, 932–947,
 1334 doi:10.1175/JHM-D-13-0200.1, 2015.

1335 Pinker, R. T., and Laszlo, I.: Modeling surface solar irradiance for satellite applications on a
 1336 global scale, *Journal of Applied Meteorology*, 31, 194–211, 1992.

1337 Rodell, M., Houser, P. R., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C. J., Arsenault,
 1338 K., Cosgrove, B., Radakovich, J., Bosilovich, M., Entin, J. K., Walker, J. P., Lohmann, D.,
 1339 and Toll, D.: The Global Land Data Assimilation System, *Bull. Amer. Meteor. Soc.*, 85,
 1340 381–394, 2004.

1341 Schmutge, T. J.: Remote sensing of soil moisture: recent advances, *IEEE T. Geosci. Remote*
 1342 *Sens.*, 21, 336–344, 1983.

1343 Scipal, K., Wagner, W., Trommler, M., and Naumann, K.: The global soil moisture archive
 1344 1992–2000 from ERS scatterometer data: first results, in: *Proceedings of the IEEE*
 1345 *International Geoscience and Remote Sensing Symposium (IGARSS)*, 24–26 June,
 1346 Toronto, Canada, 3, 1399–1401, 2002.

1347 Szczypka, C., Calvet, J. C., Maignan, F., Dorigo, W., Baret, F., and Ciais, P.: Suitability of
 1348 modelled and remotely sensed essential climate variables for monitoring
 1349 Euro-Mediterranean droughts, *Geoscientific Model Development*, 7, 931–946, 2014.

1350 Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *J.*
 1351 *Geophys. Res.*, 106, 7183–7192, 2001.

1352 Wagner, W., Dorigo, W., de Jeu, R., Fernandez, D., Benveniste, J., Haas, E., and Ertl, M.:
 1353 Fusion of active and passive microwave observations to create an Essential Climate
 1354 Variable data record on soil moisture, *ISPRS Annals of the Photogrammetry, Remote*
 1355 *Sensing and Spatial Information Sciences (ISPRS Annals)*, Volume I-7, XXII ISPRS
 1356 Congress, Melbourne, Australia, 25 August–1 September 2012, 315–321, 2012.

1357 Wang, A., Lettenmaier, D. P., and Sheffield, J.: Soil moisture drought in China, 1950–2006, *J.*
 1358 *Climate*, 24, 3257–3271, doi: 10.1175/2011JCLI3733.1, 2011.

1359 Wang, A., and Zeng, X.: Sensitivities of terrestrial water cycle simulations to the variations
 1360 of precipitation and air temperature in China, *J. Geophys. Res.*, 116, D02107,
 1361 doi:10.1029/2010JD014659, 2011.

1362 Yang, K., He, J., Tang, W. J., Qin, J., and Cheng, C. C. K.: On downward shortwave and
 1363 longwave radiations over high altitude regions: Observation and modeling in the Tibetan
 1364 Plateau, *Agr. Forest Meteorol.*, 150, 38–46, 2010.

1365 Zhu, Y. F.: The regional division of dryness/wetness over Eastern China and variations of

1366 dryness/wetness in Northern China during the last 530 years (in Chinese), *Acta Geogr.*
1367 *Sin.*, 58, supplement, 100–107, 2003.
1368

1369 | **Table 1.** Locations of the eight sub-regions in China.

Identification	Region Name	Location	Number of Observational Stations
China I	northeast China	120–135 °E, 40–50 °N	61 (40)
China II	northern North China	110–120 °E, 40–45 °N	15 (5)
China III	southern North China	110–120 °E, 34–40 °N	57 (47)
China IV	central and lower Yangtze River Basin	110–122 °E, 30–34 °N	28 (18)
China V	eastern northwest China	95–110 °E, 34–42 °N	77 (53)
China VI	western northwest China	80–95 °E, 34–50 °N	20 (8)
China VII	northern southwest China	100–110 °E, 28–34 °N	22 (12)
China VIII	southern southwest China	100–110 °E, 20–28 °N	9 (6)

1370 | ^aThe values in the brackets represent the number of sites having significant ($p < 0.05$)
1371 | positive Spearman’s correlations for both the ESA CCI SM product and CLM4.5 simulation.
1372 |

binghaojia 15/7/13 9:14 PM
已删除: 62

binghaojia 15/7/13 9:14 PM
已删除: 58

1375 **Table 2.** Averaged performance metrics and standard deviations of the ESA CCI SM product
1376 and CLM4.5 model for all available in situ sites. (Stations: Number of stations; N: Averaged
1377 value of number of valid measurements per station; ubRMSD: unbiased root mean square
1378 difference; R_{sp} : Spearman correlation coefficient).

	<u>ESA CCI SM</u>	CLM4.5
Stations	299 (212) ^a	306 (253)
N	124 (127)	126 (128)
<u>ub</u> RMSD (m ³ m ⁻³)	<u>0.053±0.017</u> (0.050±0.014)	0.050±0.017 (0.048±0.015)
R_{sp}	0.27 ± 0.21 (0.37 ± 0.14)	0.35 ± 0.21 (0.42 ± 0.14)

1379 ^aThe values in the brackets represent the average values from the sites having significant ($p <$
1380 0.05) positive Spearman correlations.

- binghaojia 15/6/29 2:27 PM
已删除: ECV-SM
- Binghao Jia 15/8/21 4:04 PM
已删除: over
- binghaojia 15/7/21 10:55 AM
已删除: Number
- binghaojia 15/7/21 10:34 AM
已删除: Bias: EVC-SM or CLM4.5 minus measurements;
- binghaojia 15/7/21 10:35 AM
已删除: ; E: centered normalized root mean square difference
- binghaojia 15/6/29 2:27 PM
已删除: ECV-SM
- binghaojia 15/7/21 10:31 AM
已删除: 301
- binghaojia 15/7/13 9:14 PM
已删除: 308
- binghaojia 15/7/21 11:37 AM
已删除: 37373
- binghaojia 15/7/21 11:28 AM
已删除: 38957
- binghaojia 15/7/21 10:33 AM
已删除: BIAS (m³ m⁻³) ... [42]
- binghaojia 15/7/21 12:51 PM
已设置格式: 左
- binghaojia 15/7/21 12:49 PM
已删除: 0.075 ± 0.034
- binghaojia 15/7/21 12:50 PM
已删除: 0.090 ± 0.041
- binghaojia 15/7/21 10:34 AM
已删除: R ... [43]
- binghaojia 15/7/21 10:33 AM
已删除: 1.13 ± 0.32
- binghaojia 15/7/21 10:33 AM
已删除: 0.93 ± 0.20
- binghaojia 15/7/21 10:51 AM
已删除: correlation coefficients
- Binghao Jia 15/8/21 4:04 PM
已删除: with
- binghaojia 15/7/21 10:51 AM
已删除: (p = 0.05)

Figures

Fig. 1 Locations of the 306 in situ soil moisture measurement sites (red dots) in China. Also shown (blue boxes) are the eight sub-regions defined in Table 1.

Fig. 2 Fraction of days with valid observations, expressed as the number of days with observations divided by the total number of days per period, split into six different merging periods: (a) 1 January 1979–31 August 1987, (b) 1 September 1987–30 June 1991, (c) 1 July 1991–31 December 1997, (d) 1 January 1998–30 June 2002, (e) 1 July 2002–31 December 2006, (f) 1 January 2007–30 September 2011, (g) 1 October 2011–30 June 2012, (h) 1 July 2012–31 December 2013; and (i) for the total period 1 January 1979–31 December 2013.

Fig. 3 Spearman rank correlation coefficients (R_{sp}) between the in situ measurements and (a) ESA CCI SM and (b) CLM4.5 from 1993 to 2012 at 306 sites in China. Open circles represent statistically significant Spearman correlation ($p < 0.05$) while closed circles are not significant.

Fig. 4 Unbiased root mean square differences (ubRMSDs) between the in situ measurements and (a) ESA CCI SM and (b) CLM4.5 from 1993 to 2012 at 306 sites in China.

Fig. 5 Statistical scores, including ubRMSD (5a and 5c) and R_{sp} (5b and 5d), for the ESA CCI SM product (red) and CLM4.5 model (blue) between 1993 and 2012 in the eight studied sub-regions of China using two evaluation strategies (S1 and S2). S1: average values of statistical metrics for the stations over each region. S2: statistical metrics with in situ observations for the soil moisture time series averaged at the stations of each region.

Fig. 6 Multi-year (1993–2012) mean monthly volumetric soil moisture (SM, $m^3 m^{-3}$) for the period between March and October as described by in situ observations, the ESA CCI SM product, and CLM4.5 simulation in the eight studied sub-regions of China.

Fig. 7 Comparison of the anomaly time series of monthly soil moisture derived from the ESA CCI SM product, CLM4.5 simulation, and in situ observations between 1993 and 2012 for

binghaojia 15/7/24 4:57 PM

已删除: Table 3. The root mean square differences of soil moisture anomalies for the ECV-SM and CLM4.5 against in situ measurements over eight sub-regions (Table 1). Unit is $m^3 m^{-3}$ [44]

binghaojia 15/7/13 9:14 PM

已删除: 308

Binghao Jia 15/8/21 4:08 PM

已删除: up according

binghaojia 15/7/30 11:16 AM

已删除: The biases (top, the ECV-SM or CLM4.5 minus in situ measurements) and root mean square differences (RMSDs, bottom) between the ECV-SM product or CLM4.5 simulation and ground-based measurements from 1993 to 2012 at 308 sites in China

Binghao Jia 15/8/21 4:12 PM

已删除: The same as Fig. 4 3 but for the unbiased root mean square differences (ubRMSDs) correlation coefficients R (top) and the centered normalized root mean square difference E (bottom).

binghaojia 15/7/30 3:32 PM

已删除: Statistical scores for the ECV-SM product and CLM4.5 simulation between 1993 and 2012 in the eight sub-regions of China defined in Table 1: (a) BIAS; (b) RMSD; (c) correlation coefficients (R)

Binghao Jia 15/8/21 4:29 PM

已删除: from the

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

binghaojia 15/7/30 3:36 PM

已删除: product

binghaojia 15/7/30 3:36 PM

已删除: simulation

Binghao Jia 15/8/21 4:30 PM

已删除: defined in Table 1

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

binghaojia 15/7/30 3:38 PM

已删除: product

binghaojia 15/7/30 3:38 PM

已删除: simulation

the eight studied sub-regions of China.

Fig. 8 Spearman correlations (R_{sp}) between the anomalies (seasonal cycle removed) of the
ESA CCI SM product (red) or CLM4.5 model (blue) and those of in situ observations
between 1993 and 2012 in the eight studied sub-regions of China using two evaluation
strategies (S1 and S2).

Fig. 9 Averaged $ubRMSD$ and R_{sp} values of the ESA CCI SM product against in situ
observations for five blended time periods. Red bars represent the sites with significant R_{sp}
 values ($p \leq 0.05$) for all periods (1993–2012) while blue bars consider sites with significant
 R_{sp} values ($p \leq 0.05$) for each blended period. Blend 3: July 1991–December 1997; Blend 4:
January 1998–June 2002; Blend 5: July 2002–December 2006; Blend 6: January
2007–September 2011; and Blend 7–8: October 2011–December 2012.

Fig. 10 Evaluation of the ESA CCI SM product and CLM4.5 simulation against in situ
observations over different types of soil moisture stations. "All" is calculated using all valid
sites (200) with significant ($p < 0.05$) positive Spearman correlation coefficients for both the
ESA CCI SM product and CLM4.5 simulation. "Rain" represents the sites (101) located in
the same 0.25° grid box as the in situ precipitation observations; "No Rain" represents the
remaining sites (99). Presented are the average values, the standard deviations (as indicated
by the box), and the maximum and minimum values (whiskers).

Fig. 11 Average statistical metrics: (a) $ubRMSD$, and (b) Spearman correlation coefficients
(R_{sp}) between the ESA CCI SM product (red) or in situ observations (OBS, blue) and
CLM4.5 simulation at different soil depths. "L1", "L1–2", "L1–2–3", "2cm", "5cm", and
"10cm" represent the first layer (0–1.75 cm), uppermost two layers (0–1.75, and 1.75–4.51
cm), and uppermost three layers (0–1.75, 1.75–4.51, and 4.51–9.06 cm), 0–2 cm, 0–5 cm,
and 0–10 cm, respectively. The average values of CLM4.5 were computed by using each soil
layer thickness as the weight; the CLM4.5 simulations were scaled to the dynamic range of

Binghao Jia 15/8/21 4:31 PM

已删除: (Table 1)

Binghao Jia 15/8/21 4:28 PM

已删除: The same as Figs. 5b and d but for the anomalies Taylor diagram summarizing the comparison results of (a) the original soil moisture values and (b) soil moisture anomalies between the ECV-SM, CLM4.5 and in situ measurements in the eight sub-regions of China defined in Table 1 between 1993 and 2012.

binghaojia 15/7/30 4:11 PM

已删除: correlation coefficients

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

Binghao Jia 15/8/21 4:31 PM

已删除: with

binghaojia 15/7/30 4:12 PM

已删除: data

Binghao Jia 15/8/21 4:32 PM

已删除: the

Binghao Jia 15/8/21 4:32 PM


已删除: defined in Section 2.2

Binghao Jia 15/8/21 4:32 PM

已删除: =

1501 | [the in situ observations or ESA CCI SM dataset using a linear rescaling method \(Eq. 1\) prior](#)
1502 | [to computing the ubRMSD values.](#)
1503


1504



1505

1506 **Fig. 1** Locations of the 306 in situ soil moisture measurement sites (red dots) in China. Also
1507 shown (blue boxes) are the eight sub-regions defined in Table 1.

1508






Fig. 2 Fraction of days with valid observations, expressed as the number of days with observations divided by the total number of days per period, split into six different merging periods: (a) 1 January 1979–31 August 1987, (b) 1 September 1987–30 June 1991, (c) 1 July 1991–31 December 1997, (d) 1 January 1998–30 June 2002, (e) 1 July 2002–31 December 2006, (f) 1 January 2007–30 September 2011, (g) 1 October 2011–30 June 2012, (h) 1 July 2012–31 December 2013; and (i) for the total period 1 January 1979–31 December 2013.

Binghao Jia 15/8/21 4:09 PM

已删除: up according





Fig. 3 Spearman rank correlation coefficients (R_{sp}) between the in situ measurements and (a) ESA CCI SM and (b) CLM4.5 from 1993 to 2012 at 306 sites in China. Open circles represent statistically significant Spearman correlation ($p < 0.05$) while closed circles are not significant.

binghaojia 15/7/21 2:12 PM

已删除:

binghaojia 15/7/20 10:26 PM



已删除:

binghaojia 15/7/20 10:22 PM

已删除: The biases (top, the EVC-SM or CLM4.5 minus in situ measurements) and root mean square differences (RMSDs, bottom) between the ECV-SM product or CLM4.5 simulation and ground-based measurements from 1993 to 2012 at 308 sites in China.





Fig. 4 Unbiased root mean square differences (ubRMSDs) between the in situ measurements and (a) ESA CCI SM and (b) CLM4.5 from 1993 to 2012 at 306 sites in China. Open circles represent statistically significant Spearman correlation ($p < 0.05$) while closed circles are not significant.



binghaojia 15/7/21 2:14 PM
已删除: correlation coefficients R (top) and the centered normalized root mean square difference E (bottom)





Fig. 5 Statistical scores, including ubRMSD (5a and 5c) and R_{sp} (5b and 5d), for the ESA CCI SM product (red) and CLM4.5 model (blue) between 1993 and 2012 in the eight studied sub-regions of China using two evaluation strategies (S1 and S2). S1: average values of statistical metrics for the stations over each region. S2: statistical metrics with in situ observations for the soil moisture time series averaged at the stations of each region.

binghaojia 15/7/30 11:16 AM



已删除:

Binghao Jia 15/8/21 4:18 PM

已删除: Statistical scores, including ubRMSD (left) and R_{sp} (right), for the ECV-SMESA CCI SM (red) product and CLM4.5 (blue) simulation between 1993 and 2012 in the eight sub-regions of China defined in Table 1 using two evaluation strategies.: S1 is the average values of statistical metrics for the stations over each region, while S2 representing the statistical metrics with in situ observations for the soil moisture time series averaged at the stations of each region(a) BIAS; (b) RMSD; (c) correlation coefficients (R).





Fig. 6 Multi-year (1993–2012) mean monthly volumetric soil moisture (SM, $\text{m}^3 \text{m}^{-3}$) for the period between March and October [as described by](#) in situ observations, [the ESA CCI SM product](#), and [the CLM4.5 simulation](#) in the eight studied sub-regions of China.



已删除:

Binghao Jia 15/8/21 4:19 PM

已删除: from the

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

binghaojia 15/7/30 3:36 PM

已删除: product

binghaojia 15/7/30 3:36 PM

已删除: simulation

Binghao Jia 15/8/21 4:20 PM

已删除: defined in Table 1




Fig. 7 Comparison of the anomaly time series of monthly soil moisture derived from the [ESA CCI SM product](#), the [CLM4.5 simulation](#), and in situ observations between 1993 and 2012 for the eight [studied](#) sub-regions of China.

binghaojia 15/7/30 3:37 PM

(a) China I —OBS

SM (m³ m⁻³)

1993 1997 2001

(c) China III

SM (m³ m⁻³)

1993 1997 2001

(e) China V

SM (m³ m⁻³)

1993 1997 2001

(g) China VII

SM (m³ m⁻³)

1993 1997 2001

已删除:

binghaojia 15/6/29 2:27 PM

已删除: ECV-SM

binghaojia 15/7/30 3:38 PM

已删除: product

binghaojia 15/7/30 3:38 PM

已删除: simulation

Binghao Jia 15/8/21 4:21 PM

已删除: (Table 1)





Fig. 8 Spearman correlations (R'_{sp}) between the anomalies (seasonal cycle removed) of the ESA CCI SM product (red) or CLM4.5 model (blue) and those of in situ observations between 1993 and 2012 in the eight studied sub-regions of China using two evaluation strategies (S1 and S2).



Binghao Jia 15/8/21 4:24 PM
 已删除: The same as Figs. 5b and d but for the anomalies. Taylor diagram summarizing the comparison results of (a) the original soil moisture values and (b) soil moisture anomalies between the ECV-SM, CLM4.5 and in situ measurements in the eight sub-regions of China defined in Table 1 between 1993 and 2012.





Fig. 9 Averaged $ubRMSD$ and R_{sp} values of the ESA CCI SM product against in situ observations for five blended time periods. Red bars represent the sites with significant R_{sp} values ($p \leq 0.05$) for all periods (1993–2012) while blue bars consider sites with significant R_{sp} values ($p \leq 0.05$) for each blended period. Blend 3: July 1991–December 1997; Blend 4: January 1998–June 2002; Blend 5: July 2002–December 2006; Blend 6: January 2007–September 2011; and Blend 7–8: October 2011–December 2012.



binghaojia 15/7/30 3:49 PM
已删除: correlation coefficients
binghaojia 15/6/29 2:27 PM
已删除: ECV-SM
Binghao Jia 15/8/21 4:34 PM
已删除: with
binghaojia 15/7/30 3:56 PM
已删除: data
Binghao Jia 15/8/21 4:34 PM
已删除: the
Binghao Jia 15/8/21 4:34 PM
已删除: defined in Section 2.2
Binghao Jia 15/8/21 4:34 PM
已删除: =
Binghao Jia 15/8/21 4:34 PM
已删除: =




Fig. 10 Evaluation of the ESA CCI SM product and CLM4.5 simulation against in situ observations over different types of soil moisture stations. "All" is calculated using all valid sites (200) with significant ($p < 0.05$) positive Spearman correlation coefficients for both the ESA CCI SM product and CLM4.5 simulation. "Rain" represents the sites (101) located in the same 0.25° grid box as the in situ precipitation observations; "No Rain" represents the remaining sites (99). Presented are the average values, the standard deviations (as indicated by the box), and the maximum and minimum values (whiskers).

Binghao Jia 15/8/21 4:39 PM

已删除: Evaluation of the ESA CCI SM and CLM4.5 against in situ observations over different types of soil moisture stations. "All" is calculated using all valid sites (200) with significant ($p = 0.05$) positive Spearman correlation coefficients for both the ESA CCI SM and CLM4.5. "Rain" represents the sites (101) locating at the same 0.25° grid box as the in situ precipitation observations while "No Rain" for the other sites (99). Presented are the average values, the standard deviations (as indicated by the box), and the maximum and minimum values (whiskers).




Fig. 11 Average statistical metrics: (a) ubRMSD, and (b) Spearman correlation coefficients (R_{sp}) between the ESA CCI SM product (red) or in situ observations (OBS, blue) and CLM4.5 simulation at different soil depths. "L1", "L1-2", "L1-2-3", "2cm", "5cm", and "10cm" represent the first layer (0–1.75 cm), uppermost two layers (0–1.75, and 1.75–4.51 cm), and uppermost three layers (0–1.75, 1.75–4.51, and 4.51–9.06 cm), 0–2 cm, 0–5 cm, and 0–10 cm, respectively. The average values of CLM4.5 were computed by using each soil layer thickness as the weight; the CLM4.5 simulations were scaled to the dynamic range of the in situ observations or ESA CCI SM dataset using a linear rescaling method (Eq. 1) prior to computing the ubRMSD values.