

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

Spatial variation of soil physical properties in adjacent alluvial and colluvial soils under Ustic moisture regime

M. Sağlam¹, H. S. Öztürk², S. Erşahin³, and A. İ. Özkan²

¹Department of Soil Science, Ondokuz Mayıs University, Samsun, Turkey

²Department of Soil Science, Faculty of Agriculture, Ankara University, 06110 Diskapı, Ankara, Turkey

³Department of Forest Engineering, School of Forestry, Çankırı Karatekin University, Çankırı, Turkey

Received: 10 March 2011 – Accepted: 11 April 2011 – Published: 28 April 2011

Correspondence to: H. S. Öztürk (hozturk@agri.ankara.edu.tr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

4261

Abstract

Soils vary spatially due to differences in soil management and soil formation factors. The soil spatial variability is an important determinant of efficiency of farm inputs and yield. This study was carried out to identify and compare spatial variation of some soil physical properties by geostatistics in alluvial and adjacent colluvial soils formed under ustic moisture regime at Gökhöyük State Farm (1750 ha), Amasya, Turkey. Seventy four soil samples were collected on a regular grid (500 × 500-m) and additional 224 samples were collected on 28 500-m fine-transects, randomly superimposed between the nodes of grids. Semivariograms and corresponding kriging maps for soil texture, soil organic matter (SOM), bulk density (BD), saturated hydraulic conductivity (K_s), and available water content (AWC) were prepared. Statistical analyses were conducted separately for colluvial and alluvial sites as well as whole area. The soils in alluvial site is rich in clay with high BD and SOM, and low in K_s and AWC; and the soils in colluvial site was designated as low in K_s , SOM, and AWC and high in BD. All variables, except SOM, showed a strong spatial dependency. In general, nugget, sill and range values of most of the studied soil variables decreased from alluvial site to colluvial site. When local (alluvial and colluvial sites separately) and global (alluvial + colluvial) kriged maps for BD, AWC, and soil textural separates, use of global semivariograms (one semivariogram for entire study area) resulted in loss of some details in colluvial sites, suggesting that local semivariograms for alluvial and colluvial soils should be used in kriging predictions at the farm. The results had significant implications for water management as AWC was spatially associated to clay content in alluvial site and to clay and sand contents in colluvial site.

1 Introduction

Soil properties can vary depending on soil forming factors (time, parent material, topography, climate, organisms, and complex interactions among them) and differences in

4262

human activities. Heterogeneity is an inherent quality of soil that typifies its distribution in space (Júnior et al., 2006). Therefore, evaluation of soil spatial variability becomes an important issue in agricultural and environmental research.

5 Geostatistics enables us to describe spatial patterns by semivariograms and to predict the values of soil attributes at unsampled locations by a set of statistical tools (Vieira et al., 1983; Trangmar et al., 1985; Warrick et al., 1986). At present, soil science research largely relies on the use of geostatistics, which, together with classical statistics, constitutes an extraordinarily important tool for agronomy (Júnior et al., 2005).

10 Soils attributes can show a considerable heterogeneity (Tyler, 1985; Trangmar et al., 1987; Isaaks et al., 1989). Soil variability was widely studied in different contexts such as, soil topography (Miller et al., 1988; Iqbal et al., 2005), evaluation of soil fertility (Yasrebi et al., 2008; Štípek et al., 2002); planning and interpretation of field research (Wilding and Drees, 1986), and the study of the physical and chemical attributes of soil under different management regimes (Mzuku et al., 2005; Wei et al., 2008).

15 Despite the importance of soil texture and its relative ease of determination using conventional methods, soil maps are produced at large scales to adequately represent their spatial distribution. Quantitative information on soil surface texture would be extremely useful for modeling, planning, and managing the soils (Scull et al., 2005). 20 Duffera et al. (2007) report some soil properties including particle size distribution (soil texture), soil water content, plant available water and cone index shows horizontal spatial structure and captured by soil map units i.e. soil texture maps yet, on the other hand, some properties such as bulk density, total porosity and saturated hydraulic conductivity are not spatially correlated and unrelated to soil map units. They also suggested 25 use of soil map units that delineate boundaries for developing management zones for site-specific crop management

Soil characteristics can highly change by various factors. In this context we mainly work to figure out the effects of the topography on soil. Characterizing spatial variation of soil variables can provide important implications in water and nutrient management

4263

and fertilizer applications in agricultural production. The objectives of this study were to characterize spatial variation of soil physical properties in a large state farm covered by alluvial and colluvial soils with known long term management history. The results showed that the source of spatial variation were highly different in alluvial site from that 5 in colluvial site.

2 Materials and methods

2.1 Site description

10 The study was carried out in an approximately 1750 ha area at Gökhöyük State Farm (between $722^{\circ}24'35''$ – $722^{\circ}24'78''$ E and $449^{\circ}03'30''$ – $449^{\circ}06'02''$ N and with an elevation of 500 m) (Fig. 1), in Amasya province of Turkey. The farm is located in North of Çekerek Creek and consists of three physiographic units; highlands, formed on limestone and sandstone; colluvial soils formed on the sediments from the highlands; and alluvial soils, formed on the deposits of the Çekerek Creek. Only colluvial and alluvial areas were included by the study. Approximately 30% of study areas is alluvial and rest 15 is colluvial soils. The area has terrestrial climate with annual precipitation of 370 mm and average temperature of 13.9°C . Considerable amount of the precipitation falls in the autumn (September–December) and in the spring (April and May). The slope of the alluvial site changes mainly from 0 to 2% and the slopes are mainly oriented in the southwest. Slope and elevation gradually increases in the north.

2.2 Field history

20 In general, crops in the alluvial areas are irrigated, while dryland (rainfed) farming is practiced in the colluvial sites. Sunflower, vetch, corn, chickpeas, onions, sugar beets, and alfalfa are the major crops grown in alluvial soils, and wheat is the main crop in colluvial site. The main crop rotation systems consisted of wheat-sunflower,

GS+ (Version 5) were used in geostatistics module of ArcView. Each time, the estimates in kriging were controlled by the procedure cross-validation to determine the most proper number of neighboring data points to be used in the estimation (Vieira et al., 1981). A maximum of 15 and a minimum of 10 neighboring data points were used along with semivariograms in kriging estimation. The surface maps of estimates from point kriging and surface maps of error values were examined to assess the quality of the estimations.

3 Results

In contrast to sand and silt content, clay content in the alluvial area is higher. The particle size gradually changes from coarse to finer toward to the river, from the northwest to the southeast. In the colluvial site where slope changes from 2 to 4%, sand was the dominant particle size. High BD was coincided with high clay, and low sand and silt contents in the alluvial and the colluvial sites. Unexpectedly, organic matter was low in the areas with very high clay content. Clay can stabilize organic matter against degradation, but this low organic matter occurred in clay rich areas was attributed to the unfavorable conditions caused by high clay content that decreased organic matter addition to the soil. On the contrary, the reason for the high soil organic matter in coarse textured areas can be attributed to more available soil water, pore space, and the aeration capacity that resulted in SOM addition in greater amounts.

Descriptive statistics for soil variables in the study area are given in Table 1. Clay contents are generally high. On the other hand, a low mean value for sand content was found especially in the alluvial site. AWC was also low in both sites. High SOM was determined in alluvial soils, whereas, lower values were detected in colluvial soils. Clay, silt, and SOM contents were more varied in alluvial site while BD and K_s were more varied in colluvial site. Table 1 presents the parameters of semivariograms for selected variables for alluvial, colluvial, and whole (alluvial + colluvial) area. The range of BD increased from 414 m in colluvial area to 1018 m in alluvial area.

4267

Figure 2 displays different increases in semivariograms and maximum with increasing lag distances. The variograms for clay, SOM and K_s in alluvial area rise abruptly and reach to their highest value in a greater range comparing with those in colluvial area. The variograms show higher values of sill and range in alluvial site (except for the range of AWC) than in colluvial soils. This increase show a greater structured variance and a longer range in the later, which may indicate a continuous depositional event in colluvial site and relatively stable conditions in alluvial site

The kriged contour maps of clay, sand, SOM, BD, K_s , and AWC for the alluvial and colluvial sites are shown in Figs. 3 and 4. Kriged maps indicated soils with high clay and low sand and with high BD were found in the SW and in the center of the alluvial site, respectively. Kriged maps for K_s show that soils with high values located on the North in both sites.

The data from the whole study area (alluvial+colluvial) were analyzed together and results were shown in kriged maps (Fig. 5). From these maps, soils with the high clay were found in the center and the SW edge of the field, high sand in the north and high SOM mainly in the center and southern part of the field.

4 Discussion

The majority of the soil properties were slightly skewed with a coefficient of skewness < 0.5 . However, their distribution can be deemed normal. Although similar mean values occurred for clay content in alluvial and colluvial soils, clay was far more variable in the alluvial soils due to the nonuniform deposition of clay by Çekerek Creek. The relatively high standard deviation on alluvial site was also support this conclusion. While soil texture differed between the sites, the mean BD values of the sites were similar. This may be due to compaction of the clay soils in the alluvial site due to the field traffic on the farm and monoculture practiced (alfalfa and wheat). The reason of high K_s in alluvial soil is due to silty soils distributed in the northeast of the field. There is also high standard deviation in K_s in alluvial soils because of the high variation in

4268

soil texture over the field. Another possible reason of high K_s in alluvial soil would be intensive plant root growth despite to high clay. These plant roots may generate macro pore flow which is very important in vertical transport of water and chemicals in soils. AWC was low in both sites because of the high clay content in some localities and high silt content in others. Comparatively higher SOM in alluvial soils was attributed to irrigation that resulted in plant residue return to the soil in greater amounts than occurred in colluvial site.

That extremely low nugget ratio (0.03%) occurred for sand in colluvial site was attributed to its deposition controlled by slope steepness. Cambardella et al. (1994) interpreted spatial continuity according to degree of nugget ratio for the semivariograms. The variable BD was strongly spatially dependent in both sides. Strong spatial dependency of BD was attributed to strong association between BD and soil textural separates, which also exhibited a strong spatial dependency in the study area. Our calculations showed that, except K_s , all soil variables studied were strongly spatially dependent as their nugget ratio was less than 25%. Lower nugget effect with lower range values for BD found in the colluvial area may indicate the combined influence of soil texture, topography, and the land use on BD. Similar conclusion was made by Cambardella et al. (1994), who stated that strongly spatially dependent properties might be controlled by intrinsic variation in soil characteristics such as texture and mineralogy.

Values for sill and range of all the variables (except the range of AWC) were greater in alluvial site than in colluvial site. This difference showed a greater structural variance and a longer range in the former, which may indicate a more patchy distribution of these variables in colluvial site. Of the soil variables studied, the greatest variation occurred for K_s in both sides. A moderate nugget effect occurred for K_s indicated that considerable amount of this variation comes from processes such as short range variation and variance caused by laboratory procedures. Experiments have shown that K_s varies widely among materials with different particle-size distribution. Indeed, replicates of the same test in the same laboratory often yield highly dissimilar results (Arya et al., 2000; Mbonimpa et al., 2002). Therefore, the high nugget variance for K_s was

4269

expected on both sides. Semivariograms for SOM indicated strong spherical structure in both sides. Greater variation in SOM in alluvial area could be due to difference in addition of organic matter to soil due to differences in agricultural practices.

The Çekerek Creek deposited its sediments in the direction normal to its flow path, resulting in a deposition trend in decreasing sand content and increasing clay content away from stream bank. In contrast to colluvial area, some faint local trends occurred for BD in alluvial area (Fig. 3). These trends were attributed to compaction due to very high clay content combined with high trafficking conventional tillage. Kriging predicted surface maps for sand, clay and BD indicated similar spatial patterns in alluvial area. Also, semivariogram with a sill smaller than general variance indicate a strong global trend for these variables in alluvial site (Fig. 3). All these suggest that a strong spatial relation existed between BD, sand, and clay contents in alluvial site. However, no similar trends occurred among these variables in colluvial site (Fig. 4).

Some faint trends occurred for sand content in the alluvial area contrasting to colluvial area. Although sand distribution in the alluvial site was mainly in the direction of SW to NE, due to the deposition of the river sediments in the direction normal to its flow path, such a similar trend was not detected in the colluvial site. Also, no trends were detected for distribution of clay at either site. Kriged maps indicated that soils with high clay content in alluvial site occur in the SW and the center of the field, whereas, clay content in the colluvial site decrease from the river bank to the hillside in the north (Figs. 3 and 4). The kriged maps of clay and sand contents represent a reciprocal distribution in alluvial soils. However, no such distribution was observed for sand and clay content in colluvial site.

Soil organic matter content in alluvial soil is generally higher than in colluvial soils, and this was attributed to poor drainage conditions, high ground water table, and more rigorous plant growth in some localities. While faint local trends were detectable for SOM in alluvial area due to differences in cropping pattern and associated tillage, no trends was detected in colluvial area due to more uniform tillage and cropping pattern.

4270

- Júnior, V. V., Carvalho, M. P., Dafonte, J., Freddi, O. S., Va'zquez, E. V., and Ingaramo, O. E.: Spatial variability of soil water content and mechanical resistance of Brazilian ferralsol, *Soil Till. Res.*, 88, 166–177, 2005.
- Mbonimpa, M., Aubertin, M., Chapuis, R. P., and Bussière, B.: Practical pedotransfer functions for estimating the saturated hydraulic conductivity, *Geotech. Geol. Eng.*, 20, 235–259, 2002.
- 5 Miller, M. P., Singer, M. J., and Nielsen, D. R.: Spatial variability of wheat yield and soil properties on complex hills, *Soil Sci. Soc. Am. J.*, 52, 1133–1141, 1988.
- Myers, D. E.: Statistical models for multiple scaled analysis, in: *Scale in remote sensing and GIS*, edited by: Quattrochi, D. A. and Goodchild, M., Lewis Publishers, 273–294, 1997.
- 10 Mzuku, M., Khosla, R., Reich, R., Inman, D., Smith, F., and MacDonald, L.: Spatial Variability of Measured Soil Properties across Site-Specific Management Zones, *Soil Sci. Soc. Am. J.*, 69, 1572–1579, 2005.
- Nielsen, D. and Wendroth, O.: *Spatial and Temporal Statistics – Sampling Field Soils and Their Vegetation*, Catena-Verlag, 2003.
- 15 Rossi, R. E., Mulla, D. J., Journel, A. G., and Franz, E. H.: Geostatistical tools for modeling and interpreting ecological spatial dependence, *Ecol. Monogr.*, 62, 277–314, 1992.
- Scull, P., Okin, G., Chadwick, O. A., and Franklin, J. A.: Comparison of methods to predict soil surface texture in an alluvial basin, *Prof. Geogr.*, 57, 423–437, 2005.
- Štípek, K., Vaněk, V., Száková, J., Černý, J., and Šilha, J.: Temporal variability of available phosphorus, potassium and magnesium in arable soil, *Plant Soil Environ.*, 50, 547–551, 2004.
- 20 Trangmar, B. B., Yost, R. J., and Uehara, G.: Application of Geostatistics to spatial studies of soil properties, *Adv. Agron.*, 38, 65–91, 1985.
- Trangmar, B. B., Yost, R. J., Wade, M. K., Uehara, G., and Sudjadi, M.: Spatial variation of soil properties and rice yield on recently cleared land, *Soil Sci. Soc. Am. J.*, 51, 668–674, 1987.
- 25 Tyler, D. D.: Soil sampling in no-tillage cropping, *Better Crops Plant Food*, 69, 6–27, 1985.
- Vauclin, M., Viera, S. R., Vachaud, G., and Nielsen, D. R.: The use of co-kriging with limited field soil observations, *Soil Sci. Soc. Am. J.*, 47, 175–184, 1983.
- Vieira, S. R., Nielsen, D. R., and Biggar, J. W.: Spatial variability of field-measured infiltration rate, *Soil Sci. Soc. Am. J.*, 1040–1048, 1981.
- 30 Vieira, S. R., Hatfield, J. L., Nielsen, D. R., and Biggar, J. W.: Geostatistical theory and application to variability of some agronomical properties, *Hilgardia*, 51, 1–75, 1983.

4273

- Warrick, A. W., Myers, D. E., and Nielsen, D. R.: Geostatistical methods applied to soil science, in: *Methods of Soil Analysis, Part 1*, 2nd edition, edited by: Klute, A., Soil Science Society of America, Madison, 53–82, 1986.
- Webster, R.: Quantitative spatial analysis of soil in the field, *Adv. Soil Sci.*, 3, 1–70, 1985.
- 5 Webster, R.: Statistics to support soil research and their presentation, *Eur. J. Soil Sci.*, 52, 331–340, 2001.
- Wei, J. B., Xiao, D. N., Zeng, H., and Fu, Y. K.: Spatial variability of soil properties in relation to land use and topography in a typical small watershed of the black soil region, northeastern China, *Environ. Geol.*, 53, 1663–1672, 2008.
- 10 Wilding, L. P. and Drees, L. R.: Spatial Variability and Pedology, in: *Pedogenesis and Soil Taxonomy: Concepts and Interactions*, vol. 1, edited by: Wilding, L. P., Smeck, N. E., and Hall, G. F., Elsevier, Amsterdam, The Netherlands, 83–116, 1983.
- Yasrebi, J., Saffari, M., Fathi, H., Karimian, N., Emadi, M., and Baghernejad, M.: Spatial variability of soil fertility properties for precision agriculture in Southern Iran, *J. Appl. Sci.*, 8, 1642–1650, 2008.
- 15

4274




Fig. 2. Semivariogram models showing the spatial dependence of clay, SOM, K_s , AWC and BD for alluvial, collovial and whole area.

4277




Fig. 3. Kriged maps of clay, sand, soil organic matter, soil bulk density, saturated hydraulic conductivity and available water content for alluvial site.

4278




Fig. 4. Kriged maps of clay, sand, soil organic matter, soil bulk density, saturated hydraulic conductivity and available water content for Colluvial site.

4279




Fig. 5. Kriged maps of clay, sand, soil organic matter, soil bulk density, saturated hydraulic conductivity and available water content for whole site.

4280