

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

Hillslope characteristics as controls of subsurface flow variability

S. Bachmair and M. Weiler

Institute of Hydrology, University of Freiburg, Germany

Received: 13 May 2012 – Accepted: 19 May 2012 – Published: 1 June 2012

Correspondence to: S. Bachmair (sophie.bachmair@hydrology.uni-freiburg.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

HESSD

9, 6889–6934, 2012

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Hillslope hydrological dynamics, particularly subsurface flow (SSF), are highly variable and complex. A profound understanding of factors controlling this variability is needed. Therefore we investigated the relationship between variability of shallow water table dynamics and various hillslope characteristics. We ask whether measurable hillslope properties explain patterns of subsurface flow variability. To approach this question shallow water table dynamics of three adjacent large-scale hillslopes were monitored with high spatial and temporal resolution over 18 months. The hillslopes are similar in terms of topography and parent material, but different in vegetation cover (grassland, coniferous forest, and mixed forest). We expect vegetation to be an important driver of water table dynamics at our study site, especially given the minor differences in topography. Various hillslope properties were determined in the field and via GIS analysis: common topography descriptors, well depth, soil properties via slug tests, and several vegetation parameters. Response variables characterizing the water table response per well were calculated for different temporal scales (entire time series, seasonal scale, event scale). Partial correlation analysis and a Random Forest machine learning approach were carried out to assess the explainability of SSF variability by measurable hillslope characteristics. We found a complex interplay of predictors, yet soil properties and topography showed the highest single explanatory power. Surprisingly, vegetation characteristics played a minor role. Solely throughfall and canopy cover exerted a slightly stronger control, especially in summer. Most importantly, the examined hillslope characteristics explained only a small proportion of the observed SSF variability. Consequently there must be additional important drivers not represented by current measurement techniques of the hillslope configuration (e.g. bedrock properties, preferential pathways). We also found interesting differences in explainability of SSF variability among temporal scales and between both forested hillslopes and the grassland hillslope.

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Hydrological dynamics of hillslopes, particularly shallow subsurface flow (SSF), are highly complex and variable in space and time (Bachmair et al., 2012). Nonetheless, a solid understanding of hydrological dynamics and soil-vegetation-atmosphere interactions at the hillslope scale is necessary for predictions of ungauged hillslopes and catchments dominated by subsurface flow. In recent years a common theme has evolved focusing on the drivers or organizing principles of spatio-temporal variability of hydrologic fluxes at a range of spatial scales (McDonnell et al., 2007; Wagener et al., 2007; Sivapalan, 2005). At the hillslope scale, a number of studies addressed the issue of systematically investigating dominant controls of subsurface flow generation and their interrelations (Hopp and McDonnell, 2009; Hopp and McDonnell, 2011; Keim et al., 2006; Tromp-van Meerveld and McDonnell, 2006b; Gerrits et al., 2009).

The dominant controls of hillslope response to rainfall or snowmelt can be classified as static factors, at least in the sense of shorter time scales such as years (parent material, surface and bedrock topography), and dynamic factors (e.g. soil moisture, vegetation). Surface and bedrock topography clearly exert key controls, since they directly govern flow processes and also the evolution of dynamic factors such as soil properties, soil moisture, microclimate, and thus vegetation patterns. A number of studies assessed the effect of surface and bedrock terrain attributes on runoff generation (e.g. Freer et al., 2002; Troch et al., 2003; Aryal et al., 2002; Bogaart and Troch, 2006; Fujimoto et al., 2008; Berne et al., 2005). Bedrock microtopography (0.1–10 m) was found to create a threshold-like SSF response (bedrock depressions have to be filled up for hillslope response, “fill-and-spill-hypothesis” (Tromp-van Meerveld and McDonnell, 2006a; Tromp-van Meerveld and Weiler, 2008)). Bedrock permeability governs bedrock infiltration and exfiltration and thus influences SSF at the event scale and the long-term water balance of the hillslope and catchment (Tromp-van Meerveld et al., 2007; Kosugi et al., 2006; Graham et al., 2010).

HESSD

9, 6889–6934, 2012

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hillslope characteristicsS. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Regarding soil properties, the hillslope SSF response is controlled by various interactions between infiltration capacity, changes in hydraulic conductivity, presence of fragipans, thickness of the unsaturated zone, drainable porosity, moisture content, and vertical and lateral preferential flow features (macropores, soil cracks, root channels, animal burrows, high-permeability areas such as the soil skeleton, organic material etc.) (e.g. Hopp and McDonnell, 2009; Weiler and McDonnell, 2004; Zimmermann et al., 2006; Nieber et al., 2006; Anderson et al., 2009; Weiler and Naef, 2003; Tromp-van Meerveld et al., 2007). The significance of preferential pathways making up a large fraction of total SSF was shown by numerous experimental studies (Jones, 1997; Anderson et al., 2009; Noguchi et al., 1999; Tsuboyama et al., 1994; Holden and Burt, 2002; Uchida et al., 2002; Uchida et al., 2005; Kienzler and Naef, 2008). Preferential networks effectively drain the hillslope, increasing peak discharge and decreasing peak lag time of the storm hydrograph (Uchida et al., 2001; Weiler and McDonnell, 2007).

The influence of vegetation on hydrological dynamics is mostly assessed either by field experiments at the plot scale focusing on tree/stand structure and their effects on precipitation redistribution, or at the hillslope scale via virtual experiments simulating throughfall/stemflow variability and their effects on SSF (e.g. Keim et al., 2006; Gerrits et al., 2009; Hopp and McDonnell, 2011). Virtual experiments are numerical experiments with a model driven by collective field intelligence (Weiler and McDonnell, 2004). Hillslope-scale field experiments investigating vegetation effects are rare. Exceptions are recent sprinkler studies by Nordmann et al. (2009) and Jost et al. (2012), who showed the effect of different tree species on SSF generation and various interactions of vegetation with static factors, e.g. how roots channel water depending on subsurface conditions.

Numerous plot-scale experiments showed the prominent effect of vegetation on precipitation redistribution and small-scale variability of hydrologic fluxes. For instance, Liang et al. (2007) measured a rapid increase in water content and pore water pressure at the downslope side of a tree trunk growing on a steep hillslope, which they attribute to locally concentrated stemflow input. The effect of funnelled rainwater input

close to stems was also observed in other studies (e.g. Chang and Matzner, 2000; Staelens et al., 2008; Herwitz, 1986). Moreover, throughfall patterns were found to create distinct patterns of wet and dry spots that persist over time (e.g. Guswa and Spence, 2011; Zimmermann et al., 2009; Keim et al., 2005; Staelens et al., 2006).

Whether such small-scale vegetation effects influence SSF at the hillslope scale yet needs to be assessed in the field. Virtual experiments provide valuable insights, yet they always depend on the choice of the perceptual and numerical model.

To highlight the effect of different hillslope controls on SSF generation we monitored shallow water table dynamics of three adjacent, relatively large-scale (33 × 75 m) hillslopes, which are side-slopes of a small v-shaped, zero-order catchment. The hillslopes are similar in terms of topography and parent material, but differ in vegetation cover (grassland, coniferous forest, and mixed forest). The aim of this study is to explore the relationship between variability of SSF, monitored as shallow water table dynamics, and various hillslope characteristics, representing both static and dynamic controls.

The following research questions will be explored:

1. Can the spatial variability of SSF dynamics be explained by measurable hillslope characteristics? In other words, is there a direct link between patterns of water table fluctuations and hillslope characteristics?

A number of studies showed the influence of vegetation on soil moisture patterns and runoff generation processes, e.g. through spatially variable rainfall input due to throughfall and stemflow, transpiration, and different root systems. We thus hypothesize that vegetation is an important driver of spatially variable SSF dynamics at our study site, especially given the small differences in topography (predominantly planar hillslopes).

2. Are there differences in explainability of water table dynamics regarding different temporal scales, i.e. time periods of different length? Put differently, does the explanatory power of hillslope characteristics depend on the temporal scale of the response variable (e.g. water table response over more than a year, over one sea-

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



son, over a single rainfall event)? Are there differences in explainability of water table dynamics among seasons and events?

3. If we separately look at SSF dynamics at the grassland hillslope versus both forested hillslopes – are there differences in explainability of water table dynamics in terms of land use type?

2 Methods

2.1 Site description

To answer our research questions, shallow water table dynamics of three adjacent hillslopes were monitored with high spatial and temporal resolution (see Fig. 1). The hillslopes represent NW-facing side slopes of a small v-shaped, zero-order catchment at the foot of the Black Forest in south-western Germany (location of catchment outlet: 47.9570° N, 7.8378° E; catchment area: 0.21 km²; catchment elevation range: 340–585 m a.s.l.). Due to their spatial vicinity the hillslopes have similar planar topography, parent material, aspect, and steep slope (Fig. 1). Geology is crystalline bedrock overlain by periglacial drift cover (Geologisches Landesamt Baden-Württemberg, 1996). Cambisols have developed in the periglacial drift cover; soil texture is sandy loam (WaBoA, 2007). The main difference between hillslopes is vegetation cover (grassland, coniferous forest, and mixed forest).

Due to the periglacial drift cover there is no clear soil-bedrock-interface at the study site. Periglacial drift cover evolved from a combination of solifluction, cryoturbation and aeolian processes (Arno et al., 1998; Völkel et al., 2001). Stratigraphically, periglacial drift cover is comprised of three lithologic units: basal layer, intermediate layer, and upper layer. The basal layer is compacted; it shows high bulk density and slope-parallel alignment of clasts. In the intermediate layer the coarse fraction displays finer-sized clasts of varying orientation. The upper layer is the finest textured lithologic unit and has lower bulk density (Arno et al., 1998; Völkel et al., 2001). Generally, lateral subsurface

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



flow in periglacial drift cover was observed to occur at the interface of the basal layer and intermediate layer, but also at the interface of the intermediate and upper layer (Arno et al., 1998; Chiffard et al., 2008; Nordmann et al., 2009).

The climate is warm temperate “Cfb” (Koeppen classification). Mean annual precipitation and temperature are 970 mm and 11 °C, respectively (time period 2007–2011; data from nearby WBI weather station by state-run viticulture institute Freiburg). Mean monthly rainfall and evaporation is highest in the summer months May until July, when convective storms dominate (Morhard, unpublished). See also Fig. 3 for an overview on rainfall characteristics for different seasons. The small creek at the foot of the hillslopes carries water at all times, even under dry summer conditions (baseflow then smaller than 1 l s^{-1}).

At the mixed forest hillslope, a mix of European beech (*Fagus sylvatica*) and fir (*Abies alba*) dominates, with some sycamore maple (*Acer pseudoplatanus*), ash (*Fraxinus excelsior*), and spruce (*Picea abies*) in between (see Fig. 1). There is little understory vegetation and no deadwood on the forest floor, which is covered by a thick layer of beech leaves. At the coniferous hillslope primarily spruce and fir is found, interspersed with some deciduous trees at the lower part of the hillslope (sycamore maple and ash). The surface is covered by a needle layer, lots of deadwood, and understory vegetation. Tree age at both forested hillslopes is between 70 and 100 years. The grassland hillslope, which is under this vegetation cover for 200–300 years, is used for sporadic sheep grazing in the summer.

2.2 Field methods

To monitor the internal hillslope response to rainfall or snowmelt with high spatial and temporal resolution we installed 90 wells distributed over the three hillslopes. At each hillslope 30 wells were drilled, organized in three transects of 10 wells each, laid out perpendicular to the slope gradient along the contour in the lower, middle, and upper part of the hillslope (see Fig. 1). The distance between wells per transect is 3 m. The

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



distance between transects is 30 m, except for the upper transect of the mixed forest hillslope, which is only spaced 15 m apart due to an old forest road cutting the hillslope.

The wells were drilled with a hand-held, gasoline-powered breaker (Cobra Standard). The depth of each well depends on below ground conditions. We aimed to drill to a maximum depth of 2 m. Many wells are shallower, however, due to resistance in the periglacial drift cover or the bedrock. The majority of wells end in dense layers of periglacial drift cover, since the actual bedrock is mostly located far beneath the drift cover. A PVC pipe (4 cm diameter) perforated over the entire length was placed into each well. The perforated PVC pipe had been wrapped in geotextile prior to field installation to prevent fine material transport into the pipe. The geotextile also covers the lower end of the PVC pipe. Since the PVC pipe is not closed with a cap vertical drainage is possible. After inserting the PVC pipe into the well, bentonite clay pellets were pressed around the PVC casing at the soil surface to seal the well against preferential flow along the pipe.

Each well was equipped with an Odyssey Capacitance Water Level Recorder (Data Flow Systems, New Zealand). The measuring interval was set to 2 min but the loggers run in compressed logging mode (recording only if water level change $> \pm 5$ mm). After downloading the data roughly every 3 months, the capacitance probes were pulled out, cleaned, and the water table height manually measured with an electronic contact gauge to validate data. Manufacturer's specifications of probe accuracy are 5 mm. However, the validation yielded unsatisfactory results for wells with low water table (water table within the lower 13 cm of a well). Since there was no continuous bias for wells with low water table, water table dynamics in the lower 13 cm of each well are omitted from the analysis to exclude uncertain data (this includes the length of the 7.5 cm brass counter weight at the lower end of the Odyssey probe).

To monitor rainfall input and further meteorological variables a Davis weather station Vantage Pro 2 was installed (10 min measurement interval). To gain insight into the spatial variability of rainfall and throughfall, 66 rainfall totalizers were positioned (one

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



totalizator per well at both forested hillslopes, two totalizators per transect at the grassland hillslope). The totalizators were mounted to the upper end of the well casings.

For determining soil hydraulic properties, one slug test per well was carried out in December 2010. Prior to slug injection, the Odyssey capacitance probe was pulled out of the well and replaced by a pressure transducer (Diver Groundwater Datalogger, Schlumberger Water Services) due to higher accuracy and smaller logging interval of the pressure transducer (1 s interval). For a later correction of hydrostatic head with barometric head, a Baro-Diver (Schlumberger Water Services) was placed next to the well at the soil surface. A small-diameter PVC tube with attached packer was inserted into the well down to a depth of 50 cm above well bottom, where the packer was inflated. The slug tests thus only represent soil properties of the lower 50 cm of the well, which represents the zone of perched water table development for most wells. Water was injected into the PVC tube via a funnel until the tube was completely filled up. There were some cases where it was not possible to fill up the PVC tube because of very rapid drainage.

At both forested hillslopes the location of each tree, type of tree, and stem diameter at breast height (DBH) were mapped. A grid out of barrier tape had been laid out on the hillslopes prior to mapping (3 m × 7.5 m grid size). The mapped area covers 36 m × 75 m at the coniferous forest hillslope and 36 m × 60 m at the mixed forest hillslope (3 m buffer at both sides of the horizontal transects, and 7.5 m buffer up- and downslope of the upper and lower transect, respectively). In addition, hemispherical pictures of canopy coverage were taken above each well with a digital camera and a fish-eye lens.

2.3 Determination of water table response variables

Water table response data is available for the time period April 2010 until September 2011 (18 months). Different water table response variables were determined per well for three different temporal scales:

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Entire time series (18 months)
- Seasonal scale covering 3 months each (winter: Jan/Feb/Mar, spring: Apr/May/June, summer: Jul/Aug/Sep, fall: Oct/Nov/Dec)
- Event scale: five selected events covering a range of total rainfall, mean and maximum rainfall intensity, and antecedent wetness conditions (see Table 1)

For each temporal scale following response variables were calculated:

- AREA_NORM: Area below the water table time series normalized to well depth
- INDEX_ACTI: Percentage of time during which a well is activated; as activated we define a water table >2 cm
- INDEX_150: Percentage of time during which the water table depth below the soil surface is <150 cm

Index range is 0–1. For the event scale, two additional response variables were determined: SLOPE (mean slope of water table rise), and LAG (time lag from begin of rainfall until maximum water table; instead of the overall maximum we chose 95% of the maximum to give less weight to response curves that slowly reach a plateau).

Several other water table response variables had been computed (e.g. mean water table height, absolute water table rise, coefficient of variation, percentage of time during which at least 10%, 30% or 50% of the well is saturated); however, since they were either meaningless (coefficient of variation) or highly correlated with AREA_NORM, INDEX_ACTI, and INDEX_150 we omitted them from further analyses. Note that the three selected response variables are highly correlated as well (AREA_NORM versus INDEX_ACTI: $r = 0.74 - 0.86$ for all temporal scales; AREA_NORM versus INDEX_150: $r = 0.77 - 0.86$ for all temporal scales; INDEX_ACTI versus INDEX_150: $r = 0.78 - 0.9$ for all temporal scales).

Missing data handling: Seasonal scale: If there were small data gaps we used the remaining time period for calculating well response variables. For comparability, all

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



response variables are normalized to the length of the time series. Wells with larger data gaps were treated as wells with missing values during this season. Event scale: Wells with missing data were omitted. Entire time series: Response variables were calculated from available data and normalized to the length of the time series. We tested the representativity of variables calculated from time series with partially missing data by applying the observed data gaps to time series with complete data. Since the representativity was reasonable to high, each well response variable is considered representative for the entire time series despite missing values. The data analysis was conducted with IDL (Interactive Data Language, ITT VIS).

2.4 Determination of hillslope characteristics

The following hillslope characteristics per well were determined in the field or with GIS analysis: upslope contributing area (UA), slope (SLOPE), plan curvature (PLAN_CURV), profile curvature (PROF_CURV), aspect (ASP), well depth (WDEPTH), hydraulic conductivity from slug tests according to HORSLEV method (K), slope of the fast part of the recession after slug injection (SLUG_HIGH), slope of the slow part of the recession after slug injection (SLUG_LOW), number of trees in vicinity (N_TREES), stemflow index (STEMF), percent throughfall (THROUGHF), and canopy coverage (CANOPY_COV). In addition, we created two categorical variables representing land use classification (LULC: 1 = forest, 2 = grassland), and well location on hillslope (TRANSECT: 1 = lower transect well, 2 = middle transect well, 3 = upper transect well).

Topography variables (upslope contributing area, slope, plan curvature, profile curvature, and aspect) were determined with a 1 m LiDAR DEM. All calculations were carried out with SAGA GIS: preprocessing of the DEM (sink removal, smoothing with simple filter (3 m radius)), standard terrain analysis (slope, aspect, plan curvature, profile curvature), and upslope contributing area (interactive module, multiple flow direction). The exact location of wells was determined by surveying with a laser theodolite.

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil properties were either directly measured in the field (WDEPTH) or calculated from slug tests (K, SLUG_HIGH, and SLUG_LOW). Slug test data was preprocessed by correcting hydrostatic head with barometric head. Water table height prior to slug injection was determined and subtracted from measured water table height. Many wells, however, were dry prior to slug injection. The time series were then manually trimmed so that the start represents the highest water table followed by a constant drop (e.g. if there was fluctuation around a plateau the last value prior to a constant decline was defined as start value). Determining hydraulic conductivity from the preprocessed time series was challenging since common methods, e.g. Bouwer and Rice (1976), Hvorslev (1951), assume a confined or unconfined aquifer. As already stated, unsaturated conditions were found in many wells. We chose to analyze the data according to the Hvorslev method (Hvorslev, 1951). We emphasize that the calculated values should not be regarded as physically true saturated hydraulic conductivity. We solely use K values as tool for intercomparing well behaviour.

Due to the limited applicability of common methods for determining hydraulic conductivity in our case, we calculated two additional variables from the slug test curves: SLUG_HIGH and SLUG_LOW, which represent the mean slope of the fast and slow part of the recession after slug injection, respectively. The calculations were carried out with IDL. First, the time series was smoothed (boxcar average) and the slope of the recession curve was calculated. A threshold of 0.5 cm s^{-1} was used to separate the fast part of the recession from the slow part. This threshold fitted best visual observations and resulted in two parts of the time series with distinct slope. The mean slope per recession part was calculated after a buffer had been applied to cut data around the breakpoint. For wells that only showed a fast part of recession (fast decline to 0 cm, hence no slow part of recession) SLUG_LOW was set to an arbitrary value of 10 cm s^{-1} indicating a large hydraulic conductivity. For wells with extremely rapid drainage, where the lower 50 cm of the well could not be filled up and SLUG_HIGH could not be determined, SLUG_HIGH was set to an arbitrary value of 150 cm s^{-1} indicating a large hydraulic conductivity.

Hillslope characteristics

S. Bachmair and
M. Weiler

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The amount of trees in the vicinity of the wells and the stemflow index were derived from tree mapping and GIS analysis. The variable N_TREES is the number of trees located within a 3 m radius of a well. Since we did not measure stemflow we calculated a stemflow index based on amount of upslope trees, distance from well, type of tree, and DBH: First, all trees (i) within 3 m distance and within the upslope quadrant or (ii) within 1 m distance and within the upslope semi-circle were identified. Trees meeting neither one of these conditions (e.g. located downslope or >1 m to the side of a well) were not considered as stemflow contributing. Next, the stemflow index per tree was calculated by adding up normalized tree distance (normalized to maximum distance of 3 m), normalized DBH (normalized to maximum DBH), and a factor representing type of tree (rough-barked spruce/fir: 0.5, smooth-barked beech/ash tree: 1). These factors were set according to literature values of stemflow for different forest types (Levia et al., 2011). The maximum stemflow index per tree is 3; the overall stemflow index is the sum of stemflow indices for all trees contributing stemflow to one well. Self-evidently the derived stemflow index is based on subjective decisions and does not represent actual stemflow volumes. Nevertheless, we assume this index suitably characterizes the proneness of a well to receive stemflow.

The variable THROUGHF represents mean throughfall of eight events per well during the leaf season in 2010/2011. Note that at the grassland hillslope THROUGHF represents mean open area rainfall. THROUGHF per event was normalized to maximum throughfall, which was usually higher than open area rainfall. The eight events cover a range of rainfall characteristics (total rainfall, mean and maximum rainfall intensity). Standard deviation of mean throughfall is <15 % for 82 % of wells; the rest of wells have a standard deviation between 15 % and 28.5 %.

Canopy coverage was derived from digital fish-eye photographs using the Gap Light Analyzer (GLA) imaging software (Frazer et al., 1999). We determined the percentage of canopy coverage per well for three different zenith angles (4.5°, 9°, and 18°). Since the canopy coverage for different zenith angles was highly correlated, we only con-

sidered the canopy coverage for 9° zenith angle for further analysis (4.5°/9°: $r = 0.96$, 9°/18°: $r = 0.82$, 4.5°/18°: $r = 0.93$).

2.5 Statistical analysis

2.5.1 Correlation analysis

5 Since the response variables and some of the predictor variables are not normally distributed, we applied non-parametric statistical analyses. Spearman rank correlation coefficients were computed between different predictor variables. To assess the influence of different predictor variables on the water table response, rank partial correlation coefficients (Spearman) were calculated. Partial correlation is the correlation between
10 a predictor variable and a response variable when the effects of all other predictor variables involved are removed. The hypothesis of no partial correlation was tested against the alternative that the partial correlation is significantly different from zero. Rank correlation, rank partial correlation, and statistical tests were carried out with MATLAB.

2.5.2 Random forests

15 In addition to correlation analysis, a random forest (RF) approach was chosen for revealing the effect of different hillslope characteristics on the observed water table response variables. A random forest represents a machine learning algorithm, where a large number of classification or regression trees (CARTs) are grown on a bootstrapped subsample of the data. The method developed by Breiman (2001) is a non-parametric multivariate technique. Generally, classification or regression trees explain the variation
20 of a response variable by recursively splitting the data into more homogeneous groups (nodes) based on combinations of explanatory variables (Breiman et al., 1984; De'Ath, 2002; Strobl et al., 2009). The advantage of CART is that no assumptions are made about the distribution of the data and both continuous and/or categorical variables can
25 be used. In addition, CART can handle interactions and nonlinearities among vari-

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ables. A disadvantage of CART is tree instability, since small changes in input data can produce highly divergent trees (Prasad et al., 2006). Thus, an ensemble of trees, a “random forest”, was shown to yield better predictions than a single tree (Strobl et al., 2008).

5 The random forest algorithm is described in detail by Breiman (2001). For our analysis we used the “RandomForest” package by Liaw and Wiener (2002) implemented in the R environment, where also details about the program can be found. The steps during the RF analysis are: (1) n_{tree} bootstrap samples are drawn from the original dataset, each sample containing about two-thirds of the data (n_{tree} is a user-specified parameter); (2) for each bootstrap sample, an un-pruned tree is grown. In regular CARTs
10 the data is split using the best split among all predictor variables. For random forest construction a subset of predictor variables is randomly chosen (amount of randomly selected predictor variables (m_{try}) specified by user). The best split among those variables is determined. New data is predicted by aggregating the predictions of all trees
15 (i.e., majority votes for classification, average for regression) (Liaw and Wiener, 2002). (3) At each bootstrap iteration, the data not included in building the tree model (“out-of-bag” data after Breiman (2001)) is predicted using the tree grown with the bootstrap sample. The “out-of-bag” predictions are aggregated and the error rate calculated. (4) The variable importance is determined for each predictor by assessing how much
20 the prediction error increases when the “out-of-bag” data for that predictor is permuted while all others are left unchanged (Liaw and Wiener, 2002).

The outcome of the random forest regression analysis that we use for interpretation is model performance (percentage of explained variance) and the variable importance measure %IncMSE (percent increase in mean square error) (Liaw and Wiener, 2002).
25 The variable importance measure indicates how much worse the prediction would be if the data for that predictor were permuted randomly (Prasad et al., 2006). Large values thus represent an important variable. Strobl et al. (2009) state that “the absolute values of the importance scores should not be interpreted or compared over different

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



studies". Instead, variable importance should be regarded as relative ranking (Strobl et al., 2009).

We set n_{tree} to 3000, since this resulted in a rather stable prediction error (default = 500); m_{try} was left at the default value (5). According to Díaz-Uriarte and De Andres (2006), changes in these parameters have in most cases negligible effects, suggesting the default values are a good option. Due to the random nature of the RF method, the model results slightly vary from run to run (e.g. standard deviation of explained variance 0.3% – 0.9% for the response variable AREA_NORM at the seasonal scale). Some authors average variable importance measures over several runs, e.g. Grimm et al. (2008) or Loos and Elsenbeer (2011). We conducted several runs per RF and chose the variable importance for the run with the highest percentage of explained variance. We found that the top-scoring variables were unaffected; variables with similar variable importance values appeared in slightly different order per run. We thus only give weight to the highest scoring variables and interpret the rank of variables rather as a range of appearance (e.g. top-scoring, medium and lower range of variable importance).

3 Results

3.1 Input variables

The water table response variables AREA_NORM, INDEX_ACTI and INDEX_150 show high spatial variability (see Fig. 2 for AREA_NORM during the entire time and at the seasonal scale, other response variables and event scale not shown). Remarkable is within-transect variability, especially at the event scale, where adjacent wells exhibit contrasting water table dynamics (dry well/weak response vs. strong response within 3 m distance). At the same time spatial patterns of water table dynamics vary among seasons and events. Moreover, there are differences among hillslopes: the mean of AREA_NORM and the mean of INDEX_ACTI per hillslope ranks according to coniferous

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



forest larger mixed forest larger grassland. The predictor variables also show high spatial variability (see supplementary material). Correlation between predictors is mostly low (Table 2). Exceptions are a moderate to high correlation between different pairs of vegetation predictors (NTREES, TROUGHF, CANOPY_COV, STEMF, and LULC), and a moderate correlation between the categorical variable transect and some topography variables (UA, SLOPE, PROF_CURV).

3.2 Partial correlation

The partial correlation coefficients depict the relationship between a predictor variable and a response variable when the effects of all other predictor variables are removed. As presented in Table 3, about half of the predictors show no significant correlation with the response variable AREA_NORM. For the predictor variables that are correlated with AREA_NORM the strength of the relationship is generally low to moderate. Influential predictors are some topographic variables (slope, transect, aspect in some cases), soil properties (SLUG_LOW = slope of the slow part of the recession after slug injection, well depth in some cases, K for very few cases), and LULC (land use classification into forest vs. grassland). The spatially variable vegetation predictors throughfall, canopy coverage, stemflow index and amount of trees are not correlated with the water table response AREA_NORM, except for few cases with very low partial correlation (N_TREES, CANOPY_COV: $r = -0.2$ for few temporal scales). For the response variables INDEX_ACTI and INDEX_150 the same tendency is found. Regarding the response variables mean slope and lag time until water table rise (available at the event scale only), a slightly different pattern is discernible: while LULC and SLUG_LOW are insignificant, the spatially variable vegetation predictors show higher correlations for few cases.

Table 3 also clearly reveals distinct differences among temporal scales. First, the correlation between different predictors and water table dynamics at the event scale tends to be lower and is more often insignificant than at the seasonal scale and for the entire period. Second, there seems to be a pattern between strength of rela-

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cept at the event scale. INDEX_150 is almost always lower than AREA_NORM and INDEX_ACTI.

Remarkable are the strong differences among temporal scales: The explained variance of the random forest models predicting AREA_NORM and INDEX_ACTI lies above 20% considering the entire time and most seasons. At the event scale, in contrast, the explained variance is always below 20% (for most cases below 10%) except for the event on 12-May 2010. A closer look at the differences among model performance reveals that models predicting the water table response for summer seasons and summer events tend to have a lower percentage of explained variance, e.g. 2_2010 (Apr–Jun) and 3_2010 (Jul–Sep) vs. 4_2010 (Oct–Dec) and 1_2011 (Jan–Mar), and the events on 12-Jul, 12-Sep, and 17-Jun (summer) vs. 12-May and 14-Dec (spring/winter). Similar to the results of the partial correlation analysis we found a significant negative correlation between RF model performance and mean rainfall intensity for AREA_NORM and INDEX_ACTI (see Table 5). This indicates that high rainfall intensity results in low explainability of water table response. As for the partial correlation analysis, antecedent wetness is positively correlated with RF model performance at the event scale; yet, the correlation is not statistically significant.

Regarding the differences among land use type, the RF models predicting the water table response AREA_NORM for the grassland hillslope ($n = 30$) show a distinctly lower percentage of explained variance than for the forested hillslopes (see Fig. 4 bottom graph). The RF models of the forested hillslopes ($n = 60$) perform slightly lower than when considering all wells ($n = 90$). The models for the other response variables are comparable, yet INDEX_ACTI shows more variability (few cases with model performance grassland hillslope > forested hillslopes and forested hillslopes > considering all wells). Since most models at the grassland hillslope perform very poorly overall (<10% explained variance), the chosen predictors cannot adequately explain the water table response at the grassland hillslope.

HESSD

9, 6889–6934, 2012

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3.2 Variable importance

Predictor variables can be considered informative and important for a RF model if their variable importance value is above the absolute value of the lowest negative-scoring variable (Strobl et al., 2009). Thus, only variables meeting this condition are displayed in the variable importance plots. As can be seen in Fig. 5, variable importance varies among models for different temporal scales. Despite this complexity a few definite trends can be identified for tree models explaining the water table response variable AREA_NORM. The predictors SLUG_LOW (slope of the slow part of the recession after slug injection) and PROF_CURV (profile curvature) are the most often occurring top-scoring variables and consequently have the highest single explanatory power. The percent increase in mean square error for SLUG_LOW (soil property variable) and PROF_CURV (topography) exceeds other variable importance values by far for the entire period, 4_2010, 1_2011, 2_2011, and 3_2011. Predominantly topography predictors (slope, plan curvature, upslope contributing area, transect, and aspect) are found in the upper third of the variable importance ranking following SLUG_LOW and PROF_CURV. Well depth is intermediate-ranking and shows high variability; other soil properties (K, SLOPE_HIGH) show lower importance, if any. Vegetation predictors (throughfall, canopy cover, stemflow, amount of trees, and LULC) primarily occur in the lower third of the variable importance ranking. Solely canopy cover and throughfall appear on higher ranks for some RF models; interestingly, these models all represent summer periods (seasons 3_2010, 2_2011, 3_2011, events on 12-May, 12-Jul, 12-Sep, and 17-Jun). Nevertheless, one has to keep in mind that for many of these summer period models the overall explanatory power is low, especially at the event scale.

RF models explaining the water table response variable INDEX_ACTI (percentage of time during which a well is activated, i.e. displays a water table >2 cm) follow a similar trend as for AREA_NORM (Fig. 6). SLUG_LOW is the highest-scoring predictor in most cases. Profile curvature is high-ranking as well but less important than for AREA_NORM. Instead, well depth plays a more important role. The general trend

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of topography (PROF_CURV, SLOPE, UA, PLAN_CURV, TRANS) and soil properties (SLUG_LOW, WDEPTH) occupying higher ranks, while most vegetation parameters score lower (LULC, STEMF, N_TREES) stays the same. However, the vegetation predictors throughfall and stemflow stick out: THROUGHF ranks second and third for 3_2011 (26% explained variance) and 2_2011 (32% explained variance), respectively. CANOPY_COV continuously occupies middle ranks. RF models for INDEX_150 (percentage of time during which the water table depth below the soil surface is <150 cm) are comparable to RF models of AREA_NORM (not shown). SLUG_LOW and PROF_CURV are most important followed by the topography predictors slope, transect, plan curvature, and upslope contributing area. WDEPTH plays a minor role, if any. While LULC, STEMF, and N_TREES show a negligible importance, THROUGHF and CANOPY_COV continuously appear on intermediate ranks.

In terms of the effect of different temporal scales, variable importance is more variable and trends are less clear at the event scale. In addition, the vegetation predictors throughfall and canopy cover are more important during summer periods. Regarding the effect of land use type, RF models explaining the water table response for the forested hillslopes differ little from RF models including all wells. SLUG_LOW, PROF_CURV and WDEPTH score highest when regarding all three response variables (AREA_NORM, INDEX_ACTI, and INDEX_150). Noticeable is that among these highest-ranking predictors well depth plays a more important role than for models considering all wells. Vegetation predictors follow the same pattern as for RF models of all wells. In contrast, the importance of predictor variables for RF models predicting the water table response at the grassland hillslope is distinctly different: profile curvature, throughfall, and transect are top-scoring with interchanging order while soil properties (SLUG_LOW, WDEPTH) do not appear at all. Interesting is that THROUGHF, which represents open area rainfall at the grassland hillslope, only marginally differs among transects (lower transect: 90.1%, middle transect: 89.8%, upper transect 86.2%; values per transect represent the mean of two rainfall totalizers over 8 events). Open area rainfall and also profile curvature (lower transect slightly concave, middle and up-

HESSD

9, 6889–6934, 2012

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



per transect more planar/convex) thus represent similar information as the predictor transect. This fits well with the observation that mostly the lower transect wells show a water table response at the grassland hillslope. Nonetheless, variable importance needs to be interpreted against the background that RF models of the grassland hillslope generally have a very poor explanatory power.

4 Discussion

4.1 Can the spatial variability of SSF dynamics be explained by measurable hillslope characteristics?

The partial correlation analysis and the random forest approach revealed similar results:

1. The observed spatio-temporal variability of water table response results from a complex set of interactions among hillslope characteristics. Soil properties and topography show the highest single explanatory power.
2. Vegetation predictors surprisingly played a minor role, if any. Throughfall and canopy cover were the most important variables among vegetation predictors.
3. The mapped hillslope characteristics explain the observed spatial variability of water table dynamics to some extent. However, since they do not sufficiently describe the response there must be additional important drivers not captured by our measurements.

Despite some differences in outcome of both analyses, the observed similar results of both methods independently validate these conclusions. The high explanatory power of topography is interesting given the predominantly planar side slopes. Clearly, hillslope topography has been identified as a dominant control (e.g. Fujimoto et al., 2008; Hopp and McDonnell, 2009; Berne et al., 2005; Bogaart and Troch, 2006); however,

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(micro-) topography differences are rather small at the studied hillslopes relative to differences within a catchment. The RF models identified profile curvature as a highly important variable. This predictor did not show any significant partial correlation with water table response. Instead, slope explained the most among topography variables.

We assured the meaningfulness of different predictors during ensemble tree construction by single regression trees and partial dependence plots of ensemble tree models, which illustrate the relationship between the response variable and a given predictor after averaging out the effects of the other predictor variables (Liaw and Wiener, 2002). Important topography predictors all showed meaningful relationships e.g. concave profile curvature or high upslope contributing area resulting in strong water table response. Soil properties (hydraulic conductivity and well depth) as influential drivers are not surprising. Interesting, however, is that SLUG_LOW was identified as most important variable among variables calculated from slug tests, and also showed highest partial correlation among them (K after Hvorslev method, SLUG_LOW, SLUG_HIGH). Consequently, SLUG_LOW seems to best represent actual hydraulic conductivity. It is possible that SLUG_HIGH (fast part of recession, duration sometimes only for a few time steps) does not stand for true physical soil properties but rather for initial effects after slug injection.

For us surprising is the subordinate effect of vegetation. Our hypothesis that vegetation exerts a major control on spatially variable SSF dynamics has to be rejected. This is remarkable given the many studies reporting on the effect of precipitation redistribution (stemflow, throughfall) and rooting patterns on hydrologic fluxes (e.g. Nordmann et al., 2009; Nikodem et al., 2010; Chang and Matzner, 2000; Jost et al., 2012; Jost et al., 2004; Gerrits, 2010; Sansoulet et al., 2008; Liang et al., 2007; Liang et al., 2011; Keim et al., 2006). In our study no significant partial correlation between stemflow index and different water table response variables could be identified and the variable importance during tree model construction was marginal. The predictor number of trees in vicinity of a well showed slightly higher variable importance and there were few cases with low but significant partial correlation. This variable may act as proxy for some vegetation

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



related parameter, such as root density or root water uptake, but may also simply mimic canopy coverage and throughfall. THROUGHF and CANOPY_COV showed the highest influence among vegetation predictors during tree model construction, yet still played a secondary control overall. Noteworthy is the relative stronger influence during summer seasons and summer events. Our experimental findings cannot confirm the clear effect of interception on subsurface saturation patterns as shown by two virtual experiments (Keim et al., 2006; Gerrits, 2010). Another virtual experiment, however, also found only a minor effect of fine-scale throughfall patterns on the hillslope hydrological response (Hopp and McDonnell, 2011).

The question is why vegetation predictors at our hillslopes exert a generally weak control on spatial variability of shallow water table dynamics, even though other studies proved the importance of vegetation on hydrologic fluxes. One reason may be comparably thick soils and hence a deep unsaturated zone, which mask or blur the effect of spatially variable input due to throughfall and stemflow. It would be interesting to assess whether the same analysis for hillslopes with shallow soil provides a different picture regarding the effect of vegetation. Other controls, e.g. highly transmissive soil, as observed for some wells, may also override effects of precipitation redistribution. Furthermore it is possible that our calculated stemflow index is not representative for actual stemflow volumes, but quantitative information was not available.

Another important outcome of our analyses is that the predictors representing many facets of the hillslope configuration do not sufficiently explain the spatial variability of water table response. The explained variance of different ensemble tree models rarely exceeded 30 percent and was often lower. Hence, there must be very important additional drivers not captured by our measurements: Bedrock topography is not too well represented by our predictors except for well depth. However, bedrock upslope contributing area and micro-relief was found to strongly govern the hillslope hydrologic response (e.g. Tromp-van Meerveld and McDonnell, 2006a; Graham et al., 2010; Freer et al., 2002). Nevertheless, high-resolution information about subsurface topography is rare and highly elaborate to determine. The periglacial deposits at our study site

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and therefore no clear soil-bedrock-interface add even more complexity. Experimental studies at sites with similar subsurface structure showed subsurface flow occurrence in different layers of periglacial deposits, and different behavior of impeding layers depending on antecedent wetness (Chiffard et al., 2008; Nordmann et al., 2009). Additional drivers may be preferential pathways (macropores, roots, soil pipes), as numerous observed (e.g. Uchida et al., 2002; Weiler and Flüher, 2004; Bachmair et al., 2009; Anderson et al., 2009; Noguchi et al., 1999; Kienzler and Naef, 2008). Deriving quantitative data representing this variable at high spatial resolution for a large hillslope seems impossible with current methods (excavations, electrical resistivity tomography). Further controls could also be spatially variable surface conditions due to litter, soil compaction, and hydrophobicity

4.2 Are there differences in explainability of water table dynamics among temporal scales and seasons/events?

Both analyses exposed that the water table response at the event scale is less explainable: partial correlation between influential predictors and water table response and explained variance of ensemble tree models was mostly lower than at the seasonal scale and for the entire time. Furthermore, trends of predictor variable importance were more variable at the event scale. Short-term variability of water table dynamics is thus less explainable than longer-term variability. A closer look at the differences among seasons and events revealed that there is a tendency of lower explainability for summer seasons and summer events (by degree of explainability we mean strength of partial correlation between water table response and influential predictors, and degree of ensemble tree model performance). Interestingly, we found a negative correlation between explainability of water table response and rainfall intensity. We also found a moderate relationship between explainability of water table response and AWI; nevertheless, due to the small sample size (AWI only available at event scale) this should be interpreted cautiously. These findings indicate that the mapped predictors explain the observed water table response for time periods with high rainfall intensity (and low AWI)

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to a lesser degree. We reason that during such periods additional drivers of SSF dynamics play a more important role, e.g. stronger hydrophobicity and thus bypass-flow, exploitation of preferential pathways initiated by high rainfall intensity.

The assumption about rainfall intensity and antecedent wetness inducing different infiltration and saturation patterns coincides with findings by Kienzler and Naef (2008). Sprinkling experiments at several hillslopes revealed differences in SSF response, which they attribute to “indirect feeding” of preferential pathways via saturated areas, or “direct feeding” via rainfall (Kienzler and Naef, 2008). We suggest that direct feeding of preferential pathways, evoked by high rainfall intensity, plays a bigger role during summer seasons and summer events, thereby reducing the correlation between water table response and measured hillslope characteristics. It would be interesting to assess the relationship between saturation patterns and subsurface flow volume. If preferential pathways were an important control in summer, saturation patterns and subsurface flow volume would be lowly correlated, since preferential flow takes places under unsaturated conditions. A recent review highlights that preferential flow does not require matrix saturation or saturation of the conduits (Nimmo, 2012). Following the concept by Kienzler and Naef (2008), we would expect indirect feeding of preferential pathways via saturated areas during seasons with low rainfall intensity and high antecedent wetness, and therefore a higher correlation between saturation patterns and subsurface flow volume, and saturation patterns and measured hillslope characteristics.

The observation that throughfall and canopy cover had a greater influence on random forest models during summer periods also highlights the seasonality effect. More pronounced throughfall patterns in summer would explain the higher importance. There is no consensus whether increasing rainfall intensity de- or increases spatial variability of throughfall, yet higher total rainfall reduces spatial variability (Levia and Frost, 2006). The differences in importance among seasons likely go back to foliation vs. defoliation of deciduous trees. Staelens (2006), for instance, found a significantly higher spatial variability of throughfall during the leafed season than during the leafless.

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.3 Are there differences in explainability of water table dynamics regarding land use?

When we split the water table response into forest hillslope response vs. grassland, distinct differences emerged. There were minor differences in explainability of water table dynamics between the base case (all wells) and forest only, whereas the grassland hillslope response was basically not explainable with the same predictors (hardly any significant partial correlation and RF model performance very poor). There are two possible explanations why shallow water table dynamics at the grassland hillslope are not attributable to hillslope properties. Either the water table response at the grassland hillslope is inherently different due to other runoff generation mechanisms for this type of land use, e.g. higher importance of preferential flow, different infiltration processes etc. Or the grassland hillslope shows distinct differences in static drivers, i.e. drivers not influenced by land use, such as (bedrock-) topography. In this case differences are coincidental and not physically attributable to vegetation cover effects.

To validate these possible explanations we carried out non-parametric statistical tests to assess whether topography and soil properties of the three hillslopes significantly differ among hillslopes (Kruskal-Wallis test comparing all hillslopes; Wilcoxon rank-sum test for pairwise tests). We found that upslope contributing area and well depth are significantly different among all hillslopes, and between each pair of hillslopes ($p < 0.05$). Plan curvature significantly differs between grassland vs. mixed forest and grassland vs. coniferous forest ($p < 0.05$), yet not between both forest hillslopes. Consequently, smaller UA, slightly convex plan curvature, and deeper soil may induce a different SSF response at the grassland hillslope. One could hypothesize there is a threshold concerning input volume until the SSF response follows the pattern of the forested hillslopes (higher storage capacity of the grassland hillslope and lower UA). On the other hand, one could argue the grassland hillslope is more homogeneous due to the homogeneous vegetation cover, whereas trees enforce small-scale spatial variability of rainfall input (and other micro-meteorological parameters), soil properties

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and soil moisture, and thus hydrologic fluxes. At the grassland hillslope there is a distinct pattern of stronger and more continuous response at the lower transect, while the upslope transects show little to no SSF response. In the forest there is more spatial variability within and among transects, and upper transect wells exhibit strong water table fluctuations (Bachmair et al., 2012). It is hence possible that there are inherent differences in spatial variability of hillslope properties due to vegetation cover, which lead to a higher explanatory power of water table variability for the forested hillslopes vs. the grassland hillslope.

5 Conclusions

Partial correlation analysis and a random forest approach elucidated the relationship between variability of shallow water table response and different hillslope characteristics. Following key points were uncovered:

- Complex interplay of predictors: The observed variability of water table response results from complex interactions among hillslope characteristics. Soil properties and topography, despite predominantly planar hillslopes, showed the highest single explanatory power.
- Minor role of vegetation: Vegetation predictors played a minor role, if any. Solely throughfall and canopy cover exerted a slightly stronger control, especially in summer.
- Low overall explanatory power: The examined hillslope characteristics do not sufficiently explain the water table response. Hence, there must be additional very important drivers not represented by our measurements of the hillslope configuration (e.g. bedrock topography, preferential pathways, hydrophobicity).
- Differences among temporal scales: Short-term variability was less explainable than longer-term variability. Additionally, the degree of explainability among sea-

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sons and events was correlated with rainfall intensity: the mapped predictors explain the water table response for time periods with high rainfall intensity to a lesser degree.

- Grassland vs. forest: Explainability of SSF variability at only the forested hillslopes was similar to the base case (all wells). In contrast, the SSF response at only the grassland hillslope was hardly explainable with the same predictors.

The fact that detailed information on the hillslope configuration did not help to sufficiently explain the SSF response raises the question about the validity of physically-based process or numerical models designed to predict subsurface flow at the observed scale. A better characterization of measurable model parameters at this spatial scale will be hard to derive. One possible reason for the mediocre overall explainability is that additional drivers such as preferential pathways, especially during high-intensity rainfall could not be derived. Further research should therefore focus on how to represent spatially and temporarily variable preferential pathways in models and how to parameterize them. Ecohydrological feedbacks also need to be further assessed. Are the minor effects of vegetation different at other hillslopes? Furthermore we need to assess the relationship between saturation patterns and subsurface flow volume, and whether this relationship varies temporally due to connectivity of saturated zones or onset of preferential pathways.

Supplementary material related to this article is available online at:
<http://www.hydrol-earth-syst-sci-discuss.net/9/6889/2012/hessd-9-6889-2012-supplement.pdf>

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. We thank all people helping with fieldwork, especially Emil Blattmann, Benjamin Grahler, Jakob Sohrt, Christian von Sperber, Matthias Ritter, and Ramona Skozilas. We thank the community of Au and several private land owners for the provision of field sites. The scholarship by the German National Academic Foundation for the first author is gratefully acknowledged.

References

- Anderson, A. E., Weiler, M., Alila, Y., and Hudson, R. O.: Dye staining and excavation of a lateral preferential flow network, *Hydrol. Earth Syst. Sci.*, 13, 935–944, doi:10.5194/hess-13-935-2009, 2009.
- Arno, K., Lindemann, J., Schellenberger, A., Beierkuhnlein, C., Kaupenjohann, M., and Peiffer, S.: Slope deposits and water paths in a spring catchment, Frankenwald, Bavaria, Germany, *Nutr. Cycl. Agroecosys.*, 50, 119–126, 1998.
- Aryal, S., O’Loughlin, E., and Mein, R.: A similarity approach to predict landscape saturation in catchments, *Water Resour. Res.*, 38, 1208, doi:10.1029/2001WR000864, 2002.
- Bachmair, S., Weiler, M., and Nützmann, G.: Controls of land use and soil structure on water movement: Lessons for pollutant transfer through the unsaturated zone, *J. Hydrol.*, 369, 241–252, 2009.
- Bachmair, S., Weiler, M., and Troch, P. A.: Intercomparing hillslope hydrological dynamics: spatio-temporal variability and vegetation cover effects, *Water Resour. Res.*, 48, W05537, doi:10.1029/2011WR011196, 2012.
- Bogaart, P. W. and Troch, P. A.: Curvature distribution within hillslopes and catchments and its effect on the hydrological response, *Hydrol. Earth Syst. Sci.*, 10, 925–936, doi:10.5194/hess-10-925-2006, 2006.
- Bouwer, H. and Rice, R.: A slug test for determining hydraulic conductivity of unconfined aquifers with completely or partially penetrating wells, *Water Resour. Res.*, 12, 423–428, 1976.
- Breiman, L.: Random forests, *Mach. Learn.*, 45, 5–32, 2001.
- Chang, S. and Matzner, E.: The effect of beech stemflow on spatial patterns of soil solution chemistry and seepage fluxes in a mixed beech/oak stand, *Hydrol. Process.*, 14, 135–144, 2000.

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Chiffard, P., Didszun, J., and Zepp, H.: Skalenübergreifende Prozess-Studien zur Abflussbildung in Gebieten mit periglazialen Deckschichten (Sauerland, Deutschland), *Grundwasser*, 13, 27–41, 2008.
- De'Ath, G.: Multivariate regression trees: a new technique for modeling species-environment relationships, *Ecology*, 83, 1105–1117, 2002.
- Díaz-Uriarte, R. and De Andres, S. A.: Gene selection and classification of microarray data using random forest, *BMC Bioinformatics*, 7, doi:10.1186/1471-2105-7-3, 2006.
- Frazer, G. W., Canham, C. D., and Lertzman, K. P.: Gap Light Analyzer (GLA), Version 2.0: Imaging software to extract canopy structure and gap light transmission indices from true-colour fisheye photographs, users manual and program documentation, Simon Fraser University, Burnaby, British Columbia, and the Institute of Ecosystem Studies, Millbrook, New York, 1999.
- Freer, J., McDonnell, J., Beven, K., Peters, N., Burns, D., Hooper, R., Aulenbach, B., and Kendall, C.: The role of bedrock topography on subsurface storm flow, *Water Resour. Res.*, 38, 1269, doi:10.1029/2001WR000872, 2002.
- Fujimoto, M., Ohte, N., and Tani, M.: Effects of hillslope topography on hydrological responses in a weathered granite mountain, Japan: comparison of the runoff response between the valley-head and the side slope, *Hydrol. Process.*, 22, 2581–2594, 2008.
- Geologisches Landesamt Baden-Württemberg (Ed.): Geologische Karte von Baden-Württemberg 1:50 000, 3rd Edn., Freiburg i. Br., 1996.
- Gerrits, A. M. J.: The role of interception in the hydrological cycle, Delft University of Technology, 2010.
- Gerrits, A. M. J., Hopp, L., McDonnell, J. J., Savenije, H. H. G., and Pfister, L.: A virtual experiment on the effect of canopy and forest floor interception on subsurface flow behaviour, *Geophys. Res. Abstr.*, 11, EGU2009-7546-1, 2009.
- Graham, C. B., Woods, R. A., and McDonnell, J. J.: Hillslope threshold response to rainfall: (1) A field based forensic approach, *J. Hydrol.*, 393, 65–76, 2010.
- Grimm, R., Behrens, T., Marker, M., and Elsenbeer, H.: Soil organic carbon concentrations and stocks on Barro Colorado Island – Digital soil mapping using Random Forests analysis, *Geoderma*, 146, 102–113, 2008.
- Guswa, A. J. and Spence, C. M.: Effect of throughfall variability on recharge: application to hemlock and deciduous forests in western Massachusetts, *Ecohydrology*, doi:10.1002/eco.281, 2011.

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hillslope characteristics

S. Bachmair and
M. Weiler

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Herwitz, S. R.: Infiltration-excess caused by Stemflow in a cyclone-prone tropical rainforest, *Earth Surf. Proc. Land.*, 11, 401–412, 1986.
- Holden, J. and Burt, T.: Piping and pipeflow in a deep peat catchment, *Catena*, 48, 163–199, 2002.
- 5 Hopp, L. and McDonnell, J.: Connectivity at the hillslope scale: Identifying interactions between storm size, bedrock permeability, slope angle and soil depth, *J. Hydrol.*, 376, 378–391, 2009.
- Hopp, L. and McDonnell, J. J.: Examining the role of throughfall patterns on subsurface stormflow generation, *J. Hydrol.*, 409, 460–471, doi:10.1016/j.jhydrol.2011.08.044, 2011.
- Hvorslev, M. J.: Time lag and soil permeability in ground-water observations, DTIC Document, 1951.
- 10 Jones, J. A. A.: The role of natural pipeflow in hillslope drainage and erosion: Extrapolating from the Maesnant data, *Phys. Chem. Earth*, 22, 303–308, 1997.
- Jost, G., Schume, H., and Hager, H.: Factors controlling soil water-recharge in a mixed European beech (*Fagus sylvatica* L.)– Norway spruce [*Picea abies* (L.) Karst.] stand, *Eur. J. For. Res.*, 123, 93–104, 2004.
- 15 Jost, G., Schume, H., Hager, H., Markart, G., and Kohl, B.: A hillslope scale comparison of tree species influence on soil moisture dynamics and runoff processes during intense rainfall, *J. Hydrol.*, 420–421, 112–124, doi:10.1016/j.jhydrol.2011.11.057, 2012.
- Keim, R., Skaugset, A., and Weiler, M.: Temporal persistence of spatial patterns in throughfall, *J. Hydrol.*, 314, 263–274, 2005.
- 20 Keim, R., Tromp-van Meerveld, H., and McDonnell, J.: A virtual experiment on the effects of evaporation and intensity smoothing by canopy interception on subsurface stormflow generation, *J. Hydrol.*, 327, 352–364, 2006.
- Kienzler, P. M. and Naef, F.: Subsurface storm flow formation at different hillslopes and implications for the ‘old water paradox’, *Hydrol. Process.*, 22, 104–116, 2008.
- 25 Kosugi, K., Katsura, S., Katsuyama, M., and Mizuyama, T.: Water flow processes in weathered granitic bedrock and their effects on runoff generation in a small headwater catchment, *Water Resour. Res.*, 42, W02414, doi:10.1029/2005WR004275, 2006.
- Levia, D. and Frost, E.: Variability of throughfall volume and solute inputs in wooded ecosystems, *Prog. Phys. Geog.*, 30, 605–632, 2006.
- 30 Levia, D. F., Keim, R. F., Carlyle-Moses, D. E., and Frost, E. E.: Throughfall and stemflow in wooded ecosystems, in: *Forest Hydrology and Biogeochemistry. Synthesis of Past Research*

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- and Future Directions, edited by: Levia, D. F., Carlyle-Moses, D. E., and Tanaka, T., Springer Science+Business Media B.V., 425–443, 2011.
- Liang, W.-L., Kosugi, K., and Mizuyama, T.: Heterogeneous soil water dynamics around a tree growing on a steep hillslope, *Vadose Zone J.*, 6, 879–889, 2007.
- 5 Liang, W.-L., Kosugi, K. I., and Mizuyama, T.: Soil water dynamics around a tree on a hill-slope with or without rainwater supplied by stemflow, *Water Resour. Res.*, 47, W02541, doi:10.1029/2010wr009856, 2011.
- Liaw, A. and Wiener, M.: Classification and Regression by randomForest, *R news*, 2, 18–22, 2002.
- 10 Loos, M. and Elsenbeer, H.: Topographic controls on overland flow generation in a forest – An ensemble tree approach, *J. Hydrol.*, 409, 94–103, doi:10.1016/j.jhydrol.2011.08.002, 2011.
- McDonnell, J. J., Sivapalan, M., Vache, K., Dunn, S., Grant, G., Haggerty, R., Hinz, C., Hooper, R., Kirchner, J., Roderick, M. L., Selker, J., and Weiler, M.: Moving beyond heterogeneity and process complexity: A new vision for watershed hydrology, *Water Resour. Res.*, 43, doi:10.1029/2006wr005467, 2007.
- 15 Morhard, A.: Kurzbeschreibung des Modells GWN-BW. Erweiterungen in Version 2.0, GIT HydroS Consult GmbH, Freiburg, unpublished work.
- Nieber, J. L., Steenhuis, T. S., Walter, T., and Bakker, M.: Enhancement of seepage and lateral preferential flow by biopores on hillslopes, *Biologia*, 61, 225–228, 2006.
- 20 Nikodem, A., Kodesova, R., Drabek, O., Bubenickova, L., Boruvka, L., Pavlu, L., and Tejnecy, V.: A Numerical Study of the Impact of Precipitation Redistribution in a Beech Forest Canopy on Water and Aluminum Transport in a Podzol, *Vadose Zone J.*, 9, 238–251, doi:10.2136/vzj2009.0083, 2010.
- Nimmo, J. R.: Preferential flow occurs in unsaturated conditions, *Hydrol. Process.*, 26, 786–789, doi:10.1002/hyp.8380, 2012.
- 25 Noguchi, S., Tsuboyama, Y., Sidle, R. C., and Hosoda, I.: Morphological Characteristics of Macropores and the Distribution of Preferential Flow Pathways in a Forested Slope Segment, *Soil Sci. Soc. Am. J.*, 63, 1413–1423, 1999.
- Nordmann, B., Göttlein, A., and Binder, F.: Einfluss unterschiedlicher Waldbestockung auf die Abflussbildung – ein Beispiel aus einem Wassereinzugsgebiet im Frankenwald, *Hydrol. Wasserbewirts.*, 53, 80–95, 2009.
- 30 Prasad, A. M., Iverson, L. R., and Liaw, A.: Newer classification and regression tree techniques: bagging and random forests for ecological prediction, *Ecosystems*, 9, 181–199, 2006.

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sansoulet, J., Cabidoche, Y. M., Cattan, P., Ruy, S., and Simunek, J.: Spatially distributed water fluxes in an Andisol under banana plants: Experiments and three-dimensional modeling, *Vadose Zone J.*, 7, 819–829, doi:10.2136/vzj2007.0073, 2008.

5 Staelens, J., De Schrijver, A., Verheyen, K., and Verhoest, N. E. C.: Spatial variability and temporal stability of throughfall water under a dominant beech (*Fagus sylvatica* L.) tree in relationship to canopy cover, *J. Hydrol.*, 330, 651–662, 2006.

Staelens, J., De Schrijver, A., Verheyen, K., and Verhoest, N.: Rainfall partitioning into throughfall, stemflow, and interception within a single beech (*Fagus sylvatica* L.) canopy: influence of foliation, rain event characteristics, and meteorology, *Hydrol. Process.*, 22, 33–45, 2008.

10 Strobl, C., Boulesteix, A. L., Kneib, T., Augustin, T., and Zeileis, A.: Conditional variable importance for random forests, *BMC Bioinformatics*, 9, 307, doi:10.1186/1471-2105-9-307, 2008.

Strobl, C., Malley, J., and Tutz, G.: An introduction to recursive partitioning: rationale, application, and characteristics of classification and regression trees, bagging, and random forests, *Psychol. Methods*, 14, 323–348, doi:10.1037/a0016973, 2009.

15 Troch, P., Paniconi, C., and Van Loon, E.: Hillslope-storage Boussinesq model for subsurface flow and variable source areas along complex hillslopes: 1. Formulation and characteristic response, *Water Resour. Res.*, 39, 1316, doi:10.1029/2002WR001728, 2003.

Tromp-van Meerveld, H. J. and McDonnell, J. J.: Threshold relations in subsurface stormflow: 2. The fill and spill hypothesis, *Water Resour. Res.*, 42, W02411, doi:10.1029/2004WR003800, 2006a.

20 Tromp-van Meerveld, H. J. and McDonnell, J. J.: On the interrelations between topography, soil depth, soil moisture, transpiration rates and species distribution at the hillslope scale, *Adv. Water Resour.*, 29, 293–310, 2006b.

Tromp-van Meerveld, H. J. and Weiler, M.: Hillslope dynamics modeled with increasing complexity, *J. Hydrol.*, 361, 24–40, 2008.

25 Tromp-van Meerveld, H. J., Peters, N. E., and McDonnell, J. J.: Effect of bedrock permeability on subsurface stormflow and the water balance of a trenched hillslope at the Panola Mountain Research Watershed, Georgia, USA, *Hydrol. Process.*, 21, 750–769, 2007.

30 Tsuboyama, Y., Sidle, R. C., Noguchi, S., and Hosoda, I.: Flow and solute transport through the soil matrix and macropores of a hillslope segment, *Water Resour. Res.*, 30, 879–890, doi:10.1029/93wr03245, 1994.

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Uchida, T., Kosugi, K., and Mizuyama, T.: Effects of pipeflow on hydrological process and its relation to landslide: a review of pipeflow studies in forested headwater catchments, *Hydrol. Process.*, 15, 2151–2174, 2001.

Uchida, T., Kosugi, K. I., and Mizuyama, T.: Effects of pipe flow and bedrock groundwater on runoff generation in a steep headwater catchment in Ashiu, central Japan, *Water Resour. Res.*, 38, 1119, doi:10.1029/2001WR000261, 2002.

Uchida, T., Tromp-van Meerveld, H. J., and McDonnell, J. J.: The role of lateral pipe flow in hillslope runoff response: an intercomparison of non-linear hillslope response, *J. Hydrol.*, 311, 117–133, 2005.

Völkel, J., Leopold, M., and Roberts, M.: The radar signatures and age of periglacial slope deposits, Central Highlands of Germany, *Permafrost Periglac.*, 12, 379–387, 2001.

WaBoA: Wasser- und Bodenatlas Baden-Wuerttemberg, Umweltministerium Baden-Wuerttemberg, Landesanstalt fuer Umwelt, Messungen und Naturschutz Baden-Wuerttemberg, Stuttgart, 2007.

Wagener, T., Sivapalan, M., Troch, P., and Woods, R.: Catchment classification and hydrologic similarity, *Geography Compass*, 1, 901–931, 2007.

Weiler, M. and Flüehler, H.: Inferring flow types from dye patterns in macroporous soils, *Geoderma*, 120, 137–153, 2004.

Weiler, M. and McDonnell, J.: Virtual experiments: a new approach for improving process conceptualization in hillslope hydrology, *J. Hydrol.*, 285, 3–18, 2004.

Weiler, M. and McDonnell, J. J.: Conceptualizing lateral preferential flow and flow networks and simulating the effects on gauged and ungauged hillslopes, *Water Resour. Res.*, 43, W03403, doi:10.1029/2006WR004867, 2007.

Weiler, M. and Naef, F.: An experimental tracer study of the role of macropores in infiltration in grassland soils, *Hydrol. Process.*, 17, 477–493, 2003.

Zimmermann, A., Zimmermann, B., and Elsenbeer, H.: Rainfall redistribution in a tropical forest: Spatial and temporal patterns, *Water Resour. Res.*, 45, W11413, doi:10.1029/2008wr007470, 2009.

Zimmermann, B., Elsenbeer, H., and De Moraes, J.: The influence of land-use changes on soil hydraulic properties: Implications for runoff generation, *Forest Ecol. Manag.*, 222, 29–38, 2006.

Hillslope characteristics

S. Bachmair and
M. Weiler

Table 1. Rainfall characteristics and antecedent wetness conditions (discharge before event, weighted rainfall last 14 days) per event.

Start of Event	Duration (h)	Total rainfall (mm)	Intensity		Antecedent wetness	
			Mean (mm/h)	Max. (mm/h)	Q before event (l/s)	P last 14 days weighted (mm)
12-May 2010	23.0	25.0	2.0	4.8	3.3	37.7
12-Jul 2010	1.3	19.2	16.5	52.8	no data	1.9
12-Sep 2010	3.3	17.6	6.6	19.2	0.4	16.9
14-Nov 2010	35.7	28.2	1.2	4.2	1.1	28.0 ^a
17-Jun 2011	5.3	25.2	10.8	82.8	0.6	33.3

^a Missing data supplemented with data from nearby WBI weather station (state-run viticulture institute Freiburg).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hillslope characteristics

S. Bachmair and
M. Weiler

Table 2. Rank correlation coefficients (Spearman) between predictor variables.

	UA	SLOPE	PLAN_CURV	PROF_CURV	ASP	K	SLUG_HIGH	SLUG_LOW	WDEPTH	N.TREES	THROUGHF	CANOPY_COV	STEMF	LULC	TRANS
UA	1.00	0.18	-0.15	-0.24	0.21	-0.10	-0.05	-0.27	-0.18	0.38	-0.25	0.37	0.15	-0.46	0.53
SLOPE		1.00		-0.18	0.59	-0.16	-0.28	-0.03	-0.10	0.13	-0.02	0.06	0.01	-0.12	0.47
PLAN_CURV			1.00	0.10	-0.09	-0.09	-0.08	-0.12	0.05	0.90	0.04	-0.11	0.20	0.26	0.00
PROF_CURV				1.00	-0.38	0.08	-0.10	0.04	-0.17	0.14	-0.31	0.14	0.16	-0.08	-0.58
ASP					1.00	-0.19	-0.17	-0.11	-0.01	0.06	-0.01	0.11	0.00	-0.16	0.35
K						1.00	0.24	0.29	-0.08	0.04	-0.12	0.17	0.09	-0.11	-0.10
SLUG_HIGH							1.00	0.17	-0.04	0.09	-0.16	0.18	0.03	-0.14	-0.06
SLUG_LOW								1.00	0.18	-0.30	0.22	-0.22	-0.13	0.31	-0.11
WDEPTH									1.00	-0.41	0.54	-0.60	-0.35	0.51	0.10
N.TREES										1.00	-0.62	0.57	0.64	-0.59	0.09
THROUGHF											1.00	-0.86	-0.47	0.75	0.24
CANOPY_COV												1.00	0.44	-0.83	-0.06
STEMF													1.00	-0.34	-0.08
LULC														1.00	0.00
TRANS															1.00

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hillslope characteristics

S. Bachmair and
M. Weiler

Table 3. Rank partial correlation coefficients (Spearman) between response parameter AREA_NORM and different predictors.

Predictors	entire time		seasonal scale					event scale				
	2.2010	3.2010	4.2010	1.2011	2.2011	3.2011	12-May 2010	12-Jul 2010	12-Sep 2010	14-Nov 2010	17-Jun 2011	
UA	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
SLOPE	-0.36**	-0.44***	-0.24*	-0.39***	-0.47***	-0.35**	-0.40**	-0.50***	n.s.	-0.29*	-0.36**	n.s.
PLAN_CURV	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
PROF_CURV	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
ASP	n.s.	0.26*	n.s.	n.s.	0.27*	n.s.	0.25.	0.30**	n.s.	n.s.	n.s.	n.s.
K	n.s.	0.30**	n.s.	n.s.	n.s.	n.s.	n.s.	0.21.	n.s.	n.s.	n.s.	n.s.
SLUG_HIGH	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
SLUG_LOW	-0.37**	-0.36**	-0.25*	-0.32**	-0.39***	-0.31**	n.s.	-0.34**	-0.22.	-0.25*	-0.25*	n.s.
WDEPTH	0.22.	0.27*	n.s.	n.s.	0.31**	0.34**	n.s.	0.28*	n.s.	n.s.	n.s.	n.s.
N_TREES	n.s.	n.s.	n.s.	-0.21.	-0.20.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
THROUGHF	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
CANOPY_COV	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	-0.21.	n.s.	n.s.	n.s.	n.s.
STEMF	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
LULC	-0.42***	-0.19.	-0.21.	-0.37***	-0.38***	-0.43***	-0.43***	-0.29*	n.s.	-0.21.	n.s.	-0.29*
TRANS	0.22.	0.43***	n.s.	0.27*	0.36**	0.20.	n.s.	0.48***	n.s.	0.23*	0.24*	n.s.

n.s. (not significant) = $p > 0.1$, . = $p < 0.1$, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hillslope characteristics

S. Bachmair and M. Weiler

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Table 4. Correlation coefficients highlighting the effect of rainfall characteristics and antecedent wetness on the strength of partial correlation coefficients. Displayed are linear correlation coefficients between partial *r* of the response variable AREA_NORM and different predictors and different rainfall characteristics and antecedent wetness. For rank partial correlation coefficients see Table 3, for rainfall characteristics and antecedent wetness per temporal scale see Fig. 3 and Table 1. Only predictors showing a significant partial *r* for most temporal scales were chosen for correlation (SLOPE, SLUG_LOW, WDEPTH, LULC, TRANS). Correlation with total rainfall *n* = 12 (all temporal scales), correlation with rainfall intensity *n* = 11 (seasonal and event scale), correlation with AWI *n* = 5 (event scale only).

Partial <i>r</i> with different predictor variables	Total rainfall (mm)	Rainfall intensity (mm/h)		AWI ^a (mm)
		Mean	Max.	
SLOPE	0.14 n.s.	-0.81**	-0.73*	0.49 n.s.
SLUG_LOW	0.29 n.s.	-0.46 n.s.	-0.76**	-0.04 n.s.
WDEPTH	0.16 n.s.	-0.49 n.s.	-0.16 n.s.	0.72 n.s.
LULC	0.47 n.s.	-0.57.	0.04 n.s.	0.90*
TRANS	-0.03 n.s.	-0.61*	-0.80**	0.54 n.s.

^a Antecedent wetness index: weighted rainfall of the last 14 days
n.s. (not significant) = *p* > 0.1, . = *p* < 0.1, * = *p* < 0.05, ** = *p* < 0.01, *** = *p* < 0.001



Hillslope characteristics

S. Bachmair and M. Weiler

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 5. Correlation coefficients highlighting the effect of rainfall characteristics and antecedent wetness on RF model performance. Displayed are linear correlation coefficients between RF model performance for different response variables (AREA_NORM, INDEX_ACTI, and INDEX_150) and different rainfall characteristics and antecedent wetness. For model performance per temporal scale see Fig. 4 top graph, for rainfall characteristics and antecedent wetness per temporal scale see Fig. 3 and Table 1. Correlation with total rainfall $n = 12$ (all temporal scales), correlation with rainfall intensity $n = 11$ (seasonal and event scale), correlation with AWI $n = 5$ (event scale only).

Explained variance of RF models for different response variables (%)	Total rainfall (mm)	Rainfall intensity (mm/h)		AWI ^a (mm)
		Mean	Max.	
AREA_NORM	0.30 n.s.	-0.73*	-0.45 n.s.	0.76 n.s.
INDEX_ACTI	0.41 n.s.	-0.52.	-0.39 n.s.	0.24 n.s.
INDEX_150	0.28 n.s.	-0.32 n.s.	-0.24 n.s.	0.67 n.s.

^a Antecedent wetness index: weighted rainfall of the last 14 days
 n.s. (not significant) = $p > 0.1$, . = $p < 0.1$, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$






Fig. 1. Overview on catchment and location of hillslopes (top graph), location of wells within hillslopes (middle graph), and location of mapped trees at the forested hillslopes (bottom graph, DBH= stem diameter at breast height). Catchment area: 0.21 km²; elevation range: 340–585 m a.s.l.; each hillslope (grassland, coniferous forest, and mixed forest hillslope) is equipped with 30 wells.

Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 2. Water table response variable AREA_NORM per hillslope and transect for the entire time series (18 months) and per season (1_2011: Jan/Feb/Mar, 2_2010/2011: Apr/May/June, 3_2010/2011: Jul/Aug/Sep 2010, 4_2010: Oct/Nov/Dec). A cross represents a dry well, a white spot missing data.

Hillslope characteristics

S. Bachmair and M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hillslope characteristics

S. Bachmair and
M. Weiler

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. Rainfall characteristics per season: total rainfall, percentage of time with rainfall, mean rainfall intensity (standard deviation denoted as error bars), and median of rainfall intensity. Mean and median of rainfall intensity are calculated from all time steps with rainfall. 1_2011: Jan/Feb/Mar, 2_2010/2011: Apr/May/Jun, 3_2010/2011: Jul/Aug/Sep 2010, 4_2010: Oct/Nov/Dec.

Hillslope characteristics

S. Bachmair and
M. Weiler




Fig. 4. Explained variance of random forest models for different temporal scales: entire time series, seasonal scale (highlighted in grey), and event scale. Top graph: Explained variance of random forest models predicting the water table response variables AREA_NORM, INDEX_ACTI, and INDEX_150. Bottom graph: Explained variance of random forest models predicting the water table response variable AREA_NORM for all wells ($n = 90$), only the grassland hillslope wells ($n = 30$), and only the forest hillslope wells ($n = 60$). The y-axis is scaled to a minimum of 0, missing bars represent 0% explained variance or negative values.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hillslope characteristics

S. Bachmair and
M. Weiler




Fig. 5. Variable importance plots of random forest models predicting the water table response variable AREA_NORM for different temporal scales (entire time, seasonal scale, and event scale). The x-axis displays the percentage of increase in mean square error (MSE) when the data for that predictor are permuted randomly while all others are held constant. The numbers in the lower right corner of each variable importance plot is the percentage of explained variance for that random forest model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hillslope characteristics

S. Bachmair and
M. Weiler




Fig. 6. Variable importance plots of random forest models predicting the water table response variable INDEX_ACTI for different temporal scales (entire time, seasonal scale, and event scale). The x-axis displays the percentage of increase in mean square error (MSE) when the data for that predictor are permuted randomly while all others are held constant. The numbers in the lower right corner of each variable importance plot is the percentage of explained variance for that random forest model.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

