

Authors' Reply to Editor's second round of comments

Reviewer # 2:

Reviewer: "Furthermore the study primarily focusses on suspended sediment driven water quality constituents like suspended sediment concentration, total nitrogen and total phosphorus but the analysis is restricted to monthly data, hence the most important short term events with high concentrations of the abovementioned compounds are not considered in the study."

You responded: "Furthermore, although targeted sampling of high flow events is very relevant for load estimation of particle-related contaminants, it is not appropriate for state-of-environment monitoring like NRWQN (e.g. Davies-Colley et al. 2011; cited), covering both dissolved and particulate constituents, for which random or pseudo-random (e.g. regular monthly as in the NRWQN) sampling is most appropriate."

My assessment: This argument is only partially convincing. High flow events also matter and are part of the state of the environment. However, low flow conditions (generally well characterized by sufficiently long time series of grab samples and median values) and high flow events have different ecological relevance (see Stamm, Jarvie & Scott 2014 for illustration of my argument). Hence, what your sampling scheme and statistical methods reproduces are conditions that prevail for most of the time in the streams and what the organisms living there experience for most of the time. Therefore, it makes sense to look at these metrics and analyse these trends. However, this is not an argument to disqualify the critique that high flow events are not/only poorly captured by the sampling and statistical strategy. High flow events may be essential for water quality assessment (depending of parameters). This holds on the one hand if you think about downstream systems (including estuaries); on the other hand, it may also be essential for in-stream processes: if you wish to understand for example bed sediments in streams with all their ecological relevance you will hardly be able to do so by only knowing what happens during low flow conditions. Much research has been done illustrating how grab samples may severely underestimate loads of compounds entering streams predominantly during high flows and how difficult it may be to detect trends in time with such a sampling strategy (e.g., Moosmann *et al.* 2005). You have to explicitly mention these aspects in your manuscript and you have to make it clear to the reader what the results actually represent and what not. In the current version this is completely lacking: there is no discussion about sampling effects on results for example.

Authors' response: We added a paragraph at the beginning of the Discussion to acknowledge that we did not capture large floods, the resulting potential uncertainty, and justification for our ability to assess median conditions and trends. We included the references you mention, but did not include any discussion on in-stream processes because it was beyond the scope of our study.

Reviewer: "Furthermore the manuscript is very long (41 pages text only) and not very specific including repetitions."

You responded: "The manuscript is long because of our comprehensive coverage of both spatial and temporal effects of land use on a wide range of river water quality variables in complex large catchments. Arguably, the paper could be split into two manuscripts, but we feel it will have a greater impact as one paper. Further, an understanding of temporal effects is necessary in order to explain some of the spatial effects, and vice versa. We do not understand the comment 'not very specific.' We did a lot of investigation on land use practices and processes that were responsible for the patterns and relationships we observed. Maybe the reviewer is referring to our scale of analysis: catchment-scale. On line 95, we state: "Most of our analyses were performed at

the catchment scale because it integrates the spatiotemporal changes that are reflected in our water quality measurements, it is the appropriate scale to analyze diffuse pollution, and it is the most appropriate spatial management unit (Howard-Williams et al., 2010).”

My assessment: I had a fresh look at the manuscript by reading it carefully once more and very much agree with the reviewer. Being comprehensive is nice but if this leads to a lengthy manuscript that distracts the reader from the essentials it has to be avoided. Being concise is beneficial to both – authors and readers: to the reader because he or she gets the relevant novel information as quickly and clearly as possible, to the author because the readers will like the paper more, which increases the probability of being cited later on. Given the fact that Reviewer # 2 called for a clear focus of the manuscript and your suggested focus and effects of land use *intensity*, I strongly recommend that the result and discussion really concentrate on this aspect. Below, I list some examples of lengthy and repetitive sections and paragraphs that should be avoided:

Authors’ response: We address all of these separately below. We have also gone through the paper carefully and removed other parts of the ms we did not feel was essential. Overall, we have shortened the manuscript considerably.

- L. 281 – 283, 740 – 741 (and elsewhere): These details on which catchments shows what is hardly relevant for the general reader. Only indicate such details if they illustrate an aspect a reader cannot not understand otherwise and which is essential for the manuscript.

Lines 281-283 were removed and the preceding two sentences were combined and condensed. Lines 740-741 were also removed.

- L. 297 – 312: This can be shortened.

We shortened this section by about 3 lines.

- L. 417 – 429: These two paragraphs do not focus on changes in land use intensity, which should be the focus of the manuscript. Hence, they can be massively shortened or even skipped.

This paragraph reports changes (or lack thereof) in important water quality variables from 1989 to 2014, which is a part of the study. Further, when talking about changes in nutrients, it is good to have information on water temperature, DO, conductivity, and pH for context. Thus, we have kept this section.

- L. 450 – 457: This paragraph can also be shortened without loss of information essential to the manuscript.

We shortened this paragraph, removing 4 sentences.

- L. 520 – 547: These two paragraphs do not focus on changes in land use intensity, which should be the focus of the manuscript. Hence, they can be massively shortened.

These paragraphs characterize the states and trends of water quality in NZ rivers, which is a key part of this paper. This Discussion on differences between lowland and upland catchments also sets up the next section on “The role of physiography in dictating land use intensity across NZ”. We feel it is important to leave this section in the paper.

- Section 5.2 (L. 597 – 634): This is not the focus of this manuscript because you focus on land use intensity. You can shorten this part substantially without loss of information.

We changed the title of this section to: The role of physiography in dictating land use intensity across NZ. This section is important because it makes the connections between physiography and land use intensity, which is then expanded on in the next section as regards river water quality.

- L. 536: This is repetitive and can be skipped (L. 405 – 406, L. 520, Tab. 5)
We deleted L536.

- L. 638 – 651: Repetitive, skip.
This entire section was removed.

- L. 667: This is repetitive and can be skipped.
Removed as suggested.

- L. 718 – 720: This is repetitive and can be skipped (L. 507, 654!).
L654 refers to high-producing grasslands, while L718-720 refers to plantation forests. To shorten this sentence, we removed the explanations of negative and positive relationships.

- Section 5.3.4 (L. 749 – 784): Lengthy descriptions of land use effects related to land use categories with little relevance for this study. Without loss of information you can either skip or shorten to 2 – 3 sentences at maximum.
We have removed this entire section.

- L. 846 – 855: This explanation of an outlier in the data set is superfluous. You already presented in quite some detail the reasons for a first outlier (L. 834 – 845). This first case may be included as a proof of concept for what the data set allows for, the second does not add any general insight. The general scientific audience is not interested in all details that may be of relevance for local or regional water managers.

We removed the 2nd paragraph, and added the mentioning of RO3 as another outlier to the previous paragraph.

- L. 898 – 902, 915 – 923: The content of the two respective paragraphs is very redundant and can be significantly shortened without loss of information.

We combined and condensed these two paragraphs.

- L. 907 – 911: This reads like a political statement for an NZ-internal audience and has no link to the actual content of the manuscript.

We removed these sentences.

Reviewer: “... and some conclusions are made without clear evidence.”

My observation: Along the line mentioned in general terms I stumbled across two points I’d like to mention here:

i) On L. 87 – 88 you make a bold statement about the NZ water quality monitoring program (“it has one of the longest comprehensive national water quality datasets in the world”). I suggest that you back this with some information that supports this statement.

We changed the wording to say: it has a long, consistent, and comprehensive national water quality dataset.

ii) ii) Your conclusion regarding the possible effects of increasing water clarity (L. 581 – 595). This is an interesting point but you do not provide evidence for your statement that “when combined with increasing nutrients, warmer water, and lower flows, the perfect recipe for toxic algae blooms is created.” (L. 581 – 583). You cite (McAllister, Wood & Hawes 2016) but these authors seem to contradict your statement by claiming: “While quantitative data on sedimentation rates in rivers is lacking at a national scale, increasing land use intensification and forestry are likely to result in increased sediment in rivers, which may be partly responsible for observed rise in Phormidium proliferations.” (McAllister, Wood & Hawes 2016, p. 292). Please provide references that support your statement.

McAllister’s point in this paragraph is that because most NZ freshwaters are strongly P-limited, increased dissolved reactive phosphorus (DRP; which has a strong association with sediment) leads to Phormidium proliferations (see 3rd sentence in their Conclusions for clarification).

McAllister acknowledges light-limitation on p.287. We have added references to support our claim on L581: Dodds and Welch, 2000; Hilton et al., 2006. We also used these references to back up our claim in the final sentence of this paragraph.

Editor comments:

Already in my initial comments I asked for scatter plots displaying the relationships between discharge and water quality parameters. You argued that this would be an overkill. As a consequence, the entire flow normalization (L. 137) that you do as the first step in the data processing is basically hidden from the reader. As a consequence, your entire section 4.3.1 reports data without showing actual data although you qualitatively describe concentration-discharge relationships. This is not satisfactory and could be easily alleviated. If you combine related water quality parameters (e.g., total P, DRP etc.) together in one plot, you can display these concentration-discharge relationships with 4 to 5 matrices of 9x9 plots for all 77 sites. You have already now plot matrices of 16 x 16 in the Supplementary material.


Therefore, providing scatterplots of the concentration-discharge relationships with the LOESS functions is doable and will provide the reader with essential access to real data. This will improve the quality of the manuscript substantially and will make the entire process of data processing much more transparent. I strongly suggest that you add this information.

Now that the paper is focused on land use intensity, we have removed section 4.3.1 (L368-403) entirely. Flow-normalization using LOESS is a commonly used technique and we don’t think the 770 plots (with 312 data points each) would add any value to the paper, especially since section 4.3.1 has been removed. If we combined two variables on one plot, that would still be 385 plots, but now with 624 data points each, which would look like a buckshot. For the Editor’s benefit, we have added a supplement below that shows the LOESS plots for just one catchment, AK1. We would have to produce this group of 10 plots 76 more times to capture all the catchments.

Because your focus is on the effects of intensity change you might consider to motivate the issue by actually including a figure in the main text that illustrates that land use did hardly change between 1990 and 2012 but that intensity did (based on Tab. 2 & 3 in the Supplementary Material).

This is a great idea. We have added this figure to the ms, now Figure 2.

Author supplement 1: LOESS plots for catchment AK1



1 **River water quality changes in New Zealand over 26 years (1989 – 2014): Response to land**
2 **use intensity and land disturbance**

3
4 Jason P. Julian*^{1,5}, Kirsten M. de Beurs^{2,5}, Braden Owsley^{2,5}, Robert J. Davies-Colley³, Anne-
5 Gaelle E. Ausseil⁴

6 ¹Department of Geography, Texas State University, San Marcos, TX, USA

7 ²Department of Geography and Environmental Sustainability, The University of Oklahoma,
8 Norman, OK, USA

9 ³National Institute of Water and Atmospheric Research Ltd (NIWA), Hamilton, New Zealand

10 ⁴Landcare Research, Palmerston North, New Zealand

11 ⁵Landscape & Land Use Change Institute (LLUCI), <http://tethys.dges.ou.edu/main/>, USA

12 *Corresponding author: Jason.Julian@txstate.edu

13

14 Abstract

15 Land use-water quality relationships are complex with interdependencies, feedbacks, and legacy
16 effects. Most river water quality studies have assessed catchment land use as areal coverage, but
17 here, we hypothesize and test whether land use *intensity* – the inputs (fertilizer, livestock) and
18 activities (vegetation removal) of land use – is a better predictor of environmental impact. We
19 use New Zealand as a case study because it has had one of the highest rates of agricultural land
20 intensification globally over recent decades. We interpreted water quality state and trends for the
21 26 years from 1989 to 2014 in the National Rivers Water Quality Network (NRWQN) –
22 consisting of 77 sites on 35 mostly large river systems with an aggregate catchment amounting to
23 half of NZ's land area. To characterize land use intensity, we analyzed spatial and temporal

24 changes in livestock density and land disturbance (i.e. bare soil resulting from vegetation loss by
25 either grazing or forest harvesting) at the catchment-scale, as well as fertilizer inputs at the
26 national scale. Using simple multivariate statistical analyses across the 77 catchments, we found
27 that visual water clarity was best predicted by areal coverage of high-producing pastures. The
28 primary predictor for all four nutrient variables, however, was cattle density, with plantation
29 forest coverage as the secondary predictor variable. While land disturbance was not itself a
30 strong predictor of water quality, it did help explain outliers of land use-water quality
31 relationships. From 1990 to 2014, visual clarity significantly improved in 34/77 catchments,
32 which we attribute mainly to increased dairy cattle exclusion from rivers (despite dairy
33 expansion) and the considerable decrease in sheep numbers across the NZ landscape, from 58
34 million sheep in 1990 to 31 million in 2012. Nutrient concentrations increased in many of NZ's
35 rivers with dissolved oxidized nitrogen significantly increasing in 27/77 catchments, which we
36 largely attribute to increased cattle density and legacy nutrients that have built up on high-
37 producing grasslands and plantation forests since the 1950s and are slowly leaking to the rivers.
38 Despite recent improvements in water quality for some NZ rivers, these legacy nutrients and
39 continued agricultural intensification are expected to pose broad-scale environmental problems
40 for decades to come.

41

42 1. Introduction

43 River water quality reflects multiple activities and processes~~all that has happened~~ within
44 its catchment, including geomorphic processes, vegetation characteristics, climate, and
45 anthropogenic land uses (Brierley, 2010). Relationships between water quality and these
46 catchment characteristics are not straightforward because all of these factors interact over both
47 space and time. For example, if intensive livestock grazing occurs on steep slopes, surface runoff

48 and consequently river turbidity is expected to be greater than if grazing occurs on flatter areas.
49 Or if fertilizers are heavily applied to sandy soils with high drainage density, rivers will likely
50 become eutrophied over a period of decades due to legacy nutrients slowly leaking to the rivers
51 through groundwater ([McDowell et al., 2008](#)). The influence of land use on water quality has
52 also been shown to vary among different climates (Larned et al., 2004). With all of the various
53 types of intensive land uses that have occurred across diverse landscapes over hundreds of years,
54 rivers with degraded water quality are now widespread.

55 Historically, water quality in rivers was managed to meet minimally [acceptable](#) standards
56 [or maximum pollutant load limits](#) (Baron et al., 2002; [Boesch, 2002](#); [Howard-Williams et al.,](#)
57 [2010](#)). However, in the last decade, a greater emphasis has been placed on maximizing the
58 ecosystem services provided by healthy rivers, which is driving efforts to [further](#) improve water
59 quality (Brauman et al., 2007; Davies-Colley, 2013). Early efforts in developed countries to
60 improve water quality focused on point-source pollution, particularly wastewater discharges
61 from factories and treatment plants (Campbell et al., 2004). While the broad-scale reduction in
62 point-source pollution elevated many water quality variables above minimal standards, most
63 rivers globally still have water quality impairments due to diffuse pollution – fine sediments,
64 nutrients, pathogens, toxicants, salts, and other contaminants that are delivered from unknown or
65 many indistinguishable sources across the catchment (Vorosmarty et al., 2010). [Although](#)
66 [considerable effort has been directed at monitoring and reducing diffuse pollution with some](#)
67 [success, the legacy of pollutants from various land uses remains](#) ([Boesch, 2002](#); [Kronvang et al.,](#)
68 [2008](#); [Zobrist and Reichert, 2006](#)). Agricultural land uses are by far the greatest contributors of
69 diffuse pollution, globally (Foley et al., 2005; Vitousek et al., 1997**b**); however, the ‘intangible’

70 sources of diffuse pollution make it difficult to assign cause-and-effect relationships (Campbell
71 et al. 2004).

72 ~~Many~~Most studies have used theoretical or numerical models to ~~that have examined~~
73 relationships between land use and water quality ~~have used theoretical or numerical models~~
74 because of the lack of consistent water quality ~~monitoring data~~ over long periods (bracketing land
75 use change). While ~~this practice~~modelling approaches can be useful for small catchments where
76 much is known about its landscape, modelling may not work well for larger catchments because
77 land-water relationships are complex with interdependencies, feedbacks, and legacy effects.
78 Empirical studies can shed light on some of these complexities, but they are only useful for their
79 particular catchments and may have limited generality or transferability. Comparisons of many
80 diverse catchments is probably most useful to advance understanding of broad-scale land-water
81 relationships (Zobrist and Reichert, 2006).

82 One of the most comprehensive empirical ~~riverine~~multi-catchment studies to date on
83 land use-water quality relationships has been Varanka and Luoto's (2012) study of 32 boreal
84 rivers in Finland. They analyzed five water quality variables over ~~10~~ten years as a function of a
85 suite of physiographic, climate, and land use variables. A similar study was conducted on many
86 of the same rivers in Finland, but with a more sophisticated temporal analysis (Ekholm et al.,
87 2015). And several other studies have used this same river water quality dataset to investigate
88 environmental drivers. In a study of 11 Swiss watersheds, Zobrist and Reichert (2006) analyzed
89 export coefficients of six water quality variables from biweekly, flow proportional, composite
90 samples over a 24-year period within the context of land use.

91 All of these studies, and most catchment land use studies, assessed land use (or land use
92 change) as areal coverage. However, land use intensity – the inputs (e.g. fertilizer, livestock) and

93 activities (e.g. vegetation removal) of land use – could be a better predictor of environmental
94 impact for being a more direct measure of impact than areal coverage (Blüthgen et al., 2012;
95 Ramankutty et al., 2006). Unfortunately, our understanding of the patterns, processes, and
96 impacts of land use intensity is inadequate because of (1) its complex, multidimensional
97 interactions with other landscape variables, and (2) the lack of appropriate datasets across broad
98 spatiotemporal scales (Kuemmerle et al., 2013; Erb et al., 2016). New Zealand (NZ) provides a
99 valuable test-bed for the patterns, processes, and impacts of land use intensity because over the
100 past three decades pasture area has decreased but livestock densities and fertilizer inputs have
101 increased (MacLeod and Moller, 2006; StatsNZ, 2015). Like Finland and Switzerland, New
102 Zealand (NZ) has an extensive long-term river water quality monitoring network, which has
103 allowed many studies on river water quality state and trends (Smith et al., 1996, 1997;
104 Scarsbrook et al., 2003; Scarsbrook, 2006; Ballantine and Davies-Colley, 2014) and effects of
105 land use areal coverage (Davies-Colley, 2013; Larned et al., 2004, 2016). However, this dataset
106 has not been assessed as regards changes in land use intensity that have occurred over the same
107 period.

108 Here, we ~~use NZ as a case study to illustrate~~ investigate long-term relationships among
109 land use intensity~~management~~, geomorphic processes, and river water quality in NZ – which
110 provides a particularly valuable case study because: (1) it has had one of the highest rates of
111 agricultural land intensificationties over recent decades and thus serves as a potential indicator
112 for ~~some developing~~ countries that are also increasing agricultural intensity; (2) it has ~~one of the~~
113 longest, consistent, and comprehensive national water quality datasets ~~in the world~~; and (3) it is
114 physiographically-diverse. We examined monthly data for a suite of water quality variables over
115 a 26-year period for 77 ~~very~~ diverse catchments. We then compared these states and trends of

116 river water quality to landscape data that characterized the catchments' geomorphology, soil
117 properties, and hydro-climatology; as well as temporal changes in land use areal coverage and
118 land use intensity, specifically livestock density and land disturbance, defined here as bare soil
119 resulting from vegetation loss. ~~of these catchments. We also assessed temporal changes in land~~
120 ~~cover/use, livestock, and land disturbance over our study period and compared these to temporal~~
121 ~~changes in water quality variables.~~ Altogether, these analyses ~~illustrated~~ reveal coincident
122 spatiotemporal patterns in land use intensity and water quality ~~in NZ rivers~~ over a quarter of a
123 century. Most of our analyses were performed at the catchment scale ~~because it~~ which integrates
124 the spatiotemporal changes that are reflected in our water quality measurements, ~~it~~ is the
125 appropriate scale to analyze diffuse pollution, and ~~it~~ is the most appropriate spatial management
126 unit (Howard-Williams et al., 2010).

127

128 2. Study area

129 New Zealand; (*Aotearoa*, “Land of the long white cloud” in the language of indigenous
130 *Maori* people); is a small island nation (~268,000 km²) located between the South Pacific Ocean
131 to the east and the Tasman Sea to the west. Its two main islands (North Island and South Island)
132 are located between 34° and 47° S latitude. Being located on the active boundary between the
133 Australian and Pacific Plates, NZ’s geology and geomorphology are very diverse, including
134 active volcanoes, karst regions, a range of high fold mountains (the Southern Alps), large coastal
135 plains, and rolling hills across both hard- and soft-rocks. Being stretched latitudinally, with
136 nowhere more than about 150 km from the sea, between two major ocean waters combined with
137 its topographic variability, NZ also has a diverse climate with regional extremes, including sub-
138 tropical in the far north, temperate in the central North Island, extremely wet on the western side

139 of the Southern Alps (up to 10 m annually), and semi-arid in the rain shadow to the east of the
140 Southern Alps.

141 New Zealand is the last major habitable landmass to be settled by humans. Eastern
142 Polynesians first arrived around 1300 AD (Wilmshurst et al., 2008). Europeans first arrived in
143 the late-1700s, but large-scale settlement did not begin until the 1840s. Broad-scale agriculture
144 spread shortly after and has been intensifying since. While we address land use changes at the
145 national scale in this study, our water quality analyses focus on 77 diverse catchments across NZ
146 (Fig. 1), which cumulatively cover about half of NZ's land area.

147

148 3. Methods

149 3.1. Water quality data

150 Water quality data was obtained from NZ's National Rivers Water Quality Network
151 (NRWQN), which is operated and maintained by the National Institute of Water & Atmospheric
152 Research (NIWA). This network represents one of the world's most comprehensive river water
153 quality datasets: thirteen water quality and two biomonitoring variables have been measured
154 monthly (via in situ measurements and grab samples), with supporting flow estimation, from
155 1989-2014 at 77 sites whose catchments cumulatively drain approximately half of New
156 Zealand's land surface (Davies-Colley et al., 2011). Further, this dataset has been operationally
157 stable throughout its history, which allows us to calculate trends over this period. For this study,
158 we focused on eleven water quality variables and their coincident flow (Table 1). We did not
159 analyze ammoniacal nitrogen (NH₄) because early NH₄ samples were biased high by laboratory
160 contamination (Davies-Colley et al., 2011).

161 All water quality variables, except water temperature (T_w), were flow-normalized (for
162 each site separately) in JMP® Pro (v 11.2.1) with local polynomial regression (LOESS) using a
163 quadratic fit, a tri-cube weighting function, a smoothing window (alpha) of 0.67, and a four-pass
164 robustness to minimize the weights of outliers (Cleveland and Devlin, 1988); where, flow-
165 adjusted value = raw value – LOESS value + median value. With LOESS, there is no assumption
166 about the water quality variable’s relationship with flow. For example, although visual clarity
167 usually decreases systematically with increasing flow (Smith et al., 1997), algae blooms at low
168 flows can sometimes reduce clarity. LOESS also allowed us to examine relative water quality
169 changes over long periods.

170

171 3.2. Physiographic data

172 Water quality metrics and trends were compared to a suite of landscape variables (Table
173 2). Catchment morphometrics (area, slope, ruggedness) were obtained from a 30-m digital
174 elevation model (DEM) that we rescaled (in order to align with other gridded spatial datasets)
175 from the 25-m DEM produced by Landcare Research. This 25-m DEM was interpolated from
176 20-m contours of the national TOPOBASE digital topographic dataset supplied by Land
177 Information New Zealand (LINZ; scale: 1:50,000). Catchment area (A) is the drainage area (in
178 km^2) above the NRWQN station, derived using Arc Hydro tools in ArcGIS 9.3.1 in combination
179 with the River Environment Classification (REC, v2.0), the national hydrography dataset derived
180 from a 30-m hydrologically correct DEM produced by NIWA (Snelder et al., 2010). Mean
181 catchment slope (S_c) was derived from the same software package, using a 3x3 cell window. We
182 defined ruggedness (R_r) as the standard deviation of the 30-m slope grid for each catchment

183 (*sensu* Grohmann et al., 2011). Drainage density (D_d) was calculated from the ratio of the total
184 length of REC streams ~~over~~to catchment area (in km/km²).

185 Soils data was obtained from the 1:50,000 Fundamental Soils Layers (FSL), which is
186 maintained by Landcare Research. Methods and data descriptions for this soils database are
187 described in Webb and Wilson (1995) and Newsome et al. (2008). Catchment-scale soil
188 variables (mean value across catchment) that we included in our analysis for being expected to
189 be related to water quality were: soil depth (Z_s), percent of catchment dominated by silty and
190 clayey surface soils ($SC\%$), soil pH (pH_s), cation exchange capacity (CEC), organic matter
191 percentage ($OM\%$), and phosphate retention (P_{ret}). Phosphate retention is a measure (in %) of the
192 amount of phosphate that is removed from solution by the soil via sorption ([Saunders, 1965](#)).
193 Thus, soils with high P_{ret} have low P-availability for plant growth.

194 Median annual precipitation (MAP), median annual temperature (MAT), and median
195 annual sunshine (MAS) averaged across each catchment was obtained from NIWA's National
196 Climate Database, which contained 5-km gridded daily weather data (Tait and Turner, 2005).
197 Our values for these three variables represent the median annual precipitation (total mm/y),
198 temperature (mean °C), and sunshine (hours/y) for the period 1981-2010. Relative water storage
199 (RWS) was calculated as the proportion of the annual catchment water yield (i.e. total volume of
200 water leaving the catchment in a year) stored in lakes and reservoirs. Reservoir/lake storage was
201 obtained from the Freshwater Ecosystems of New Zealand (FENZ) Database, described in
202 Snelder (2006). The last hydro-climatological variable we included in our analyses was the
203 median discharge (Q_{50}), which was calculated from the NRWQN 'flow stamping' at times of
204 water quality sampling from 1989-2014.

205

206 3.3 Land use areal coverage, intensity, and disturbance data

207 There are two national land use datasets for New Zealand. The Land-Use and Carbon
208 Analysis System (LUCAS) was developed by the NZ Ministry for the Environment (MfE, 2012)
209 for reporting and accounting of carbon fluxes and greenhouse gas emissions, as required by the
210 United Nations Framework on Climate Change and the Kyoto Protocol. Accordingly, LUCAS
211 uses 1990 as its reference year and maps land use in 12 classes for 2008 and 2012 ~~as well for 12~~
212 ~~classes~~. The Land Cover Database (LCDB) was developed by Landcare Research (LCR), with
213 contributions from MfE, Department of Conservation (DOC), Ministry for Primary Industries
214 (MPI), and Regional Councils (LCR, 2015). LCDB contains 35 land use classes for 1996, 2001,
215 2008, and 2012. Both datasets use a minimum mapping area of 1 hectare, and use many of the
216 same data and methods to map land use. There are however, some key differences in their class
217 designations and classifications that are important to our analyses: (1) LUCAS includes
218 Manuka/Kanuka as forest, whereas LCDB designates Manuka/Kanuka as shrub; (2) LUCAS
219 lumps all post-1989 forests into one class, whereas LCDB differentiates between indigenous and
220 plantation forests; (3) LUCAS uses a conservative approach to mapping high-producing
221 grasslands, whereas LCDB uses phenological information to provide more accurate estimations
222 of high-producing grassland. Because of our focus on (water quality-impacting) plantation
223 forests and high-producing grasslands, we used d the LCDB (v4.1) for our spatial and statistical
224 analyses. We used d LUCAS only to quantify long-term changes from 1990 to 2012, before the
225 LCDB was initiated in 1996. Table 3 describes the land use classes we used in this research,
226 which classes are included from both datasets, and the national comparison between LUCAS and
227 LCDB for 2012.

228 There are numerous metrics for land use intensity (Erb et al., 2013). At the catchment-
229 scale, we used livestock density as a metric for all grasslands; and we used land disturbance,
230 defined here as bare soil resulting from vegetation loss, as a metric for high-producing grasslands
231 and plantation forests. We also used national-scale annual fertilizer data (1989-2014) from
232 StatsNZ (2015) to compare long-term trends of river nutrient concentrations to nutrient inputs.
233 Livestock numbers for dairy cattle, beef cattle, sheep, and deer (at 1 ha resolution) for each
234 catchment were derived from maps provided by Ausseil et al. (2013), which is representative for
235 the year 2011. To assess total livestock impact on land disturbance, we multiplied each livestock
236 type by its AgriBase stock unit (SU) coefficient: sheep = 0.95 SU, deer = 1.9 SU, beef cattle =
237 5.3 SU, and dairy cattle = 6.65 SU (Woods et al., 2006). The total SU for each catchment was
238 then normalized by total catchment area, expressed as stock unit density (*SUD*) in SU/ha.

239 Changes in *SUD* from 1990 to 2012 (*SUD*₂₀₁₂₋₁₉₉₀) were assessed using district-level data
240 from StatsNZ (2015) on total numbers of sheep, deer, beef cattle, and dairy cattle. These
241 livestock numbers were then aggregated for each catchment and multiplied by their respective
242 SU coefficient. Stock units per hectare were then compared between 1990 and 2012 to assess
243 change in livestock impacts in each catchment. For Whakatane and Kawerau Districts, 1993 was
244 used because 1990 data was unavailable.

245 Land disturbance (i.e. bare soil) resulting from vegetation loss was quantified for all
246 high-producing grasslands (*D_{HG}*) and plantation forests (*D_{PF}*), as well as the whole catchment
247 (*D_C*) for the period 2000 - 2013. The methods for calculating and validating disturbance are
248 described in de Beurs et al. (2016). Briefly, MODIS BRDF corrected reflectance data
249 (MCD43A4) at 463 m spatial resolution and eight day temporal resolution was used to calculate
250 Tasseled Cap brightness, greenness and wetness based on the coefficients following Lobser and

251 Cohen (2007). These indices consist of linear combinations of all seven MODIS reflectance
252 bands to represent general image brightness which is comparable to albedo, image greenness
253 which is comparable to the better known vegetation indices such as NDVI and EVI, and image
254 wetness which is linked to the amount of water captured in the vegetation, most comparable to
255 normalized difference water indices. Missing pixels were ignored. We then calculated the mean
256 and standard deviation of each tasseled cap index for each combination of land cover class (LCR,
257 2015) and climatic region for each 8-day time period. We then used these measures to
258 standardize the calculated tasseled cap indices. To determine how disturbed each pixel was at
259 any point in time, we then calculated the forest and grassland disturbances. The forest
260 disturbance index is calculated as the standardized brightness minus the standardized greenness
261 and wetness. The idea is that disturbed forests appear brighter and less green and less wet than
262 undisturbed forests. The grassland index is the negative sum of all indices, indicating that
263 disturbed grasslands appear darker, less green and less wet than undisturbed grasslands. MODIS
264 disturbance data were visually validated against 7500 random pixels from Landsat imagery and
265 corresponding 15 high resolution Orbview-3 and Ikonos images. The overall accuracy of the
266 disturbance index based on Landsat data was 98%.

267

268 3.4 Statistical methods

269 We used nonparametric Spearman rank correlation coefficients (r_s) ~~instead of actual~~
270 ~~values~~ to look at relationships between variables; because many of the relationships ~~are~~ were
271 curvilinear. Statistical significance was taken to be an alpha of 0.05. Bivariate comparisons
272 between all variables (Tables 1-3) were performed to explore for associations and identify
273 correlated variables before later multivariate analyses. Median values (from the 26-y monthly

274 time-series) for water quality variables at each site were used when compared to physiographic
275 and land use variables of their corresponding catchment. Stepwise regression was then used to
276 rank-order the relative contributions of multiple landscape variables associated with each major
277 water quality variable. Stepwise regression was used because it accounts for correlations among
278 the independent landscape variables. The order of variables in the stepwise regression model and
279 the sign of their coefficient (proportional [+] vs. inverse [-]) provides an objective measure of the
280 contribution of each landscape variable to river water quality. The level of entry into the model
281 was set to $p = 0.05$. All the above statistical analyses were performed in JMP® Pro (v 11.2.1).

282 Temporal trends in water quality (1989 – 2014) and disturbance (2000 – 2013) data
283 were assessed with the seasonal Kendall test which was corrected for temporal autocorrelation
284 using the rkt R package; missing values were ignored. We also calculated the Seasonal Kendall
285 slope estimators (SKSE) using the same R package. Because some NRWQN sites had multiple
286 measurements in some months, a few records (no more than five) were removed from each site
287 in order to ensure 12 monthly values for each year for the SKSE test. There were also occasional
288 missing values for some variables throughout the time-series, particularly in the early years. Of
289 particular note, there were no *TN* values for 1994 as a result of contamination by leaking
290 ammonia refrigerant during storage of frozen subsamples. HV1 did not have data for 18 months
291 from 2012-2014.

292 In order to make trend comparisons among sites and derive an estimate of percent change
293 per year, we normalized SKSE values by dividing them by the raw data median to give the
294 relative SKSE (RSKSE) in percent change per year (Smith et al., 1996). Given that water
295 temperature (T_w) uses an arbitrary scale in °C, we only report SKSE values for this variable. We
296 also used the trend categories of Scarsbrook (2006): (1) no significant trend – the null hypothesis

297 for the Seasonal Kendall test was not rejected ($p > 0.05$); (2) significant increase/decrease – the
298 null hypothesis for the Seasonal Kendall test was rejected ($p < 0.05$); and (3) ‘meaningful’
299 increase/decrease – the trend was significant, and the magnitude of the trend (RSKSE) was
300 greater than 1% per year. According to Ballantine and Davies-Colley (2014), a 1% change per
301 year translates to slightly more than 10% change per decade (due to compounding), a rate of
302 change that is easily detectable and observable.

303

304 4. Results

305 4.1. Physiographic characteristics

306 The 77 NRWQN catchments were physiographically diverse in terms of morphometric,
307 soil, and hydro-climatological variables (Table 4; Supplement Table 1). Most notable with
308 regards to its direct influence on runoff and water quality was median annual precipitation
309 (*MAP*), which ranged from 533 to 7,044 mm/y. When combined with the wide range of
310 catchment areas (*A*), median discharge (Q_{50}) varied over three orders of magnitude, from 0.4 to
311 515 m³/s, and annual water yield from 103 to 3,475 mm/y. In terms of soil, about a quarter of the
312 catchments had very sandy surface soils ($SC\% < 10$) and a quarter had fine-textured soils ($SC\%$
313 > 70). Phosphate retention (P_{ret}), an important variable for fertilizer management and
314 consequently water quality, was particularly high ($>57\%$; 10th percentile) for catchments HM2,
315 HM5, HM6, WA1, WA2, WA3, and WN5.

316 Several physiographic variables (Table 2) displayed strong latitudinal trends from North
317 to South (r_s): ~~*MAT* (-0.83), *MAS* (-0.61), *R_r* (-0.58), *Z_s* (-0.57), and *P_{ret}* (-0.52). M and many of
318 the physiographic variables were strongly correlated ($p < 0.001$; Supplement Fig. 1). Notable
319 ones include (r_s): ~~*A* v *Q₅₀* (0.89), *S_e* v *D_d* (-0.79), *R_r* v *S_e* (-0.67), *Q₅₀* v *R_r* (-0.57), *RWS* v *Q₅₀*~~~~

320 ~~(0.55), $RWS \ v \ A$ (0.54), $R_s \ v \ D_d$ (-0.52), $OM\% \ v \ Z_s$ (0.47), $MAP \ v \ S_e$ (0.47), $Z_s \ v \ S_e$ (-0.42), $Z_s \ v$~~
321 ~~$SC\%$ (-0.41), $P_{ret} \ v \ pH_s$ (0.40), $MAP \ v \ P_{ret}$ (0.39), and $MAT \ v \ OM\%$ (0.38).~~ In consideration of
322 these relationships and perceived importance for water quality (*sensu* Varanka and Luoto, 2012),
323 we used the following subset of minimally correlated physiographic variables for subsequent
324 multivariate analyses: catchment slope (S_c), silt-clay percentage ($SC\%$), phosphate retention
325 (P_{ret}), and median flow (Q_{50}).

327 4.2. Land use areal coverage and temporal changes ~~and disturbance~~

328 Land use in NZ, like physiography, varied widely; and our 77 catchments captured this
329 diversity (Fig. 1; Supplement Table 2). Thirteen catchments were dominated (>50%) by non-
330 plantation forests (NF), with one (WN2) containing more than 94%. Thirteen other catchments
331 were dominated by shrub/grassland (SG) that was not intensively managed. The most dominant
332 land use was grasslands that were intensively managed (hereafter high-producing grasslands;
333 HG), covering the majority of the area for 31 catchments. Together, these three land uses made
334 up 84% of the catchments' areas. Plantation forest (PF) was the majority land use for three
335 catchments: ~~(RO3, RO5, and RO2), all in the volcanic plateau of central North Island.~~ Open
336 water (OW) was the majority land use for one catchment (RO1) and relatively high (>10%) for
337 two others (RO6, DN10). Barren/other (BO), which was largely bare rock, was relatively high
338 (>10%) for 13 mountainous catchments. Urban (UR) coverage rarely exceeded 1%, with only
339 one catchment greater than 2% (WN1). Annual cropland (AC) exceeded 1% in 11 catchments,
340 but never exceeded 8%. Vegetated wetland (VW) and perennial cropland (PC) were minimal in
341 all catchments, each rarely exceeding 1%.

342 In general, ~~non-plantation forest (NF)~~, ~~shrub/grassland (SG)~~ and ~~barren (BO)~~ areas
343 dominated mountainous catchments with high S_c and low Z_s ; while ~~high-producing grasslands~~
344 ~~(HG)~~ dominated most lowland catchments with low S_c , high Z_s , and high pH_s . Like ~~HG~~,
345 ~~plantation forest (PF)~~ mostly occurred on flat areas ($r_s = -0.48$ with S_c) with thick soils (0.35
346 with Z_s) that were less acidic (0.31 with pH_s). ~~PF was also significantly proportional to P_{ret} ($r_s =$~~
347 ~~0.24)~~. Given the relative dominance of catchment land use, relationships with physiographic
348 variables, and potential effects on water quality in NZ rivers (Davies-Colley, 2013; Howard-
349 Williams et al., 2010), the land use variables used for subsequent multivariate analyses were ~~NF~~,
350 ~~SG~~, ~~HG~~, ~~PF~~, and ~~OW~~.

351 Land use areal coverage did not change much in the 77 catchments from 1990 to 2012
352 across NZ (Fig. 2) or in many catchments ~~was usually minor~~ (Supplement Table 2). The greatest
353 change was a 13.4% increase in ~~PF~~ in GS1, which was almost entirely accounted for by a 13%
354 decrease in ~~SG~~. Thirteen other catchments experienced small increases (3.0 - 6.6%) in ~~PF~~,
355 accounted for by decreases in ~~SG~~ or ~~HG~~ or both. HM3 and HM4 had the greatest increases in
356 ~~HG~~ at 3.4% and 2.0%, respectively. High-producing grasslands (~~HG~~) for the other 75 catchments
357 remained virtually unchanged ($< 0.4\%$) or decreased. WH3 had the greatest decrease in ~~HG~~ at -
358 4.8%. Land use areal coverage change in other catchments was negligible.

359

360 4.3. Land use intensity and temporal changes

361 Changes in total stock unit density between 1990 and 2012 ($SUD_{2012-1990}$) were also
362 minor with only two catchments (AK1 and AK2: both -5.1 SU/ha owing to urban fringe
363 expansion) changing more than 1.6 SU/ha over this period (Supplement Table 3). Temporal
364 changes in $SUD_{2012-1990}$ for 56 of the 77 catchments were within the range of -1.0 to 1.0 SU/ha.

365 Although land use areal coverage and total livestock densities changed little ~~in~~ 1990-2012,
366 livestock *types* changed considerably for many catchments (Supplement Table 3) and across NZ
367 (Fig. 2). The general pattern was dairy cattle replacing sheep. The number of dairy cattle from
368 1990 to 2012 increased in 72 catchments, with a mean increase of 0.6 SU/ha for all catchments;
369 while the number of sheep decreased in all 77 catchments (mean = -0.9 SU/ha). Deer and beef
370 cattle numbers changed little: 0.0 and -0.2 SU/ha, respectively.

371 When 2011 livestock densities were compared with physiographic variables, the
372 strongest relationships were found with combined *SUD* of dairy and beef cattle (hereafter
373 *SUD_{cattle}*; Supplement Fig. 2). *SUD_{cattle}* decreased strongly with increasing slope, S_c ($r_s = -0.79$),
374 but increased with Z_s (0.43), pH_s (0.32), and P_{ret} (0.27). *SUD_{cattle}* also increased with *MAT*
375 (0.68) and *MAS* (0.42), but decreased with *MAP* (-0.34). Thus, highest cattle densities were
376 found in catchments such as WA3 (with the highest *SUD_{cattle}* at 15.7 SU/ha) that were relatively
377 flat, warm, sunny, and dry, with deep soils that had relatively high pH and high P-retention.
378 High-producing grasslands (*HG*) had similar, but less strong, correlations with these same
379 physiographic variables.

380 Catchment disturbance (D_C) varied widely over both space and time between 2000 and
381 2013 (Supplement Table 4). The maximum amount of D_C at one time was 35.7% for WN3 on
382 07-Apr-2003, almost entirely due to bare pastures. D_C exceeded 15% on six other occasions (264
383 days in total) in this catchment. In general, the North Island (Fig. 23) had a greater extent and
384 intensity of disturbance than the South Island (Fig. 34). The most intense disturbances occurred
385 as a result of plantation forest harvests, and these disturbances were on average visible for about
386 1.5 y up to about 4 y, with exceptions lasting more than 6 y. Indeed, D_C was strongly correlated
387 to *PF* coverage ($r_s = 0.51$). The catchment with the highest median D_C (10.5%) was RO3, which

388 had 69.8% of its catchment in *PF* and 17.7% in *HG*. Fourteen other catchments had D_C above
389 5%, and two-thirds of these were dominated by either *PF* or *HG*.

390 We also analyzed disturbance of plantation forests (D_{PF}) and high-producing grasslands
391 (D_{HG}) separately for each catchment. For catchments with at least 21.4-km² (100 MODIS pixels,
392 for the sake of statistical robustness) of plantation forest, the mean (\pm SD) D_{PF} (from 2000 to
393 2013) was $10.6 \pm 5.6\%$. The catchments with the highest D_{PF} were those with low mean annual
394 precipitation, MAP ($r_s = -0.42$). There were no significant relationships between D_{PF} and any of
395 the other physiographic variables. For catchments with at least 21.4-km² of high-producing
396 grasslands, the mean (\pm SD) D_{HG} was $6.0 \pm 6.4\%$. The catchments with the highest D_{HG} were
397 those with low mean annual sunshine (*MAS*; $r_s = -0.25$), low mean annual temperature (*MAT*; -
398 0.30), high catchment slope (S_c ; 0.25), and high ruggedness (R_r ; 0.31). The six catchments with
399 the highest D_{HG} (>15%) all had low phosphate retention (P_{ret} ; <32%). While it is assumed that
400 greater densities of livestock lead to greater pasture disturbance, we did not find a proportional
401 relationship between stock unit density (*SUD*) and D_{HG} ~~across space (i.e. among catchments)~~. In
402 fact, the highest median D_{HG} was found for catchments with *low SUD* ($r_s = -0.45$). Over time
403 however, we observed a fairly strong trend ($r_s = 0.50$) of lower D_{HG} with decreasing *SUD* (-
404 $SUD_{2012-1990}$). In all there were seven catchments with significant or meaningful decreases in
405 D_{HG} from 2000 to 2013 (assessed with Seasonal Kendall slope; SKSE), all of which had a
406 negative $SUD_{2012-1990}$.

407

408 4.3.4.4. Water quality characteristics and trends

409 4.3.1 Flow relationships ———

410 All water quality variables (per site) had strong relationships with flow (Q) except water
411 temperature (T_w), which instead followed a seasonal pattern. Conductivity ($COND$) generally
412 decreased with Q for most sites, with exceptions being AX1, DN2, DN10, NN5, RO1, RO6, and
413 TK1. For several sites, $COND$ was high for flood flows. Water pH (pH_w) decreased with Q for
414 most sites likely due to relatively acidic rainfall, with exceptions being AX3, AX4, RO5, and
415 RO6. Several sites experienced high pH_w during high flows. The typical pattern for dissolved
416 oxygen (DO) for most sites was a wide range at low flows, and high flows converging to near
417 100% DO . The exceptions were sites where DO decreased with flow (DN1, HM4, HM5, WH4)
418 and lake fed sites where DO was high (>90%) for virtually all flows (AX2, AX4, DN4, D10,
419 RO1, RO6, TK4).

420 Visual clarity ($CLAR$) had a strong (mean r^2 of 0.53 among all sites) exponential decay
421 trend with flow for almost all sites, as has been reported previously (Smith et al., 1997). Four
422 sites, all lake fed, had their highest $CLAR$ for intermediate flows (DN10, RO1, RO6, HM3). Of
423 these four, the first three had high $CLAR$ (> 2m) for virtually all flows. Turbidity ($TURB$) had
424 generally the opposite trend of $CLAR$ (as could be expected given the inverse relationship of
425 these variables), and increased near linearly with Q (albeit with more scatter than $CLAR$).
426 Several of the lake fed sites had relatively low $TURB$ at high flows (AX1, AX2, DN1, DN10,
427 RO1, RO2, RO6). Colored dissolved organic matter ($CDOM$) generally increased with flow as
428 has been reported previously by Smith et al. (1997); the lake fed sites of RO1, RO2, and RO6
429 were exceptions. $CDOM$ was sometimes low during floods, likely due to a dilution effect.

430 Total nitrogen (TN) generally increased with Q , but with a high degree of scatter, for
431 almost all sites. The exceptions were AX1, AX2, AX4, DN10, RO1, RO6, and TK4, where TN
432 was low for all flows, usually less than 100 mg/m³. The trends of oxidized nitrogen (NO_x) with

433 ~~flow varied widely among the sites. For many sites (26/77), NO_x increased with Q , usually with~~
434 ~~a positive logarithmic trend (i.e. asymptotes at high flows) due to dilution effects at high flows.~~
435 ~~A couple sites displayed a concave upward parabolic trend where NO_x concentrations were~~
436 ~~lowest for intermediate flows and high for both low and high flows (CH2, DN6), which we were~~
437 ~~unable to explain but is likely due to source of flow. Total phosphorous (TP) generally increased~~
438 ~~with Q at 73 of the sites, reflecting mobilization of suspended matter (containing P) with Q .~~
439 ~~Exceptions were the lake fed sites of DN10, RO1, and RO6, where TP was low for all flows,~~
440 ~~usually ≤ 10 mg/m³. At the lake fed site of RO2, TP actually decreased with Q . Dissolved~~
441 ~~reactive phosphorus (DRP) generally increased with Q for most sites; however, there were many~~
442 ~~exceptions. Twenty sites had no detectable trend with Q ($r^2 < 0.10$). DRP actually decreased with~~
443 ~~Q at four sites (HM5, RO2, TU2, WA3).~~

444 _____

445 4.3.2-4.4.1. Catchment characteristics

446 Median monthly values of water quality variables for the 77 catchments ranged widely
447 (Table 5; Supplement Table 5). Some rivers had exceptional water quality all around, while
448 others had either current issues with multiple variables or worsening temporal trends (assessed
449 with SKSE from 1989 to 2014; Table 6). Because of the dependence of water quality on flow,
450 we first assessed temporal trends in Q . Only two catchments had significant increases in Q
451 (AX4, WH4), with the latter also being ‘meaningful.’ Three catchments had significant decreases
452 in Q (HM3, HM5, TU2) and five others also had ‘meaningful’ decreases in Q (CH2, GY4, HM4,
453 RO3, RO4).

454 Water temperatures (T_w) were not particularly high for any of the catchments; however,
455 21 rivers had significant increases in T_w , possibly the signature of climate change. The highest

456 rates of T_w increase ($0.04^\circ\text{C}/\text{y} < \text{SKSE} < 0.08^\circ\text{C}/\text{y}$) were for large alpine rivers in the central
457 South Island covered mostly by shrub/grasslands (TK3, TK4, TK6, AX3). Because of its strong
458 latitudinal trend (stronger than any land use effect), T_w was not analyzed further. Dissolved
459 oxygen (DO) was close to 100% for most catchments, but was particularly low (<90%) for two
460 catchments: RO2 which was affected by discharge from a large pulp mill at Kawerau, and AK2
461 which is on the Auckland fringe and thus affected by various peri-urban activities. DO was very
462 high (>110%) for one catchment (HV2) due to supersaturation from high periphyton in this
463 nutrient-enriched river. Temporal trends in DO from 1989 to 2014 were relatively minor
464 ($\text{RSKSE} < 0.5\%/y$), except RO2 which had a significant increase ($\text{RSKSE} = 0.7\%/y$) attributable
465 to progressive improvements in treatment of organic waste from its large pulp mill. Conductivity
466 ($COND$) was relatively low (<115 $\mu\text{S}/\text{cm}$) for all South Island catchments and varied
467 considerably for the North Island (54-528 $\mu\text{S}/\text{cm}$). Most catchments (52/77) experienced
468 significant or ‘meaningful’ increases in $COND$ from 1989 to 2014. Water pH (pH_w) was neutral
469 to alkaline for all rivers, which have been described as calcium-sodium bicarbonate waters by
470 Close and Davies-Colley (1990), and only displayed minor changes ($\text{RSKSE} < \pm 0.1\%/y$) over
471 the 26-year study period.

472 Median visual water clarity ($CLAR$) was exceptionally high (>5 m) for seven catchments
473 and very low (<1 m) for 22 catchments. Since 1989, $CLAR$ improved in almost half of the rivers,
474 and worsened in 4 rivers (Table 6; Supplement Table 5). $TURB$ was strongly inversely
475 proportional to $CLAR$ ($r_s = -0.97$) and generally followed opposite trends of $CLAR$. However,
476 fewer of its trends were significant and it had a disproportionately large number of ‘meaningful’
477 increases (17 catchments compared to only 2 ‘meaningful’ decreases in $CLAR$). $CDOM$ was low
478 for most of the rivers, with only five catchments greater than 2.0 m^{-1} . Nineteen of the catchments

479 experienced significant or ‘meaningful’ decreases in CDOM since 1989, possibly due to the loss
480 of wetlands across NZ. Only one catchment had a ‘meaningful’ increase in *CDOM* (TK3).

481 Total nitrogen (*TN*) was relatively high ($>250\text{--}455\text{ mg/m}^3$) for ~~more than~~ almost a
482 third of the catchments, with the vast majority (~~30~~17/~~39~~23) of these being lowland
483 catchments ($<150\text{ m}$ in elevation). Most of these catchments also had relatively high NO_x .
484 Thirty-three catchments had significant or ‘meaningful’ increases in *TN* from 1989 to 2014,
485 while only five had significant or ‘meaningful’ decreases in *TN* (Table 6). NO_x had a similar
486 number of increasing temporal trends, but also had ‘meaningful’ decreases for 12 catchments.

487 Total phosphorus (*TP*) followed a similar geographical pattern as *TN*. Eighteen of the 23
488 catchments with relatively high *TP* ($>30\text{ mg/m}^3$) were lowland catchments. Most of the
489 catchments with relatively high *TP* (18/23) also had relatively high *DRP* ($>9.5\text{ mg/m}^3$).
490 Seventeen catchments had ‘meaningful’ increases in *DRP*, compared to only three with
491 ‘meaningful’ decreases. There was more of a balance in temporal trends of *TP*, with eight
492 ‘meaningful’ increases and seven ‘meaningful’ decreases.

493 In addition to the expected correlations between *CLAR* and *TURB*, and among the
494 nitrogen and phosphorus constituents, several other significant relationships existed among the
495 water quality variables (Supplement Fig. 3). ~~*TP* was correlated with *CLAR* ($r_s = -0.77$), *TURB*~~
496 ~~(0.73), *TN* (0.71), *NO_x* (0.61), *CDOM* (0.62), and *COND* (0.65). *DRP* was also correlated with~~
497 ~~*TN* (0.71), *NO_x* (0.65), and *CDOM* (0.58). *CDOM* was correlated with *TN* (0.63). Finally, *COND*~~
498 ~~and *T_w* were correlated (0.67).~~ Taking into consideration this broad multicollinearity, we focus
499 our multivariate analyses on several key water quality variables, particularly those that
500 experienced the most changes from 1989 to 2014 (Table 6): *CLAR*, *TN*, NO_x , *TP*, and *DRP*.

501

502 4.4.5. Water Quality relationships with physiography, land use, and disturbance

503 ~~There was a predictable relationship between catchment area (A) and Q_{50} ($r_s = 0.89$; all~~
504 ~~following parentheses in this section are r_s , unless specified), and Visual water clarity ($CLAR$)~~
505 generally decreased with A (-0.37; all following parentheses in this section are r_s , unless
506 specified). Except for $TURB$ (0.32), no other water quality variables had significant relationships
507 with catchment area. Several water quality variables correlated with catchment slope (S_c),
508 including: TN (-0.72), TP (-0.63), and DRP (-0.65), meaning N and P concentrations were
509 relatively high in lowland (low slope) catchments. DRP (0.65) and TP (0.61) were directly
510 proportional to mean annual temperature (MAT), but this association probably arises because the
511 highest phosphorus values occurred mainly in lowland catchments and some of the northernmost
512 catchments, temperature being strongly correlated with altitude and latitude. DRP also had a
513 (counterintuitive) significant relationship with soil phosphate retention, P_{ret} (0.35). No other
514 strong physiographic relationships emerged from our analyses.

515 The strongest relationships between water quality and land use areal coverage (Table 7)
516 included high-producing grasslands (HG), which had strong positive relationships with several
517 water quality variables except $CLAR$ which decreased as HG increased. The lesser-managed
518 shrub/grasslands (SG) had generally opposite relationships with water quality, but note that SG
519 did not have significant relationships with $TURB$ or $CLAR$. Non-plantation forest (NF) followed
520 the same trends as SG , but had fewer significant relationships with water quality. Plantation
521 forest (PF), on the other hand, followed the same trends as HG , with poorer water quality being
522 associated with greater coverage of PF ; although correlations were not as strong as HG . $CDOM$,
523 DRP , and all N-constituents had significant negative correlations with open water (OW),

524 meaning that water quality improved with greater *OW* coverage, plausibly due to entrapment of
525 fine sediment and nutrients.

526 Water quality was significantly correlated with all stock unit density (SUD) metrics
527 (Table 7; Supplement Fig. 4), except deer (SUD_{de}) which only had relatively weak relationships
528 with *TN* and NO_x . The nutrients and CDOM had the strongest correlations with SUD_{cattle} , which
529 includes both dairy and beef cattle. *COND*, *CLAR*, and *TURB* had the strongest (slightly)
530 correlations with SUD_{be} . Overall, degraded water quality was strongly associated with high
531 livestock densities, even stronger than areal coverage of high-producing grasslands.

532 No significant correlations between water quality and total catchment disturbance (D_c)
533 were found; however, there were significant associations when disturbance was isolated by high-
534 producing grasslands (D_{HG}) and plantation forest (D_{PF} ; Table 7). Unexpectedly, *CLAR* and
535 *TURB* were not correlated to D_{HG} , and surprisingly, the rest of the water quality variables had a
536 significant *inverse* relationship with D_{HG} . Conversely, *CLAR* was the only water quality variable
537 correlated to plantation forest disturbance, D_{PF} ($r_s = -0.27$). Some interesting results emerged
538 when temporal trends in water quality (via SKSE) were assessed for catchments with high
539 disturbance. Of the 15 catchments with D_c greater than 5%, six had ‘meaningful’ increases in
540 *TURB* (RO3, HM4, RO6, WA6, HV6, HM2; all in North Island); while only one (HV5) had a
541 ‘meaningful’ decrease in *TURB*. Most of these 15 catchments also experienced significant
542 increases in *TN* (9 catchments; 7/9 also ‘meaningful’) and NO_x (10 catchments; 8/10 also
543 ‘meaningful’). Interestingly, *TP* and *DRP* significantly increased in only two of these highly
544 disturbed catchments.

545

546 4.5-6. Multivariate water quality relationships

547 In order to build on the above correlation analyses, the water quality variables of *CLAR*,
548 *TN*, *NO_x*, *TP*, and *DRP* were each assessed in a multivariate stepwise regression, using the
549 following ten physiographic and land use independent variables: *S_c*, *SC%*, *P_{ret}*, *Q₅₀*, *NF*, *SG*,
550 *HG*, *PF*, *OW*, and *SUD_{cattle}* (Table 8). The residual plots for all five water quality variables met
551 the assumptions of normality and linearity, but displayed heteroscedasticity with wide scatter for
552 high values. *CLAR* was correlated to *-HG*, followed by *+OW*, *-Q₅₀*, and *-PF*, where signs
553 represent whether the relationship is positive (+) or inverse (-). Thus, water clarity was
554 predictably lower for larger rivers that drain larger areas of high-producing grasslands and/or
555 plantation forests, but improved with increased open water coverage (Fig. 45).

556 The combined stock unit density for beef and dairy cattle (*SUD_{cattle}*) was the primary
557 predictor for all four nutrient variables, with *TN*, *TP*, and *DRP* also being proportional to
558 plantation forest coverage (*PF*; Table 8). Dissolved oxidized nitrogen (*NO_x*) was not
559 proportional to *PF*, or any other independent variable in the stepwise regression. Coverage of
560 high-producing grasslands (*HG*) and silt-clay surface soils (*SC%*) were also proportional factors
561 for *TN*. ~~In sum~~ Whether intensity or areal coverage, land use was the primary and secondary
562 predictor for all five water quality variables (Fig. 45).

563

564 5. Discussion

565 5.1. River water quality states and trends

566 We characterized water quality states and trends for 77 river sites across New Zealand
567 (NZ) using a wide range of flows and water quality conditions for each site, including some
568 small floods. We acknowledge that our analyses did not fully capture large floods due to their
569 short durations, unlikelihood of occurring during the preset monthly sampling, and the fact that

570 we relied on grab samples. These episodic floods are particularly important for the water quality
571 of downstream waters such as lakes and estuaries (Stamm et al., 2014). The uncertainty
572 surrounding our lack of flood samples could have been mitigated by composite samples or
573 supplemental flood samples; however, our 26 years of monthly samples for each site (n = 312)
574 did allow us to confidently report median conditions and temporal trends in water quality
575 (Moosmann et al., 2014).

576 ~~We found~~There was a wide range of water quality across NZ rivers (Table 5), with
577 drastic differences between upland and lowland rivers, distinguished by the 150 m elevation
578 threshold. For example, visual water clarity (*CLAR*), which is often used as a ‘master variable’
579 for overall water quality (Davies-Colley et al., 2003; Julian et al., 2008), was high for upland
580 rivers (mean = 3.2 m), with only two [alpine glacial flour-affected] rivers below the ANZECC
581 (2000) guideline of 0.6 m (CH3, AX3). Many of the upland rivers (7/33) had very high water
582 clarity (> 5 m), including one of the clearest non-lake-fed rivers in the world – Motueka River
583 (NN2) with a median *CLAR* of 9.8 m. The lowland rivers, in contrast, had a mean *CLAR* of 1.2
584 m, with 17 (39%) below the ANZECC guideline of 0.8 m. Note that these ANZECC (2000)
585 guidelines, which are statistical derivations (i.e. 20th-percentile of the first decade of the
586 NRWQN record for ‘reference’ sites), are merely ‘trigger values’ that when exceeded trigger a
587 management response to protect ecosystem health (Hart et al., 1999). Although these ‘trigger
588 values’ are not effects-based standards (which would be difficult to define for the wide variety of
589 NZ ecosystems), they do provide a useful reference for comparing water quality states and
590 trends. Save for a few borderline exceptions, the same sites that were below visual clarity
591 guidelines also exceeded the turbidity trigger values of 4.1 and 5.6 NTU for upland and lowland
592 rivers, respectively.

593 ~~Nutrient concentrations in NZ rivers also varied widely (Table 5), again with high~~
594 ~~concentrations typically in lowland catchments and low concentrations in upland catchments.~~
595 Nine of the ten catchments with the highest TN ($>740 \text{ mg/m}^3$) were lowland catchments. In all,
596 13 lowland catchments exceeded the ANZECC TN guideline of 614 mg/m^3 and 8 upland
597 catchments exceeded the guideline of 295 mg/m^3 . Almost three quarters of these catchments
598 (15/21) also exceeded the NO_x guideline of 444 mg/m^3 (lowland) and 167 mg/m^3 (upland). There
599 were a similar number of sites exceeding ANZECC guidelines for TP ($33/26 \text{ mg/m}^3$ for
600 lowland/upland) and DRP ($10/9 \text{ mg/m}^3$ for lowland/upland), each with at least 20 and most of
601 these were corresponding. Our results on the state and trends of the 77 NRWQN catchments
602 generally accord with earlier NRWQN studies (e.g. Ballantine and Davies-Colley, 2014) and a
603 recent publication by Larned et al. (2016), which analyzed water quality states and trends for 461
604 NZ river sites for the period 2004-2013.

605 Based on ANZECC (2000) trigger values, we have organized the catchments into four
606 classes (Fig. 56): I. clean river with high visual water clarity (*CLAR*) and low dissolved inorganic
607 nutrients (DIN); II. sediment-impacted river with low *CLAR* and low DIN; III. nutrient-impacted
608 river with high *CLAR* and high DIN; and IV. sediment- and nutrient-impacted river with low
609 *CLAR* and high DIN. Note that the term ‘sediment-impacted’ is a connotation for total suspended
610 solids (TSS), which includes organic matter as well. In agriculture-dominated catchments, both
611 mineral sediment and particulate organic matter can greatly increase TSS (Julian et al., 2008).
612 We use *CLAR* as a preferred metric for suspended matter because TSS is not routinely measured
613 in the NRWQN (or other monitoring networks) while *CLAR* correlates strongly to TSS ($r = -$
614 0.92), and better than *TURB* ($r = 0.87$) (Ballantine et al., 2014). Further, *CDOM* in NZ rivers is
615 low with minimal impact on *CLAR*. We use NO_x as our preferred metric for DIN because it is

616 least affected by suspended sediment and soil properties (compared to *DRP*). However,
617 catchments that exceed ANZECC guidelines for *DRP* are indicated in Fig. 5-6 by grey-filled
618 markers.

619 When this classification is combined with the SKSE trend analyses (Table 6), we obtain a
620 clear picture of the current and potential state of NZ rivers (Fig. 56). Before individual rivers are
621 discussed ~~(next section)~~, we first point out key differences between the upland and lowland
622 catchments, which will later be placed within the context of physiography and land use intensity.
623 Most obvious, and consistent with the findings of Larned et al. (2004), was that lowland rivers
624 were much more degraded, particularly by sediment. More than a third of the lowland
625 catchments were either Class II or IV (17/44); whereas, only two upland catchments were Class
626 II. None of the upland catchments were Class IV, and more than two-thirds were clean rivers
627 (Class I). Both types had a similar number of nutrient-impacted rivers (Class III). ~~Another major
628 difference is that all but three of the upland catchments are far from class boundaries, meaning
629 that they are relatively stable in terms of water quality. Further, almost all of the upland
630 catchments that have had significant increases in NO_x were already nutrient impacted.
631 Conversely, many of the lowland catchments are very close to class boundaries, with most of
632 these having recently changed classes or likely crossing over in the near future.~~ Particularly
633 concerning is that almost half of the lowland rivers (19/44) are currently experiencing
634 ‘meaningful’ increases (>1% per year) in NO_x , *DRP*, or both. The other striking trend is that
635 many of the lowland rivers are becoming clearer, with 18/44 experiencing ‘meaningful’
636 increases (>1% per year) in *CLAR* – which, plausibly, has been attributed to increasing riparian
637 fencing to exclude cattle from channels (Davies-Colley, 2013; Ballantine and Davies-Colley,
638 2014; Larned et al., 2016).

639 While clearer rivers are seen as an improvement in water quality; when combined with
640 increasing nutrients, warmer water, and lower flows, the perfect recipe for toxic algae blooms is
641 created ([Dodds and Welch, 2000](#); [Hilton et al., 2006](#)). Only recently has the widespread problem
642 of toxic algae blooms in NZ rivers been evidenced (Wood et al., 2015; McAllister et al., 2016),
643 and our results indicate that this problem could worsen given the increasing trends we found in
644 water temperatures, ~~DIN~~[inorganic nutrients](#), and most influential in our opinion, water clarity.
645 ~~Eutrophication~~[Nutrient enrichment](#) and global warming receive the most attention when it
646 comes to degraded water quality, but rivers have increasingly become light-limited (Hilton et al.,
647 2006; Julian et al., 2013) such that when clarity improves in warm, nutrient-rich rivers, algae can
648 proliferate. Particularly problematic for NZ is that its lowland catchments, which are warmer
649 (mean median T_w of 13.6 v 10.8 °C for upland rivers), have much greater ~~DIN~~[DRP and \$NO_x\$](#) , and
650 have longer water residence times, are the ones becoming appreciably clearer (Fig. 56). If
651 droughts become more frequent and intense in NZ, toxic algae blooms are also likely to become
652 more frequent, more widespread, and more problematic. However, this algae response is
653 complex and depends on a number of interacting factors such that the apparent potential for
654 increasing algal nuisance might not necessarily be realized in some rivers ([Dodds and Welch,](#)
655 [2000](#); [Hilton et al., 2006](#)).

656

657 5.2. The role of physiography in dictating land use [intensity](#) across NZ

658 While physiography did not emerge as a significant independent variable in the
659 multivariate analyses (except TN with $SC\%$), physiography is important because it largely
660 controls the location and intensity of agricultural land uses. The greatest coverages of high-
661 producing grasslands (HG) and the highest densities of cattle (SUD_{cattle}), the two primary

662 explanatory variables for all five major water quality variables (Table 8), were both found
663 predominantly in flat areas with deep soils located in warm, sunny, and relatively dry climates.
664 Livestock in NZ depend almost exclusively on pasture grasses and thus their productivity is
665 maximized when pasture productivity is maximized. The very large cattle are not well suited for
666 steep slopes, particularly dairy cattle which can weigh more than 500 kg. Deep soils are
667 important because they absorb and hold more water for plant uptake, and are not as susceptible
668 to waterlogging, especially in wetter climates. Year-round and intense grazing is best supported
669 by warm and sunny climates where pasture grasses are highly productive and recover quickly
670 following intense grazing such as strip/rotational grazing which is common in NZ dairy farms.

671 Another soil property we found to be positively correlated to SUD_{cattle} was phosphate
672 retention (P_{ret}). The highest dairy cow densities were found on Allophanic volcanic soils with
673 high P_{ret} , likely because these soils respond favorably to P-fertilizer and thus can be managed
674 more intensively. However, soils with high P_{ret} require more P-fertilizer, and thus generally have
675 higher export of DRP to rivers. Our finding of a significant positive correlation between these
676 two variables is consistent with this interpretation. Further, we found that high-producing
677 pastures with high P_{ret} had the lowest disturbance (D_{HG}), indicating that these intensively
678 managed pastures recover quickly following grazing. In a more comprehensive study of land
679 disturbance across the North Island of NZ, de Beurs et al. (2016) also found that Allophanic soils
680 had the least disturbance among all soil orders. Where high livestock densities occur in less than
681 ideal conditions, land disturbance is likely. Our catchment-scale analyses limit our interpretation
682 of specific situations, but based on our results, field observations and previous remote sensing
683 analyses, pasture disturbance in NZ will likely be highest during droughts on steep, south-facing

684 slopes with thin soils being heavily grazed by sheep. Under these conditions, grasses will be
685 grazed down to bare soil and recover very slowly.

686 Plantation forests (*PF*) in NZ also correlated with thick soils with relatively high P_{ret} on
687 flat areas, particularly the pumice soils of the central North Island. The porous nature of the
688 pumice soils allows them to efficiently hold and regulate nutrients, water, and air; while being
689 well-draining and resistant to compaction and flooding. Under these conditions, radiata pine (the
690 dominant *PF* species in NZ) grows rapidly (mean harvest cycle of 28 y) and can be harvested
691 year-round. Since 1990 however, many of the *PF* additions have occurred on steeper slopes in
692 response to carbon credit incentives, greater economic demand for wood products (PCE, 2013),
693 and the need for soil erosion control on steep pasture susceptible to land-sliding (Parkyn et al.,
694 2006).

695

696 5.3 Land use intensity and water quality in New Zealand rivers

697 ~~5.3.1 Land use diversity and effectiveness~~

698 ~~Water quality in NZ rivers has been related to regional differences in climate and source-~~
699 ~~of flow (Larned et al., 2016); however, we focus here on the role of land use because (1) the vast~~
700 ~~majority of our catchments were large (only five less than 100 km²) and thus their surface water~~
701 ~~quality was likely dominated by catchment characteristics (Julian and Gardner, 2014); (2) the~~
702 ~~changes we observed in water quality have been linked to land use globally (Foley et al., 2005;~~
703 ~~Vitousek et al., 1997a; Bennett et al., 2001; Walling, 2006); and (3) our results indicate that land~~
704 ~~use was the dominant source of diffuse pollutants, and thus influence on spatial and temporal~~
705 ~~patterns in river water quality across NZ. Before describing relationships, we would like to first~~
706 ~~point out that the 77 NRWQN catchments captured the diversity of land use in NZ, with *NF*, *SG*,~~

707 ~~and *HG* (the three dominant land uses of NZ) accounting for 84% of both the 77 catchments and~~
708 ~~NZ as a whole. Our empirical study was also an excellent natural experiment in which to assess~~
709 ~~the effects of land use on water quality because we had an assortment of dominant land uses~~
710 ~~(\rightarrow 50% area) among our catchments (Fig. 1): 24 *HG*, 13 *SG*, 13 *NF*, 2 *PF*, and 25 mixed (i.e. no~~
711 ~~single dominant land use).~~

712

713 5.3.2-1 High-producing pastures and livestock densities

714 High-producing grassland coverage (*HG*) was the primary explanatory variable for visual
715 clarity (*CLAR*; Table 8, Fig. 45). *CLAR* in NZ rivers is mostly influenced by mineral and organic
716 particulates (Davies-Colley et al., 2014). Livestock reduce visual clarity in multiple ways,
717 especially in NZ where high densities of multiple types of livestock tread year-round on
718 relatively steep slopes with highly erodible soils vegetated by shallow-root introduced grasses
719 which are susceptible to destabilization (McDowell et al., 2008). The year-round treading is
720 particularly important because most NZ regions during winter are very wet with short days,
721 which increases soil disturbance (pugging and compaction) and slows recovery times. Where
722 livestock have direct access to rivers, their trampling of riverbanks and instream disturbance is
723 often the main contributor to reduced *CLAR* (Trimble and Mendel, 1995; McDowell et al., 2008).

724 The lowland flatter areas in NZ have high *HG* coverage and high cattle stock densities
725 (SUD_{cattle}). These lowlands also have high drainage densities – often increased by artificial
726 drainage. The influence of *HG* on *CLAR* is thus exacerbated by this interaction of high SUD_{cattle}
727 and artificial drainage, ~~which explains the high negative correlation between *HG* and *CLAR* ($-$~~
728 ~~0.45). Interestingly, SUD_{cattle} was not an explanatory variable for *CLAR* in the stepwise~~
729 regression, which is likely a result of two factors. First, *HG* and SUD_{cattle} are highly correlated,

730 and stepwise regression does not include secondary variables that are explaining the same
731 proportion of variance as the primary independent variable. Second, we found that *CLAR* has
732 actually *improved* in catchments where SUD_{cattle} is high and/or has increased (Fig. 56), which we
733 ~~noted earlier could be a result of increased attribute to the promotion of~~ riparian fencing, ~~across~~
734 ~~NZ since~~In 2003, ~~NZ implemented when~~ the *Dairying and Clean Streams Accord*, ~~which has led~~
735 ~~to the exclusion of dairy cattle from 87% (as of 2012) of perennial rivers greater than 1 m in~~
736 ~~width was implemented~~ (Bewsell et al., 2007; Howard-Williams et al., 2010; Gunn and
737 Rutherford, 2013). By excluding (dairy) cattle from channels and riparian zones, the contribution
738 of riverbank and bed erosion to degraded *CLAR* has likely been mitigated and reduced over time
739 (Trimble and Mendel, 1995; Hughes and Quinn, 2014). Indeed, *CLAR* has been significantly and
740 meaningfully improving in many of NZ's rivers (Table 6), even those with increasing SUD_{cattle} ,
741 albeit from a fairly degraded condition. Of the 34 catchments with significant increases in *CLAR*,
742 all but 5 had increases in SUD_{cattle} from 1990 to 2012.

743 Another potential explanation for improved water clarity at numerous sites is the
744 considerable decrease in sheep density across the NZ landscape. NZ had 57.65 million sheep in
745 1990. By 2012, that number had been reduced by almost half, to 31.19 million (StatsNZ, 2015).
746 Although cattle are larger and have a greater treading impact per animal, the much greater
747 number of sheep means that stock unit density (SUD) may be broadly comparable as regards
748 environmental impact. Another difference is that sheep are generally placed on steeper, less
749 stable slopes in NZ, where headwater stream channels are located. Where there are breaks in
750 slope (even small ones), sheep create tracks of bare soil with their hooves and hillside scars with
751 their bodies (for scratching and shelter), both of which can enhance soil erosion (Evans, 1997).
752 Further, cattle (using their tongues) leave approximately half the grass height on the pasture after

753 grazing; whereas sheep (using their teeth) graze approximately 80% of grass height (down to
754 bare soil in dire conditions), leaving it exposed to erosion (Woodward, 1998). Considering all
755 these factors, sheep can have a greater impact on sediment runoff into rivers, and consequently
756 visual clarity, than suggested by their aversion to water *versus* cattle's attraction to water.
757 Although not isolated in our analyses, the particulate fractions of *TN* and *TP* have likely been
758 affected by similar processes as *CLAR* and may follow the same temporal trends (Ballantine and
759 Davies-Colley, 2014).

760 While *HG* was also strongly correlated to river nutrient concentrations (Table 7), the
761 primary explanatory variable for all four major nutrient metrics (Table 8, Fig. 45) was land use
762 intensity as measured by the livestock density of beef and dairy cattle (*SUD_{cattle}*). The difference
763 between these two explanatory variables may seem trivial, however the distinction is important if
764 we want to understand future trends and effectiveness of water quality management strategies.
765 As we demonstrated, the area of land used for high-producing grasslands (*HG*) has not changed
766 much since 1990 (Fig. 2). In fact, it has decreased or stayed virtually the same in all but two of
767 the 77 catchments. Yet, nutrient concentrations have been increasing in many of the rivers (Table
768 6), which we attribute to (1) increasing numbers of cattle (mostly dairy) on both *HG* and *SG*, and
769 (2) legacy nutrients being slowly delivered to the rivers in groundwater. From 1990 to 2012, NZ
770 approximately doubled its number of dairy cattle, exceeding 6.4 million. (StatsNZ, 2015). This
771 enormous addition to a country that is only 268,000 km² in area, has been accompanied by more
772 than 1.426 million tonnes of P-based fertilizers and 335,000 tonnes of N-based fertilizers
773 annually (1990-2012 mean; StatsNZ, 2015). Of the nutrients consumed by lactating dairy cows,
774 approximately 79% of N and 66% of P are returned to the landscape in the form of urine and
775 feces (Monaghan et al., 2007). This results, potentially, in about 260,000 tonnes of N-based and

776 940,000 tonnes of P-based diffuse pollution. Some of these nutrients will be transported to rivers
777 during subsequent storms, but a majority will remain (building up) in the landscape to be slowly
778 added to rivers over decadal time-scales (Howard-Williams et al., 2010).

779

780 5.3.3-2. Plantation forests

781 All water quality variables were significantly correlated to plantation forest coverage
782 (*PF*; Table 7), with a negative relationship with *CLAR* (~~i.e. *CLAR* was lower for higher *PF*~~) but
783 positive for all other variables (~~i.e. nutrients increased with *PF*~~). From the stepwise regression,
784 *PF* emerged as an explanatory variable for all major water quality variables except *NO_x* (Table
785 8), suggesting that its dominant impact on river water quality was from surface runoff. Plantation
786 forestry activities can add a considerable amount of sediment and nutrient pollution to rivers,
787 especially during and immediately following harvesting (Fahey et al., 2003; Croke and Hairsine,
788 2006; Davis, 2005). This harvesting period of maximum soil disturbance usually lasts about two
789 years (Fahey et al., 2003), but the land cover may remain sparsely vegetated and susceptible to
790 erosion for several years (but usually not more than 5 y; de Beurs et al., 2016). The greatest *PF*
791 impact on sediment runoff, and thus potentially *CLAR*, is usually from road sidecast/runoff,
792 shallow landslides, and channel scouring/gullyng (Fahey et al., 2003; Motha et al., 2003;
793 Fransen et al., 2001).

794 Rivers receive a pulse of nutrients during the forest harvest, but fertilizers are also
795 applied at time of re-planting and sometimes routinely to enhance growth (Davis, 2005). Radiata
796 pine in the pumice soils of the central North Island, the dominant area of *PF* in NZ, are
797 particularly responsive to both N- and P-fertilizers and thus likely receive ample supplements.
798 Like pasture fertilizers, some of these nutrients may be delivered to rivers during intense

799 precipitation, but there is also a legacy of nutrients left behind. Fertilizers have been applied to
800 plantation forests in NZ since the 1950s, with an intense period of application in the 1970s
801 (Davis, 2005). While fertilization rates (tonnes/ha/y) have decreased since 1980, the amount of
802 NO_x leaving catchments mostly covered in *PF* has significantly and ‘meaningfully’ increased
803 since 1989: ~~RO3 (69.8% *PF*, 3.0%/y RSKSE), RO5 (53.3% *PF*, 1.7%/y RSKSE), and RO2~~
804 ~~(42.5% *PF*, 1.2%/y RSKSE)~~. None of these catchments had more than 17.7% *HG*, none had
805 major increases in *HG* ($< 0.3\%$), none had major increases in SUD_{cattle} (< 0.7 SU/ha), and none
806 had a significant increase in D_{PF} . What the catchments did have in common were all had
807 gravelly/sandy pumice soils (< 4.5 *SC%*) and all were intensively managed as reported by Davis
808 (2005) and as indicated by high D_C ($> 6.8\%$). The extended periods of nonvegetated land due to
809 weed control also increases the amount of nutrients delivered to rivers over the long term (Davis,
810 2005).

811

812 ~~5.3.4 Other land uses~~

813 ~~Open water (*OW*) in the form of lakes can remove sediment, nutrients, and *CDOM* by a~~
814 ~~range of processes (Schallenberg et al., 2013; Wetzel, 2001). Consistent with this concept, our~~
815 ~~bivariate comparisons showed that catchments with more *OW* had lower *CDOM*, *TN*, NO_x , and~~
816 ~~*DRP* (Table 7). Our multivariate analyses found *OW* to be an explanatory variable for *CLAR*~~
817 ~~(Table 8, Fig. 4), which we attribute to several of the stations with high *CLAR* being located~~
818 ~~downstream of large lakes (AX1, DN10, RO1, RO6). If these 4 catchments are removed, the~~
819 ~~relationship between *OW* and *CLAR* is not significant. While lakes can improve downstream~~
820 ~~water quality, many lakes in NZ, particularly shallow lakes, are experiencing eutrophication and~~

821 other water quality issues (Larned et al., 2016; Abell et al., 2011), which can cause regime shifts
822 (Schallenburg and Sorrell, 2009) and degrade downstream river water quality.

823 An important land use for nutrient/sediment fluxes that was missing from our analyses
824 was vegetated wetlands (VW), which was a consequence of exceptionally low VW coverage in
825 NRWQN catchments (0.1% on average and a maximum of 2.2%). With such a miniscule
826 coverage, these residual wetlands do not provide a detectable water quality improvement
827 function at the catchment scale (Mitsch and Gosselink, 2000). Historically, wetlands covered
828 approximately 10% of mainland NZ (Ausseil et al., 2011). This considerable loss (> 90% of pre-
829 European extent) of wetlands has deprived NZ rivers of many valuable ecosystem services,
830 especially the filtration/processing of sediment and nutrients (Clarkson et al., 2013; Verhoeven et
831 al., 2006). If some of these wetlands could be restored, some of the alarming eutrophication
832 trends we have documented here (Table 6, Fig. 5) could be mitigated. For example, Mitsch et al.
833 (2001) found that just adding 10% of wetland coverage can reduce up to 40% of the nitrogen
834 entering receiving waters.

835 The other important land use missing from our analyses was urban (UR), also because
836 very little of NZ's land area is urban (Table 3), accounting on average for only 0.35% of our
837 catchment areas (maximum 5.8%). However, urban water management did have major effects on
838 three of our catchments by reducing *DRP* point sources. The 'meaningful' decrease of *DRP*
839 ($RSKSE = -4.6\%/y$) in the Manawatu River below Palmerston North (WA9) was due to
840 progressive improvements in the city's wastewater treatment, particularly after 2008 when a new
841 main wastewater treatment plant (incorporating P-removal) became fully operational. *DRP* was
842 also 'meaningfully' reduced ($RSKSE = -5.3\%/y$) for the Ohinemuri River below Waihi (HM6)
843 when P-removal was added to the Waihi wastewater treatment plant in 2005. And *DRP* for Hutt

844 ~~River at Bouleott (WN1) was ‘meaningfully’ reduced (RSKSE = -3.1%/y) with progressive~~
845 ~~improvements to the Hutt Valley wastewater treatment, which were completed in 2002. It is~~
846 ~~important to note that these point discharge-affected sites were the only ones with meaningful~~
847 ~~reductions in *DRP*.~~

848

849 5.3.5.3. Land disturbance and water quality

850 So far, we have discussed how land use affects water quality, with a focus on sediment
851 and nutrient runoff from high-producing grasslands (*HG*) and plantation forests (*PF*). When land
852 is disturbed (i.e. bare soil), sediment/nutrient mobilization can be enhanced. The most intense
853 and longest lasting disturbances occurred during plantation forest harvests. Following harvest,
854 we found that the land remained disturbed for 1-6 years, with a mean of 1.5 years. The overall
855 mean and median D_{PF} among all catchments was 10%, which means that plantation forestry
856 leaves large areas of disturbed land at any one time. When this bare land is exposed to intense
857 precipitation, large quantities of sediment and nutrients can be mobilized into the rivers. This
858 happened in the Motueka Catchment (NN1) in 2005 when a 50-y storm fell on some recently-
859 harvested plantation forests. For one of NN1’s sub-catchments, the post-harvest disturbed land
860 caused a five-fold increase in sediment yield compared to pre-harvest events. Following this
861 event, sediment yields at NN1 were elevated by a factor of 2-3 over the next 3 years (Basher et
862 al., 2011). Similar sediment erosion events for plantation forests during the post-harvest
863 disturbance have been documented for other catchments across NZ (Hicks et al., 2000; Phillips et
864 al., 2005). Because these disturbances only last a few years, they typically do not show up as
865 temporal trends (via SKSE); however it is possible that they produce enough readily available
866 sediment to impact water quality for longer periods ([Kamarinas et al., 2016](#)).

867 The coincidence of rainstorms on disturbed pasture could have the same effect on
868 sediment/nutrient runoff if the pasture is connected to the stream network via steep slopes or
869 adjacent channels/canals (Dymond et al., 2010; [Kamarinas et al., 2016](#)). Pastures become
870 disturbed from overgrazing, strip grazing, pugging/soil compaction, tilling/reseeding,
871 cropping/harvesting, or landsliding on steep slopes. Given the high intensity of grazing
872 management in NZ, all of these are common. While D_{HG} was lower than D_{PF} on average, D_{HG}
873 had a higher maximum (Table 4). Spatiotemporal patterns in disturbance between these two land
874 uses were also different (de Beurs et al., 2016). D_{PF} covered large areas and lasted years at a
875 time; whereas D_{HG} had two patterns: (1) one related to dairy cattle strip grazing, which were
876 short-lived due to quick recovery times of grasses in fertilized soils; and (2) more widespread
877 and longer continuous disturbances occurring on steeper slopes grazed by sheep and beef cattle,
878 particularly following drought periods. Because our disturbance analyses had a spatial resolution
879 of 463 m, we likely missed some paddock-scale disturbances. Future work could use Landsat
880 imagery (30-m resolution) to assess disturbance (*sensu* de Beurs et al., 2016).

881 All six catchments with ‘meaningful’ increases in D_{HG} had large increases in dairy cattle
882 density 1990-2012 (mean of ± 1.0 SU/ha across the catchment). Not surprisingly, all six
883 catchments suffered impacts to water quality. Five of the six had ‘meaningful’ increases in DRP
884 and three had meaningful increases in NO_x and TN . One had a ‘meaningful’ increase in $TURB$
885 and three had significant reductions in DO . One of these catchments, in particular, may provide a
886 glimpse into NZ’s future if agricultural intensification continues. The Waingongoro River
887 catchment (WA3) is covered almost entirely by HG (91.2%), with practically all of this land
888 being used for intensive strip grazing. The SUD_{da} was 15.0 SU/ha in 1990 and increased to 15.4
889 SU/ha by 2012. The D_{HG} from 2000-2013 had a strong increasing trend of 9.8%/y RSKSE,

890 associated with the intensification of dairy operations (Wilcock et al., 2009). The result of all this
891 intensification was that WA3 had ‘meaningful’ increases in *TP*, *DRP*, and *TN*. The only reason
892 *NO_x* did not display a significant trend is because the catchment was already overloaded with a
893 median river concentration of 1,852 mg/m³. Noteworthy is that these significant trends of
894 increasing *SUD_{da}*, *D_{HG}*, and nutrients are occurring not only in lowland catchments on the North
895 Island (WA3, HV2), but also in upland catchments of the North Island (RO6), as well as both
896 lowland (TK1) and upland (CH3, TK2) catchments on the South Island.

897 While disturbance was not itself a strong predictor of water quality, it did help explain
898 outliers of land use-water quality relationships. For example, streams with high *DRP* (> 20
899 mg/m³; 10th percentile) had one of two dominant land uses, either plantation forest, *PF* (RO2,
900 RO3) or high-producing grassland, *HG* (HM5, WA3, WA9, HM4, HM2). The one exception was
901 RO4, which had relatively low coverage of *PF* (11.2%) and *HG* (2.9%). In fact, RO4 is
902 dominated by *NF* (79.1%). Upon closer examination, we found that the small areas of *PF* and
903 *HG* in RO4 were disturbed frequently. Further, most of the disturbed forestry occurred on steep
904 slopes and most of the disturbed pastures (practically all sheep and beef) occurred on hilly terrain
905 adjacent to stream channels. Our high temporal-resolution analyses of disturbance showed that
906 even though this catchment is mostly indigenous forest, intense disturbances on small
907 proportions of developed land can have a considerable impact on water quality. RO4 is also
908 experiencing significant increases in *TURB* and *TP*, as well as a significant decrease in *Q*.
909 Another outlier example was RO3, which was the only non-*HG*-dominated catchment with
910 ~~extremely~~ high *NO_x* (634 mg/m³). RO3 was dominated by *PF* (69.8%), but it had the highest
911 median disturbance (10.5%) of all catchments. ~~As discussed previously, disturbance in plantation~~
912 ~~forests is correlated with harvest frequency and management intensity. In addition to the many~~

913 ~~pulses of NO_x from the forest harvests and post-harvest storms over a vegetation-cleared soil~~
914 ~~surface, all of the replantings in the N-deficient pumice soils would have been accompanied by~~
915 ~~routine N-fertilizer applications (Davis, 2005). And the catchment's well-drained sandy/gravelly~~
916 ~~soils meant that this dissolved N was transported to streams without much attenuation.~~ This
917 catchment also exceeds ANZECC guidelines for *DRP* and has experienced meaningful increases
918 in *TURB*, *TN*, and *NO_x*.

919 We believe that land disturbance and consequently river eutrophication and reduced
920 visual clarity will continue to worsen in some NZ catchments based on the following. More
921 plantation forests were planted 1993-1997 (3,810 km²) than any other 5-y period in NZ history
922 (NZFFA, 2014). With a 28-y mean age of harvest, NZ will experience its greatest coverage and
923 intensity of forest disturbance around 2025, less than 10 years from now. When combined with
924 drought and intense storms, the potential for nutrient and sediment mobilization from these lands
925 into NZ's rivers is high, especially given that approximately 45% of these plantings occurred on
926 high-producing grasslands (NZFFA, 2014) where many of the legacy nutrients will be exported
927 to rivers during forest harvest (Davis, 2014). Many of these plantings also occurred on steep
928 slopes, which exacerbates sediment runoff. If carbon prices continue to stay low, there will be a
929 high likelihood that many of the harvested forests will be converted to pasture, adding even more
930 nutrients to NZ rivers (PCE, 2013). Given that the Central Government created a national policy
931 goal of nearly doubling the export to GDP ratio by the year 2025 (MBIE, 2015), NZ is likely to
932 see continued increases in livestock density, fertilizer usage, and supplemental feed to support
933 these extra livestock, all of which will add even more pressure and risks of eutrophication on
934 NZ's rivers.

935

936 Conclusions

937 This study had the overall goal of describing how changes in land use ~~and land~~
938 ~~disturbance~~intensity impact river water quality across broad scales and over long periods. To
939 address this goal we used a combination of ‘brute force’ statistical analyses (in terms of hundreds
940 of analyses using a suite of physiographic, land use, and water quality data for 77 catchments
941 over 26 years) and careful examination (using multi-resolution data to find patterns and
942 relationships among these variables). This goal was ambitious and we likely missed some
943 relationships and details of water quality changes. However, we found empirical evidence for
944 several key relationships among land use intensity, ~~land disturbance~~geomorphic processes, and
945 water quality, which we now place into a broader perspective.

946 The greatest negative impact on river water quality in New Zealand (NZ) in recent
947 decades has been high-producing pastures that require large amounts of fertilizer to support high
948 densities of livestock. While this claim has been previously published (Davies-Colley, 2013;
949 Howard-Williams et al., 2010; and references within), our results and supporting information
950 show that the relationship between high-producing pastures and water quality is complicated,
951 being dependent on ~~physiography (particularly soil type)~~, livestock type/density, ~~and~~ disturbance
952 regime, and physiography, particularly soil type. Dairy cattle receive much of the blame for
953 degraded water quality because of their high nutrient requirements (Howard-Williams et al.,
954 2010), but beef cattle can also strongly degrade water quality due to comparable required inputs
955 and grazing on steeper land with a higher potential for runoff (McDowell et al., 2008). Further,
956 pasture designations/boundaries are becoming increasingly blurred by modern cattle
957 management, with greater movements of dairy and beef cattle among pastures, greater use of
958 high-producing pastures for beef, over-wintering of dairy cattle on beef pastures, and cross-

959 breeding (Morris, 2013). While riparian fencing has ~~no doubt~~plausibly improved the clarity of
960 NZ rivers, the removal of millions of sheep from steep slopes has also likely played a role that
961 should be investigated further.

962 New Zealand is the global leading exporter of whole milk powder, butter, and sheep
963 products; and NZ's prominence in these industries is likely to continue over the next decade
964 (OECD/FAO, 2015). In their~~this~~ most recent environmental review by the Organisation for
965 Economic Co-operation and Development (2015), NZ had the highest percent increase (1990-
966 2005) in agricultural production out of 29 OECD countries, the highest percent increase in N-
967 fertilizer use, and the 2nd highest increase in P-fertilizer use. This agricultural intensification
968 massive application of nutrients to the NZ landscape over our study period is reflected in overall
969 nutrient enrichment of NZ rivers (Fig. 5; Table 6). If cattle continue to be added at the rates we
970 documented, additional fertilizers and supplemental feed will be needed. Even if best
971 management practices are adopted to reduce nutrient export to rivers, there is already a half-
972 century legacy of nutrients distributed across the NZ landscape that will continue to leak to the
973 rivers (Larned et al., 2016). Indeed, the full impact of agricultural intensification on river water
974 quality will not be fully appreciated for another several decades (Howard-Williams et al., 2010;
975 Vant and Smith, 2004). Having an extensive national network like the NRWQN to document and
976 study these water quality changes will be important.

977 However due to legacy/lag effects, notably the slow delivery of nutrients to rivers from
978 land and groundwaters (Larned et al., 2016), the full impact on river water quality will not be
979 fully appreciated for another several decades (Howard-Williams et al., 2010; Vant and Smith,
980 2004). ~~Because NZ's economy is heavily dependent on agricultural production, the agricultural~~
981 ~~intensification that we have documented since 1990 may be expected to continue, with greater~~

982 livestock densities being supported by supplemental feed and fertilizers. Even if best
983 management practices are adopted to reduce nutrient export to rivers, there is already a half-
984 century legacy of nutrients distributed across the NZ landscape that will continue to leak to the
985 rivers. Having an extensive national network like the NRWQN to document and study these
986 water quality changes is important, but unfortunately the NRWQN is being down-sized at the
987 time of writing. Less than half of the 77 sites are to be retained by NIWA in a 'benchmark'
988 network, with 'excess' sites being transferred to regional operation or closed. Although regional
989 management agencies in NZ conduct much water quality monitoring (e.g. Larned et al., 2016),
990 the quality (of some) and consistency of their datasets falls short of the NRWQN—which was
991 also longer running than all but a very few regional sites.

992 In response to public concerns on water quality, New Zealand released its National Policy
993 Statement on Freshwater Management in 2011. Data and evidence based science is now needed
994 to support and facilitate limit settings for water quality standards, especially for diffuse pollution
995 (Duncan, 2014). ~~In their most recent environmental review by the Organisation for Economic
996 Co-operation and Development (2015), NZ had the highest percent increase (1990-2005) in
997 agricultural production out of 29 OECD countries, the highest percent increase in N fertilizer
998 use, and the 2nd highest increase in P fertilizer use. This massive application of nutrients to the
999 NZ landscape over our study period is reflected in overall nutrient enrichment of NZ rivers (Fig.
1000 5; Table 6). However due to legacy/lag effects, notably the slow delivery of nutrients to rivers
1001 from land and groundwaters (Larned et al., 2016), the full impact on river water quality will not
1002 be fully appreciated for another several decades (Howard-Williams et al., 2010; Vant and Smith,
1003 2004).~~

1004

1005 Author contribution

1006 J. Julian designed the study and performed most of the analyses. K. de Beurs developed the
1007 disturbance dataset and performed all trend analyses, both with assistance from B. Owsley. R.
1008 Davies-Colley provided water quality dataset and guidance on its use. A.-G. Ausseil developed
1009 the stock unit density dataset and provided guidance on land use analyses. J. Julian prepared the
1010 manuscript with contributions from all co-authors.

1011

1012 Acknowledgments

1013 The ~~impetus~~inspiration for this research was J.P. Julian's Fulbright Senior Scholar
1014 Fellowship, which was hosted by the National Institute of Water & Atmospheric Research
1015 (NIWA) in Hamilton, NZ in 2012. This work was funded by NASA LCLUC grant
1016 NNX14AB77G and NSF Geography grants #1359970 and #1359948 (Co-PIs Julian and de
1017 Beurs). Andrew Tait (Climate Principal Scientist at NIWA) provided climate data. Agricultural
1018 production and other data essential to this manuscript was collected by William Wright
1019 (Landcare Research; LCR). Many other people in New Zealand provided expert advice,
1020 including Suzie Greenhalgh (LCR), Sandy Elliot (NIWA), Andrew Hughes (NIWA), Deborah
1021 Ballantine (then of NIWA), Graham McBride (NIWA), Murray Hicks (NIWA), David Hamilton
1022 (University of Waikato), Les Basher (LCR), Ian Fuller (Massey University), Roger Young
1023 (Cawthron Institute), Rien Visser (University of Canterbury), and David Lee-Jones (USDA
1024 FAS). Support for this project was also provided by numerous Regional Councils/Districts,
1025 including Auckland, Canterbury, Horizons, Tasman, and Waikato.

1026

1027 **References**

1028 ~~Abell, J. M., Özkundakci, D., Hamilton, D. P., and Miller, S. D.: Relationships between land use~~
1029 ~~and nitrogen and phosphorus in New Zealand lakes, Marine and Freshwater Research, 62,~~
1030 ~~162-175, 2011.~~

1031 ~~Ausseil, A. G. E., Chadderton, W. L., Gerbeaux, P., Stephens, R. T. T., and Leathwick, J. R.:~~
1032 ~~Applying systematic conservation planning principles to palustrine and inland saline~~
1033 ~~wetlands of New Zealand, Freshwater Biology, 56, 142-161, 2011.~~

1034 Ausseil, A. G. E., Dymond, J. R., Kirschbaum, M. U. F., Andrew, R. M., and Parfitt, R. L.:
1035 Assessment of multiple ecosystem services in New Zealand at the catchment scale,
1036 Environmental Modelling & Software, 43, 37-48, 10.1016/j.envsoft.2013.01.006, 2013.

1037 Australian and New Zealand Environment and Conservation Council (ANZECC): Australian and
1038 New Zealand Guidelines for Fresh and Marine Water Quality, Volume 1: The Guidelines,
1039 National Water Quality Management Strategy, Paper No. 4, Onehunga, 314 pp., 2000.

1040 Ballantine, D. J., and Davies-Colley, R. J.: Water quality trends in New Zealand rivers: 1989–
1041 2009, Environmental Monitoring and Assessment, 186, 1939-1950, 10.1007/s10661-013-
1042 3508-5, 2014.

1043 Ballantine, D. J., Hughes, A. O., and Davies-Colley, R. J.: Mutual relationships of suspended
1044 sediment, turbidity and visual clarity in New Zealand rivers, in: Proceedings of the
1045 International Association of Hydrological Sciences, New Orleans, 265-271, 2014.

1046 Baron, J. S., Poff, N. L., Angermeier, P. L., Dahm, C. N., Gleick, P. H., Hairston, N. G., Jackson,
1047 R. B., Johnston, C. A., Richter, B. D., and Steinman, A. D.: Meeting ecological and
1048 societal needs for freshwater, Ecological Applications, 12, 1247-1260, 2002.

1049 Basher, L. R., Hicks, D. M., Clapp, B., and Hewitt, T.: Sediment yield response to large storm
1050 events and forest harvesting, Motueka River, New Zealand, *New Zealand Journal of*
1051 *Marine and Freshwater Research*, 45, 333-356, 2011.

1052 ~~Bennett, E. M., Carpenter, S. R., and Caraco, N. F.: Human Impact on Erodable Phosphorus and~~
1053 ~~Eutrophication: A Global Perspective: Increasing accumulation of phosphorus in soil~~
1054 ~~threatens rivers, lakes, and coastal oceans with eutrophication, *BioScience*, 51, 227-234,~~
1055 ~~10.1641/0006-3568(2001)051[0227:hioepa]2.0.co;2, 2001.~~

1056 Bewsell, D., Monaghan, R. M., and Kaine, G.: Adoption of stream fencing among dairy farmers
1057 in four New Zealand catchments, *Environmental Management*, 40, 201-209, 2007.

1058 Blüthgen, N., Dormann, C. F., Prati, D., Klaus, V. H., Kleinebecker, T., Hölzel, N., Alt, F.,
1059 Boch, S., Gockel, S., Hemp, A., Müller, J., Nieschulze, J., Renner, S. C., Schöning, I.,
1060 Schumacher, U., Socher, S. A., Wells, K., Birkhofer, K., Buscot, F., Oelmann, Y.,
1061 Rothenwöhrer, C., Scherber, C., Tschardtke, T., Weiner, C. N., Fischer, M., Kalko, E. K.
1062 V., Linsenmair, K. E., Schulze, E.-D., and Weisser, W. W.: A quantitative index of land-
1063 use intensity in grasslands: Integrating mowing, grazing and fertilization, *Basic and*
1064 *Applied Ecology*, 13, 207-220, <http://dx.doi.org/10.1016/j.baae.2012.04.001>, 2012.

1065 Boesch, D. F.: Challenges and opportunities for science in reducing nutrient over-enrichment of
1066 coastal ecosystems, *Estuaries*, 25, 886-900, 10.1007/bf02804914, 2002.

1067 Brauman, K. A., Daily, G. C., Duarte, T. K., and Mooney, H. A.: The nature and value of
1068 ecosystem services: An overview highlighting hydrologic services, in: *Annual Review of*
1069 *Environment and Resources*, *Annual Review of Environment and Resources*, Annual
1070 *Reviews*, Palo Alto, 67-98, 2007.

- 1071 Brierley, G. P.: Landscape memory: the imprint of the past on contemporary landscape forms
1072 and processes, *Area*, 42, 76-85, 2010.
- 1073 Campbell, N., D'Arcy, B., Frost, A., Novotny, V., Sansom, A.: Diffuse pollution. An
1074 introduction to the problems and solutions, IWA publishing, London, 322 pp., 2004.
- 1075 ~~Clarkson, B. R., Ausseil, A. G. E., and Gerbeaux, P.: Wetland ecosystem services, in: Ecosystem~~
1076 ~~Services in New Zealand—Conditions and Trends, edited by: Dymond, J. R., Manaaki~~
1077 ~~Whenua Press, Lincoln, 192-202, 2013.~~
- 1078 Cleveland, W. S., and Devlin, S. J.: Locally weighted regression: An approach to regression
1079 analysis by local fitting, *Journal of the American Statistical Association*, 83, 596-610,
1080 1988.
- 1081 Close, M. E., and Davies-Colley, R. J.: Baseflow water chemistry in New Zealand rivers 1.
1082 Characterisation, *New Zealand Journal of Marine and Freshwater Research*, 24, 319-341,
1083 10.1080/00288330.1990.9516428, 1990.
- 1084 Croke, J. C., and Hairsine, P. B.: Sediment delivery in managed forests: a review, *Environmental*
1085 *Reviews*, 14, 59-87, 10.1139/a05-016, 2006.
- 1086 Davies-Colley, R. J.: River water quality in New Zealand: An introduction and overview, in:
1087 *Ecosystem Services in New Zealand - Conditions and Trends*, edited by: Dymond, J. R.,
1088 Manaaki Whenua Press, Lincoln, 432-447, 2013.
- 1089 Davies-Colley, R. J., Vant, W. N., and Smith, D. G.: Colour and Clarity of Natural Waters:
1090 Science and Management of Optical Water Quality, Blackburn Press, Caldwell, 310 pp.,
1091 2003.
- 1092 Davies-Colley, R. J., Smith, D. G., Ward, R. C., Bryers, G. G., McBride, G. B., Quinn, J. M., and
1093 Scarsbrook, M. R.: Twenty years of New Zealand's National Rivers Water Quality

1094 Network: Benefits of careful design and consistent operation, Journal of the American
1095 Water Resources Association, 47, 750-771, 2011.

1096 Davies-Colley, R. J., Ballantine, D. J., Elliott, A. H., Swales, A., Hughes, A. O., and Gall, M. P.:
1097 Light attenuation – a more effective basis for the management of fine suspended
1098 sediment than mass concentration?, Water Science and Technology, 69, 1867-1874. doi:
1099 10.2166/wst.2014.096, 2014.

1100 Davis, M.: Nutrient losses from forestry in the Lake Taupo catchment, Environment Waikato
1101 Technical Report 2005/37, Hamilton, 22, 2005.

1102 Davis, M.: Nitrogen leaching losses from forests in New Zealand, New Zealand Journal of
1103 Forestry Science, 44, 1-14, 10.1186/1179-5395-44-2, 2014.

1104 de Beurs, K. M., Owsley, B. C., and Julian, J. P.: Disturbance analyses of forests and grasslands
1105 with MODIS and Landsat in New Zealand, International Journal of Applied Earth
1106 Observation and Geoinformation, 45, Part A, 42-54,
1107 <http://dx.doi.org/10.1016/j.jag.2015.10.009>, 2016.

1108 ~~Duncan, R.: Regulating agricultural land use to manage water quality: The challenges for science
1109 and policy in enforcing limits on non-point source pollution in New Zealand, Land Use
1110 Policy, 41, 378-387, 2014.~~

1111 Dodds, W. K., and Welch, E. B.: Establishing nutrient criteria in streams, Journal of the North
1112 American Benthological Society, 19, 186-196, 2000.

1113 Dymond, J. R., Betts, H. D., and Schierlitz, C. S.: An erosion model for evaluating regional land-
1114 use scenarios, Environmental Modelling & Software, 25, 289-298,
1115 10.1016/j.envsoft.2009.09.011, 2010.

1116 Ekholm, P., Rankinen, K., Rita, H., Raike, A., Sjoblom, H., Raateland, A., Vesikko, L., Cano
1117 Bernal, J. E., and Taskinen, A.: Phosphorus and nitrogen fluxes carried by 21 Finnish
1118 agricultural rivers in 1985–2006, *Environmental Monitoring and Assessment*, 187, 1-17,
1119 10.1007/s10661-015-4417-6, 2015.

1120 Erb, K.-H., Haberl, H., Jepsen, M. R., Kuemmerle, T., Lindner, M., Muller, D., Verburg, P. H.,
1121 and Reenberg, A.: A conceptual framework for analysing and measuring land-use
1122 intensity, *Current Opinion in Environmental Sustainability*, 5, 464-470,
1123 <http://dx.doi.org/10.1016/j.cosust.2013.07.010>, 2013.

1124 Erb, K.-H., Fetzel, T., Haberl, H., Kastner, T., Kroisleitner, C., Lauk, C., Niedertscheider, M.,
1125 and Plutzer, C.: Beyond Inputs and Outputs: Opening the Black-Box of Land-Use
1126 Intensity, in: *Social Ecology: Society-Nature Relations across Time and Space*, edited by:
1127 Haberl, H., Fischer-Kowalski, M., Krausmann, F., and Winiwarter, V., Springer
1128 International Publishing, Cham, 93-124, 2016.

1129 Evans, R.: Soil erosion in the UK initiated by grazing animals, *Applied Geography*, 17, 127-141,
1130 [http://dx.doi.org/10.1016/S0143-6228\(97\)00002-7](http://dx.doi.org/10.1016/S0143-6228(97)00002-7), 1997.

1131 Fahey, B. D., Marden, M., and Phillips, C. J.: Sediment yields from plantation forestry and
1132 pastoral farming, coastal Hawke's Bay, North Island, New Zealand, *Journal of Hydrology*
1133 (New Zealand), 42, 27-38, 2003.

1134 Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S.,
1135 Coe, M. T., Daily, G. C., Gibbs, H. K., Helkowski, J. H., Holloway, T., Howard, E. A.,
1136 Kucharik, C. J., Monfreda, C., Patz, J. A., Prentice, I. C., Ramankutty, N., and Snyder, P.
1137 K.: Global consequences of land use, *Science*, 309, 570-574, 2005.

1138 Forest Owners Association (FOA): New Zealand plantation forest industry: Facts and figures
1139 2011/2012, Ministry for Primary Industries, Wellington, 46 pp., 2012.

1140 Fransen, P. J. B., Phillips, C. J., and Fahey, B. D.: Forest road erosion in New Zealand:
1141 overview, *Earth Surface Processes and Landforms*, 26, 165-174, 10.1002/1096-
1142 9837(200102)26:2<165::AID-ESP170>3.0.CO;2-#, 2001.

1143 Grohmann, C. H., Smith, M. J., and Riccomini, C.: Multiscale analysis of topographic surface
1144 roughness in the Midland Valley, Scotland, *IEEE Transactions on Geoscience and*
1145 *Remote Sensing*, 49, 1200-1213, 2011.

1146 [Gunn, A., and Rutherford, C.: The Dairying and Clean Streams Accord: Snapshot of Progress](#)
1147 [2011/2012, Ministry for Primary Industries, Wellington, 12, 2013.](#)

1148 Hart, B. T., Maher, B., and Lawrence, I.: New generation water quality guidelines for ecosystem
1149 protection, *Freshwater Biology*, 41, 347-359, 10.1046/j.1365-2427.1999.00435.x, 1999.

1150 Hicks, D. M., Gomez, B., and Trustrum, N. A.: Erosion thresholds and suspended sediment
1151 yields, Waipaoa River Basin, New Zealand, *Water Resources Research*, 36, 1129-1142,
1152 10.1029/1999WR900340, 2000.

1153 Hilton, J., O'Hare, M., Bowes, M. J., and Jones, J. I.: How green is my river? A new paradigm of
1154 eutrophication in rivers, *Science of the Total Environment*, 365, 66-83, 2006.

1155 Howard-Williams, C., Davies-Colley, R., Rutherford, K., and Wilcock, R.: Diffuse pollution and
1156 freshwater degradation: New Zealand Perspectives, International Conference of the IWA
1157 Diffuse Pollution Specialist Group, Beaupre, Quebec, 2010.

1158 [Hughes, A. O., and Quinn, J. M.: Before and After Integrated Catchment Management in a](#)
1159 [Headwater Catchment: Changes in Water Quality, *Environmental Management*, 54,](#)
1160 [1288-1305, 10.1007/s00267-014-0369-9, 2014.](#)

1161 ~~Julian, J. P., and Gardner, R. H.: Land cover effects on runoff patterns in eastern Piedmont~~
1162 ~~(USA) watersheds, Hydrological Processes, 28, 1525-1538, 10.1002/hyp.9692, 2014.~~

1163 Julian, J. P., Doyle, M. W., Powers, S. M., Stanley, E. H., and Riggsbee, J. A.: Optical water
1164 quality in rivers, Water Resources Research, 44, W10411,
1165 doi:10.1029/2007WR006457, 2008.

1166 Julian, J. P., Davies-Colley, R. J., Gallegos, C. L., and Tran, T. V.: Optical Water Quality of
1167 Inland Waters: A Landscape Perspective, Annals of the Association of American
1168 Geographers, 103, 309-318, 2013.

1169 Kamarinas, I., Julian, J. P., Hughes, A. O., Owsley, B. C., and de Beurs, K. M.: Nonlinear
1170 changes in land cover and sediment runoff in a New Zealand catchment dominated by
1171 plantation forestry and livestock grazing, Water, 8, 436, doi:10.3390/w8100436, 2016.

1172 Kronvang, B., Andersen, H. E., Børgesen, C., Dalgaard, T., Larsen, S. E., Bøgestrand, J., and
1173 Blicher-Mathiasen, G.: Effects of policy measures implemented in Denmark on nitrogen
1174 pollution of the aquatic environment, Environmental Science & Policy, 11, 144-152,
1175 <http://dx.doi.org/10.1016/j.envsci.2007.10.007>, 2008.

1176 Kuemmerle, T., Erb, K., Meyfroidt, P., Müller, D., Verburg, P. H., Estel, S., Haberl, H., Hostert,
1177 P., Jepsen, M. R., Kastner, T., Levers, C., Lindner, M., Plutzer, C., Verkerk, P. J., van der
1178 Zanden, E. H., and Reenberg, A.: Challenges and opportunities in mapping land use
1179 intensity globally, Current Opinion in Environmental Sustainability, 5, 484-493,
1180 <http://dx.doi.org/10.1016/j.cosust.2013.06.002>, 2013.

1181 Landcare Research (LCR): New Zealand Land Cover Database, available
1182 at: <http://www.lcdb.scinfo.org.nz/>, 2015.

1183 Larned, S. T., Scarsbrook, M. R., Snelder, T. H., Norton, N. J., and Biggs, B. J. F.: Water quality
1184 in low-elevation streams and rivers of New Zealand: recent state and trends in contrasting
1185 land-cover classes, *New Zealand Journal of Marine and Freshwater Research*, 38, 347-
1186 366, 2004.

1187 Larned, S., Snelder, T., Unwin, M., and McBride, G.: Water quality in New Zealand rivers:
1188 current state and trends, *New Zealand Journal of Marine and Freshwater Research*, DOI:
1189 10.1080/00288330.2016.1150309, 2016.

1190 Lobser, S. E., and Cohen, W.B.: MODIS tasselled cap: land cover characteristicsexpressed
1191 through transformed MODIS data, *Int. J. Remote Sens.*, 28,5079–5101, 2007.

1192 McAllister, T. G., Wood, S. A., and Hawes, I.: The rise of toxic benthic *Phormidium*
1193 proliferations: a review of their taxonomy, distribution, toxin content and factors
1194 regulating prevalence and increased severity, *Harmful Algae*, 55, 282-294, 2016.

1195 McDowell, R. W., Houlbrooke, D. J., Muirhead, R. W., Muller, K., Shepherd, M., and Cuttle, S.
1196 P.: *Grazed Pastures and Surface Water Quality*, Nova Science Publishers, New York, 238
1197 pp., 2008.

1198 Ministry of Business, Innovation & Employment (MBIE): *Business Growth Agenda: 2015*,
1199 available at: [http://www.mbie.govt.nz/info-services/business/business-growth-
1200 agenda/pdf-and-image-library/towards-2025/mb13078-1139-bga-report-01-export-
1201 markets-09sept-v17-fa-web.PDF](http://www.mbie.govt.nz/info-services/business/business-growth-agenda/pdf-and-image-library/towards-2025/mb13078-1139-bga-report-01-export-markets-09sept-v17-fa-web.PDF), 2015.

1202 Ministry for the Environment (MfE): *Land-use and carbon analysis system (LUCAS): Satellite*
1203 *imagery interpretation guide for land-use classes (2nd edition)*, Ministry for the
1204 Environment, Wellington, 73 pp., 2012.

1205 ~~Mitsch, W. J., and Gosselink, J. G.: The value of wetlands: importance of scale and landscape~~
1206 ~~setting, *Ecological Economics*, 35, 25–33, [10.1641/0006-3568\(2001\)051\[0373:rlttg\]2.0.co;2](http://dx.doi.org/10.1016/S0921-
1207 8009(00)00165-8, 2000.</p><p>1208 Mitsch, W. J., Day, J. W., Gilliam, J. W., Groffman, P. M., Hey, D. L., Randall, G. W., and
1209 Wang, N.: Reducing Nitrogen Loading to the Gulf of Mexico from the Mississippi River
1210 Basin: Strategies to Counter a Persistent Ecological Problem: Ecotechnology—the use of
1211 natural ecosystems to solve environmental problems—should be a part of efforts to
1212 shrink the zone of hypoxia in the Gulf of Mexico, <i>BioScience</i>, 51, 373–388,
1213 <a href=), 2001.~~

1214 Monaghan, R. M., Hedley, M. J., Di, H. J., McDowell, R. W., Cameron, K. C., and Ledgard, S.
1215 F.: Nutrient management in New Zealand pastures— recent developments and future
1216 issues, *N. Z. J. Agric. Res.*, 50, 181-201, [10.1080/00288230709510290](http://dx.doi.org/10.1080/00288230709510290), 2007.

1217 Moosmann, L., Müller, B., Gächter, R., Wüest, A., Butscher, E., and Herzog, P.: Trend-oriented
1218 sampling strategy and estimation of soluble reactive phosphorus loads in streams, *Water*
1219 *Resources Research*, 41, W01020, [10.1029/2004WR003539](http://dx.doi.org/10.1029/2004WR003539), 2005.

1220 Morris, S. T.: Sheep and beef cattle production systems, in: *Ecosystem Services in New Zealand*
1221 *- Conditions and Trends*, edited by: Dymond, J. R., Manaaki Whenua Press, Lincoln, 79-
1222 84, 2013.

1223 Motha, J. A., Wallbrink, P. J., Hairsine, P. B., and Grayson, R. B.: Determining the sources of
1224 suspended sediment in a forested catchment in southeastern Australia, *Water Resources*
1225 *Research*, 39, 1056, doi:10.1029/2001WR000794, 2003.

1226 Newsome, P. F. J., R.H., W., and Willoughby, E. J.: *Land Resource Information System Spatial*
1227 *Data Layers: Data Dictionary*, Palmerston North, 75, 2008.

1228 New Zealand Farm Forestry Association (NZFFA): National Exotic Forest Description: as at 1
1229 April 2014, Ministry for Primary Industries, Wellington, 77 pp., 2014.

1230 OECD/FAO: OECD-FAO Agricultural Outlook 2015-2024, OECD/Food and Agriculture
1231 Organization of the United Nations Paris, 148, 2015.

1232 Parkyn, S. M., Davies-Colley, R. J., Scarsbrook, M. R., Halliday, N. J., Nagels, J. W., Marden,
1233 M., and Rowan, D.: Pine afforestation and stream health: a comparison of land-use in two
1234 soft rock catchments, East Cape, New Zealand, New Zealand Natural Sciences, 31, 113-
1235 135, 2006.

1236 Parliamentary Commissioner for the Environment (PCE): Water quality in New Zealand: land
1237 use and nutrient pollution, Parliamentary Commissioner for the Environment,
1238 Wellington, 82 pp., available at: [http://www.pce.parliament.nz/media/1275/pce-water-](http://www.pce.parliament.nz/media/1275/pce-water-quality-land-use-web-amended.pdf)
1239 [quality-land-use-web-amended.pdf](http://www.pce.parliament.nz/media/1275/pce-water-quality-land-use-web-amended.pdf), 2013.

1240 Phillips C. J., Marden M., and Rowan D.: Sediment yield following plantation harvesting,
1241 Coromandel Peninsula, North Island, New Zealand, Journal of Hydrology (NZ), 44, 29-
1242 44, 2005.

1243 Ramankutty, N., Graumlich, L., Achard, F., Alves, D., Chhabra, A., DeFries, R. S., Foley, J. A.,
1244 Geist, H., Houghton, R. A., Goldewijk, K. K., Lambin, E. F., Millington, A., Rasmussen,
1245 K., Reid, R. S., and Turner, B. L.: Global Land-Cover Change: Recent Progress,
1246 Remaining Challenges, in: Land-Use and Land-Cover Change: Local Processes and
1247 Global Impacts, edited by: Lambin, E. F., and Geist, H., Springer Berlin Heidelberg,
1248 Berlin, Heidelberg, 9-39, 2006.

1249 Saunders, W. M. H.: Phosphate retention by New Zealand soils and its relationship to free
1250 sesquioxides, organic matter, and other soil properties, N. Z. J. Agric. Res., 8, 30-57,
1251 DOI: 10.1080/00288233.1965.10420021, 1965.

1252 Scarsbrook, M. R.: State and trends in the National River Water Quality Network (1989-2005),
1253 Ministry for the Environment, ME number 778, 48 pp., available
1254 at: <http://www.mfe.govt.nz/publications/ser/water-quality-network-nov06/index.html>,
1255 2006.

1256 Scarsbrook, M. R., McBride, C. G., McBride, G. B., and Bryers, G. G.: Effects of climate
1257 variability on rivers: Consequences for long term water quality analysis, Journal of the
1258 American Water Resources Association, 39, 1435-1447, 2003.

1259 ~~Schallenberg, M. and Sorrell, B.: Regime shifts between clear and turbid water in New Zealand~~
1260 ~~lakes: Environmental correlates and implications for management and restoration, New~~
1261 ~~Zealand Journal of Marine and Freshwater Research, 43, 701-712, DOI:~~
1262 ~~10.1080/00288330909510035, 2009.~~

1263 ~~Schallenberg, M., de Winton, M. D., Verburg, P., Kelly, D. J., Hamill, K. D., and Hamilton, D.~~
1264 ~~P.: Ecosystem services of lakes, in: Ecosystem Services in New Zealand—Conditions and~~
1265 ~~Trends, edited by: Dymond, J. R., Manaaki Whenua Press, Lincoln, 203-225, 2013.~~

1266 Smith, D. G., McBride, G. B., Bryers, G. G., Wisse, J., and Mink, D. F. J.: Trends in New
1267 Zealand's National River Water Quality Network, New Zealand Journal of Marine and
1268 Freshwater Research, 30, 485-500, 10.1080/00288330.1996.9516737, 1996.

1269 Smith, D. G., Davies-Colley, R. J., Knoeff, J., and Slot, G. W. J.: Optical characteristics of New
1270 Zealand rivers in relation to flow, Journal of the American Water Resources Association,
1271 33, 301–312, 1997.

1272 Snelder, T. H.: Definition of a multivariate classification of New Zealand Lakes, National
1273 Institute of Water & Atmospheric Research Ltd, Christchurch, 32, 2006.

1274 Snelder, T., Biggs, B., and Weatherhead, M.: New Zealand River Environment Classification
1275 User Guide (REC, v2), Produced for the Ministry for the Environment by the National
1276 Institute of Water and Atmospheric Research (NIWA), Publication number ME 1026,
1277 2010.

1278 Stamm, C., Jarvie, H. P., and Scott, T.: What's More Important for Managing Phosphorus:
1279 Loads, Concentrations or Both?, Environmental science & technology, 48, 23-24,
1280 10.1021/es405148c, 2014.

1281 Statistics New Zealand (StatsNZ): 2012 Agricultural Census tables, available
1282 at: [http://www.stats.govt.nz/browse_for_stats/industry_sectors/agriculture-horticulture-](http://www.stats.govt.nz/browse_for_stats/industry_sectors/agriculture-horticulture-forestry/2012-agricultural-census-tables/livestock.aspx)
1283 [forestry/2012-agricultural-census-tables/livestock.aspx](http://www.stats.govt.nz/browse_for_stats/industry_sectors/agriculture-horticulture-forestry/2012-agricultural-census-tables/livestock.aspx), 2015.

1284 Tait, A., and Turner, R.: Generating Multiyear Gridded Daily Rainfall over New Zealand,
1285 Journal of Applied Meteorology, 44, 1315-1323, 10.1175/JAM2279.1, 2005.

1286 Trimble, S. W., and Mendel, A. C.: The cow as a geomorphic agent—A critical review,
1287 Geomorphology, 13, 233–253, 1995.

1288 Vant, B., and Smith, P.: Nutrient concentrations and water ages in 11 streams flowing into
1289 Lake Taupo, Environment Waikato Technical Report 2002/18R, 26, 2004.

1290 Varanka, S., and Luoto, M.: Environmental determinants of water quality in boreal rivers based
1291 on partitioning methods, River Research and Applications, 28, 1034-1046,
1292 10.1002/rra.1502, 2012.

1293 ~~Verhoeven, J. T. A., Arheimer, B., Yin, C., and Hefting, M. M.: Regional and global concerns~~
1294 ~~over wetlands and water quality, Trends in ecology & evolution, 21, 96-103,~~
1295 ~~<http://dx.doi.org/10.1016/j.tree.2005.11.015>, 2006.~~

1296 ~~Vitousek, P. M., Aber, J. D., Howarth, R. W., Likens, G. E., Matson, P. A., Schindler, D. W.,~~
1297 ~~Schlesinger, W. H., and Tilman, G. D.: Human alteration of the global nitrogen cycle:~~
1298 ~~Sources and consequences, Ecological Applications, 7, 737-750, 1997a.~~

1299 Vitousek, P. M., Mooney, H. A., Lubchenco, J., and Melillo, J. M.: Human domination of Earth's
1300 ecosystems, Science, 277, 494, 1997b.

1301 Vorosmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P.,
1302 Glidden, S., Bunn, S. E., Sullivan, C. A., Liermann, C. R., and Davies, P. M.: Global
1303 threats to human water security and river biodiversity, Nature, 467, 555-561, 2010.

1304 ~~Walling, D. E.: Human impact on land-ocean sediment transfer by the world's rivers,~~
1305 ~~Geomorphology, 79, 192-216, 2006.~~

1306 Webb, T. W., and Wilson, A. D.: A Manual of Land Characteristics for Evaluation of Rural land,
1307 Landcare Research New Zealand Ltd, Lincoln, 1995.

1308 ~~Wetzel, R. G.: Limnology: Lake and River Ecosystems, Academic Press, San Diego, 1006 pp.,~~
1309 ~~2001.~~

1310 Wilcock, R. J., Betteridge, K., Shearman, D., Fowles, C. R., Scarsbrook, M. R., Thorrold, B. S.,
1311 and Costall, D.: Riparian protection and on-farm best management practices for
1312 restoration of a lowland stream in an intensive dairy farming catchment: A case study,
1313 New Zealand Journal of Marine and Freshwater Research, 43, 803-818,
1314 doi:10.1080/00288330909510042, 2009.

1315 Wilmshurst, J. M., Anderson, A. J., Higham, T. F. G., Worthy, T. H.: Dating the late prehistoric
1316 dispersal of Polynesians to New Zealand using the commensal Pacific rat, Proceedings of
1317 the National Academy of Sciences, 105, 7676-7680, 2008.

1318 Wood, S., Hawes, I., McBride, G., Truman, P., and Dietrich, D.: Advice to inform the
1319 development of a benthic cyanobacteria attribute, Cawthron Institute, Report No. 2752,
1320 Nelson, 91, 2015.

1321 Woods, R., Bidwell, V., Clothier, B., Green, S., Elliott, S., Shankar, U., Harris, S., Hewitt, A.,
1322 Gibb, R., Parfitt, R., and Wheeler, D.: The CLUES Project: Predicting the effects of land-
1323 use on water quality - Stage II, National Institute of Water & Atmospheric Research,
1324 Christchurch, 2006.

1325 Woodward, S. J. R.: Bite mechanics of cattle and sheep grazing grass-dominant swards, Applied
1326 Animal Behaviour Science, 56, 203-222, 1998.

1327 Zobrist, J., and Reichert, P.: Bayesian estimation of export coefficients from diffuse and point
1328 sources in Swiss watersheds, Journal of Hydrology, 329, 207-223,
1329 <http://dx.doi.org/10.1016/j.jhydrol.2006.02.014>, 2006.

1330

1331

1332

1333

1334

1335 **Tables**

1336

1337 Table 1. Water quality variables measured by the National River Water Quality Network
 1338 (NRWQN) obtained from monthly **grab** samples from 1989 to 2014 for 77 catchments. [Details](#)
 1339 [on analytical methods can be found in Davies-Colley et al. \(2011\).](#)

Variable	Definition (units)
<i>Q</i>	Water discharge (m ³ /s)
<i>T_w</i>	Water temperature (°C)
<i>DO</i>	Dissolved oxygen (%)
<i>COND</i>	Water conductivity (µS/cm)
<i>pH_w</i>	Water pH (-log ₁₀ [H ⁺])
<i>CLAR</i>	Horizontal visual water clarity from black disc sighting range (m)
<i>TURB</i>	Water turbidity (NTU)
<i>CDOM</i>	Colored dissolved organic matter, measured as spectrophotometric absorbance of a membrane filtrate at 440 nm (m ⁻¹)
<i>TN</i>	Total nitrogen (mg/m ³)
<i>NO_x</i>	Oxidized nitrogen in nitrate and nitrite forms (mg/m ³)
<i>TP</i>	Total phosphorus (mg/m ³)
<i>DRP</i>	Dissolved reactive phosphorus (mg/m ³)

1340

1341

1342 Table 2. Landscape variables characterizing the 77 catchments of the National River Water
 1343 Quality Network (NRWQN). More details on sources for these data can be found in Methods
 1344 section.

Variable	Definition (units)	Source (resolution/scale)
Morphometric variables		
Area (A)	Total catchment area above monitoring site (km^2)	National Elevation Dataset (30 m)
Drainage density (D_d)	Total length of streams per catchment area (km/km^2)	River Environment Classification, v2 (1:24,000)
Catchment slope (S_c)	Mean slope across entire catchment (degrees)	National Elevation Dataset (30 m)
Ruggedness (R_r)	Standard deviation of catchment slope (degrees)	National Elevation Dataset (30 m)
Soil variables		
Silt-clay percentage ($SC\%$)	Percentage of catchment surface soils dominated by clayey or silty soils (%)	Fundamental Soil Layers (1:63,360)
Soil depth (Z_s)	Mean maximum potential rooting depth across catchment (m)	Fundamental Soil Layers (1:63,360)
Soil pH (pH_s)	Mean pH at 0.2-0.6 m depth across catchment ($-\log_{10}[\text{H}^+]$)	Fundamental Soil Layers (1:63,360)
Cation exchange capacity (CEC)	Weighted mean CEC at 0-0.6 m depth across catchment (cmoles $[\text{+}]/\text{kg}$)	Fundamental Soil Layers (1:63,360)
Organic matter percentage ($OM\%$)	Weighted mean of total carbon at 0-0.2 m depth across catchment (%)	Fundamental Soil Layers (1:63,360)
Phosphate retention (P_{ret})	Weighted mean of phosphate retention at 0-0.2 m depth across catchment (%)	Fundamental Soil Layers (1:63,360)
Hydro-climatological variables		
Median annual precipitation (MAP)	Median annual precipitation averaged across catchment (mm/y)	NIWA National Climate Database (5 km)
Median annual temperature (MAT)	Median annual temperature averaged across catchment ($^{\circ}\text{C}$)	NIWA National Climate Database (5 km)
Median annual sunshine (MAS)	Median annual sunshine hours averaged across catchment (hours/y)	NIWA National Climate Database (5 km)
Median discharge (Q_{50})	Median discharge from NRWQN samples during 1989-2014 (m^3/s)	NRWQN (catchment)

Relative water storage (<i>RWS</i>)	Proportion of annual Q_{50} stored in reservoirs/lakes (m^3/m^3)	Freshwater Environments New Zealand (1:50,000)
--	---	---

Land Use and Land Disturbance variables

Land use	Percent of catchment that is occupied by each land use (%); see Table 3 for land uses	Land Cover Database (LCDB, v 4.1), 2001 (1 ha)
High-producing pasture disturbance (<i>D_{HG}</i>)	Percent of high-producing grasslands within catchment that is disturbed (%), based on aggregate of 463-m pixels within catchment	de Beurs et al., 2016 (463 m; 8-day)
Plantation forestry disturbance (<i>D_{PF}</i>)	Percent of plantation forestry within catchment that is disturbed (%), based on aggregate of 463-m pixels within catchment	de Beurs et al., 2016 (463 m; 8-day)
Catchment disturbance (<i>D_C</i>)	Percent of catchment that is disturbed (%), based on aggregate of 463-m pixels within catchment	de Beurs et al., 2016 (463 m; 8-day)
Stock unit density (<i>SUD</i>)	Catchment-averaged stock unit density for dairy (<i>da</i>), beef (<i>be</i>), deer (<i>de</i>), and sheep (<i>sh</i>) in 2011 (SU/ha); subscripts are used to isolate SUD by livestock type	Ausseil et al., 2013 (1 ha)
Change in stock unit density (<i>SUD₂₀₁₂₋₁₉₉₀</i>)	Difference between SUD in 2012 and 1990 (SU/ha)	Statistics NZ (territorial authority)

1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362

1363 Table 3. Land use classification used in this study, aggregated from the LUCAS (v11) and
 1364 LCDB (v4.1) land use/cover datasets.

Class (abbreviation)	Description	LUCAS classes	LCDB classes	2012 national coverage (%) LUCAS / LCDB
Non-plantation forest (NF)	All non-plantation forests $\geq 5\text{m}$; does not include Manuka/Kanuka	71	68, 69	29.2 / 23.9
Plantation forest (PF)	All forests that are planted for the purpose of harvesting	72,73	64, 71	7.9 / 7.6
Shrub/Grassland (SG)	All shrubs $< 5\text{m}$ and grasses that are not intensively managed	74, 76	41-44, 50-58	33.0 / 25.4
High-producing grassland (HG)	High-quality pasture grasses that are intensively managed	75	40	21.6 / 33.0
Perennial cropland (PC)	Orchards and vineyards	77	33	0.4 / 0.4
Annual cropland (AC)	All annual crops and cultivated bare ground	78	30	1.4 / 1.4
Open water (OW)	Rivers, lakes/reservoirs, ponds, and estuaries	79	20-22	1.9 / 2.0
Vegetated wetland (VW)	Herbaceous or woody vegetation periodically flooded; includes mangroves	80	45-47, 70	0.5 / 0.7
Urban (UR)	Built-up areas, infrastructure, transportation networks, and urban parks/open spaces	81	1-5	0.8 / 0.9
Barren/Other (BO)	Bare rock, sand, gravel and other areas not dominated by vegetation; includes mining and permanent ice/snow	82	6-16	3.3 / 4.8

1365
 1366

1367 Table 4. Statistical description of landscape variables for the 77 NRWQN catchments. Refer to
 1368 Tables 2 and 3 for variable descriptions.

Variable	Units	Minimum	Median	Maximum	Mean \pm SD
Morphometric Variables					
Area (<i>A</i>)	km ²	26	1126	20539	2639 \pm 3714
Drainage density (<i>D_d</i>)	km/km ²	1.30	1.59	2.61	1.60 \pm 0.16
Catchment slope (<i>S_c</i>)	degrees	3.4	15.9	30.3	16.3 \pm 6.8
Ruggedness (<i>R_r</i>)	degrees	3.4	10.8	15.8	10.6 \pm 2.4
Soil Variables					
Silt-clay percentage (<i>SC%</i>)	%	0	47.3	98.7	44.0 \pm 31.6
Soil depth (<i>Z_s</i>)	m	0.55	0.96	1.50	1.02 \pm 0.22
Soil pH (<i>pH</i>)	$-\log_{10}[\text{H}^+]$	4.8	5.6	6.5	5.6 \pm 0.3
Cation exchange capacity (<i>CEC</i>)	cmoles [+]/kg	11.6	18.7	33.5	18.8 \pm 4.6
Organic matter percentage (<i>OM%</i>)	%	2.8	6.7	23.2	7.2 \pm 2.9
Phosphate retention (<i>P_{ret}</i>)	%	19.9	39.0	77.8	41.5 \pm 12.2
Hydro-climatological Variables					
Median annual precipitation (<i>MAP</i>)	mm/y	533	1652	7044	1778 \pm 873
Median annual temperature (<i>MAT</i>)	°C	5.0	9.9	15.1	9.9 \pm 2.4
Median annual sunshine (<i>MAS</i>)	hours/y	1325	1856	2116	1841 \pm 146
Median discharge (<i>Q₅₀</i>)	m ³ /s	0.4	26.0	515.0	69.6 \pm 112.6
Relative water storage (<i>RWS</i>)	m ³ /m ³	0	0	29.2	1.1 \pm 3.7
Land Use Variables					
Non-plantation forest (<i>NF</i>)	%	0.1	20.5	94.1	26.7 \pm 23.3
Plantation forest (<i>PF</i>)	%	0	3.3	69.8	8.2 \pm 12.3
Shrub/Grassland (<i>SG</i>)	%	0.4	21.7	82.3	26.6 \pm 20.2
High-producing grassland (<i>HG</i>)	%	0	21.6	91.2	30.9 \pm 26.2
Perennial cropland (<i>PC</i>)	%	0	0	1.3	0.1 \pm 0.2
Annual cropland (<i>AC</i>)	%	0	0.1	7.9	0.6 \pm 1.4
Open water (<i>OW</i>)	%	0	0.4	25.6	1.9 \pm 4.3

Vegetated wetland (<i>VW</i>)	%	0	0.1	2.2	0.3 ± 0.4
Urban (<i>UR</i>)	%	0	0.1	5.8	0.4 ± 0.7
Barren/Other (<i>BO</i>)	%	0	1.3	30.0	4.4 ± 6.5
Land Disturbance Variables					
Catchment disturbance (<i>D_C</i>)	%	0	3.4	10.5	3.6 ± 2.1
<i>HG</i> disturbance (<i>D_{HG}</i>)	%	0	4.4	34.9	6.0 ± 6.4
<i>PF</i> disturbance (<i>D_{PF}</i>)	%	0	9.9	27.8	10.4 ± 6.7
Stock unit density (<i>SUD</i>)	SU/ha	0	2.2	16.1	3.2 ± 3.1
Dairy <i>SUD</i> (<i>SUD_{da}</i>)	SU/ha	0	0.2	15.4	1.2 ± 2.4
Beef <i>SUD</i> (<i>SUD_{be}</i>)	SU/ha	0	0.5	3.5	0.7 ± 0.8
Sheep <i>SUD</i> (<i>SUD_{sh}</i>)	SU/ha	0	0.6	4.5	1.2 ± 1.3
Deer <i>SUD</i> (<i>SUD_{de}</i>)	SU/ha	0	0	0.2	0 ± 0

1369

1370

1371

1372 Table 5. Statistical description of medians of water quality variables for the 77 NRWQN
 1373 catchments. Note that the ratio of mean/median can be used as an index of data skewness.

Variable	Units	Minimum	Median	Maximum	Mean \pm SD
<i>T_w</i>	°C	7.2	12.2	16.9	12.4 \pm 2.4
<i>DO</i>	%	75.5	100.8	113.1	100.0 \pm 4.7
<i>COND</i>	μ S/cm	39	92	528	113 \pm 83
<i>pH_w</i>	$-\log_{10}[\text{H}^+]$	6.9	7.7	8.5	7.7 \pm 0.3
<i>CLAR</i>	m	0.1	1.5	9.8	2.1 \pm 1.8
<i>TURB</i>	NTU	0.3	2.1	82	4.2 \pm 9.4
<i>CDOM</i>	m^{-1}	0.1	0.7	4.6	0.9 \pm 0.8
<i>TN</i>	mg/m ³	40	259	2162	369 \pm 361
<i>NO_x</i>	mg/m ³	1	107	1852	230 \pm 302
<i>TP</i>	mg/m ³	3	15	115	24 \pm 24
<i>DRP</i>	mg/m ³	0.5	5.0	66.2	8.6 \pm 11.2

1374

Table 6. River water quality trends from 1989-2014. The table reports numbers of sites (out of 77) in different categories of water quality time trend. All variables were flow-adjusted except flow and water temperature. Significant trends were taken to be those with a p-value < 0.05 in the Seasonal Kendall test. Meaningful trends were taken to be those which also had a magnitude (RSKSE) greater than 1% per year.

Direction of trend	River Water Quality Variable (1989-2014)											
	<i>Q</i>	<i>T_w</i>	<i>DO</i>	<i>COND</i>	<i>pH_w</i>	<i>CLAR</i>	<i>TURB</i>	<i>CDOM</i>	<i>TP</i>	<i>DRP</i>	<i>TN</i>	<i>NO_x</i>
Meaningful Increase	1	0	0	4	0	29	17	1	8	17	27	24
Significant Increase	1	21	6	48	12	5	1	1	6	3	6	3
No Significant Trend	67	54	42	19	48	39	50	56	52	49	39	37
Significant Decrease	3	2	29	6	17	2	0	13	4	5	3	1
Meaningful Decrease	5	0	0	0	0	2	9	6	7	3	2	12

Table 7. Correlations of water quality (median values) vs. the major land uses, livestock densities, and median catchment disturbance of the 77 NRWQN catchments. All values represent Spearman correlation coefficients (r_s). Nonsignificant relationships ($p \geq 0.05$) are denoted by *NS*. T_w was not included because of its strong latitudinal trend. DO and pH_w were not included because they had no significant relationships with land use. SUD_{cattle} is the combination of dairy and beef cattle.

	<i>HG</i>	<i>SG</i>	<i>NF</i>	<i>PF</i>	<i>OW</i>	SUD_{da}	SUD_{be}	SUD_{cattle}	SUD_{sh}	SUD_{de}	D_C	D_{HG}	D_{PF}
<i>COND</i>	0.57	-0.53	<i>NS</i>	0.53	<i>NS</i>	0.44	0.63	0.60	0.35	<i>NS</i>	<i>NS</i>	-0.25	<i>NS</i>
<i>CLAR</i>	-0.45	<i>NS</i>	0.28	-0.31	<i>NS</i>	-0.41	-0.49	-0.49	-0.40	<i>NS</i>	<i>NS</i>	<i>NS</i>	-0.27
<i>TURB</i>	0.46	<i>NS</i>	-0.27	0.28	<i>NS</i>	0.38	0.50	0.48	0.40	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>
<i>CDOM</i>	0.56	-0.55	<i>NS</i>	0.24	-0.29	0.48	0.53	0.57	0.24	<i>NS</i>	<i>NS</i>	-0.33	<i>NS</i>
<i>TN</i>	0.82	-0.56	-0.37	0.46	-0.25	0.79	0.75	0.85	0.60	0.26	<i>NS</i>	-0.40	<i>NS</i>
NO_x	0.70	-0.53	-0.25	0.44	-0.25	0.77	0.65	0.79	0.51	0.28	<i>NS</i>	-0.39	<i>NS</i>
<i>TP</i>	0.66	-0.54	-0.32	0.48	<i>NS</i>	0.58	0.66	0.72	0.42	<i>NS</i>	<i>NS</i>	-0.24	<i>NS</i>
<i>DRP</i>	0.59	-0.65	<i>NS</i>	0.50	-0.43	0.58	0.58	0.66	0.31	<i>NS</i>	<i>NS</i>	-0.32	<i>NS</i>

Table 8. Stepwise regressions of water quality variables (median values) on landscape descriptors (forward selection, $p < 0.05$). Signs of coefficients indicate whether the relationship is proportional (+) or inverse (-). Int is model intercept. [Scatterplots that characterize the primary and secondary explanatory variables are displayed in Figure 5.](#)

Water Quality Variable	Step	Landscape Variable	Model Estimate	Multivariate sequential r^2
<i>CLAR</i>	1	<i>HG</i>	-0.03	0.17
	2	<i>OW</i>	0.18	0.27
	3	<i>Q₅₀</i>	-0.01	0.35
	4	<i>PF</i>	-0.03	0.39
	Int		3.16	
<i>TN</i>	1	<i>SUD_{cattle}</i>	77.05	0.62
	2	<i>HG</i>	4.26	0.68
	3	<i>PF</i>	5.16	0.69
	4	<i>SC%</i>	1.80	0.72
	Int		-33.95	
<i>NO_x</i>	1	<i>SUD_{cattle}</i>	86.15	0.58
	Int		62.65	
<i>TP</i>	1	<i>SUD_{cattle}</i>	5.47	0.41
	2	<i>PF</i>	0.64	0.52
	Int		7.75	
<i>DRP</i>	1	<i>SUD_{cattle}</i>	2.23	0.31
	2	<i>PF</i>	0.38	0.48
	Int		1.14	

Figures

Figure 1. Land use and location of the 77 National River Water Quality Network (NRWQN) catchments. Catchment ID colors refer to dominant land use (>50%). Catchments with no dominant land use are black.

Figure 2. Changes in land use areal coverage, livestock, and fertilizer inputs across New Zealand 1989/1990 vs. 2011/2012. Nitrogen fertilizers include urea and ammonium sulphate. Phosphorus fertilizers include superphosphate and diammonium phosphate.

Figure 23. Disturbance frequency of North Island per 463-m pixel, based on interpretation of MODIS data 2000-2013.

Figure 34. Disturbance frequency of South Island per 463-m pixel, based on interpretation of MODIS data 2000-2013.

Figure 45. Multivariate relationships between major water quality variables (median value for each site) and land use variables. For each plot, the primary explanatory variable from the stepwise regression (Table 8) is the x-axis, with bubble color representing the secondary explanatory variable. Note that oxidized nitrogen (NO_x) did not have a secondary explanatory variable. Selected catchments discussed in the text are labeled.

Figure 56. River water quality classes for upland (A) and lowland (B) catchments in New Zealand: I. clean river with high visual water clarity (CLAR) and low dissolved inorganic nutrients (DIN); II. sediment-impacted river with low CLAR and low DIN; III. nutrient-impacted river with high CLAR and high DIN; and IV. sediment- and nutrient-impacted river with low CLAR and high DIN. Classes are organized by ANZECC (2000) trigger values for water clarity (x-axis) and, ~~DIN trigger values can be discriminated for~~ NO_x (y-axis). Catchments that exceed ANZECC guidelines for DRP are indicated in by ~~and DRP~~ (grey-filled markers). Arrows

indicate direction of~~whether the~~ trend over the 26 years inclusive from 1989~~from 1989-2014~~
if~~was~~ significant (dashed) or meaningful (solid). No arrow means the trend was not significant.