



Climate change and climate-driven disturbances in the San Juan River sub-basin of the Colorado River

- 5 Katrina E. Bennett<sup>1</sup>, Theodore Bohn<sup>2,3</sup>, Kurt Solander<sup>1</sup>, Nathan G. McDowell<sup>1</sup>, Chonggang Xu<sup>1</sup>, Enrique Vivoni<sup>3,4</sup>  
and Richard S. Middleton<sup>1</sup>

<sup>1</sup>Earth and Environmental Sciences, Los Alamos National Lab, Los Alamos, NM, 87545

<sup>2</sup>Julie Ann Wrigley Global Institute of Sustainability, Arizona State University, Tempe, AZ, 85287

- 10 <sup>3</sup>School of Earth and Space Exploration, Arizona State University, Tempe, AZ, 85287

<sup>4</sup>School of Sustainable Engineering and the Built Environment, Arizona State University, Tempe, AZ, 85287

- 15 Correspondence to: Katrina E. Bennett, kbennett@lanl.gov, Earth and Environmental Sciences, Los Alamos  
National Lab, Los Alamos, NM, 87545, 505-664-0698



## Abstract

Accelerated climate change and associated forest disturbances in the Southwestern USA are anticipated to have substantial impacts on regional water resources. Few studies have quantified the impact of both climate change and land cover disturbances on water balances at the basin scale, and none at the regional scale. In this work, we evaluate the impacts of forest disturbances and climate change for a headwater basin to the Colorado River, the San Juan River watershed, using a robustly-calibrated (Nash Sutcliffe 0.80) hydrologic model run with updated formulations that improve estimates of evapotranspiration for semi-arid regions. Our results show that future disturbances will have a substantial impact on streamflow with implications for water resource management. Our findings are in contradiction with conventional thinking that forest disturbances reduce ET and increase streamflow. In this study, annual average regional streamflow under the coupled climate-disturbances scenarios is at least 6–11% lower than those scenarios accounting for climate change alone, and for forested zones of the San Juan River basin streamflow is 15–21% lower. The monthly signals of altered streamflow point to an emergent streamflow pattern related to changes in forests of the disturbed systems. Exacerbated reductions of mean and low flows under disturbance scenarios indicate a high risk of lower water availability for forested headwater systems to the Colorado River basin. These findings also indicate that explicit representation of land cover disturbances is required in modelling efforts that consider the impact of climate change on water resources.



## 1 Introduction

Widespread forest disturbances are projected to increase with climate change (McDowell et al., 2016; Allen et al., 2010; Van Mantgem et al., 2009) and this will have major implications for ecosystem services (Anderegg et al., 2013). These ecosystems services impact the provision of food, water, and energy and therefore necessitate a more robust understanding of how the landscape will respond to the associated shifts in timing and frequency of streamflow and a firm understanding of the dominant processes driving these changes. However, the impacts of changes in forest ecosystems on water resources remain understudied, despite its importance for natural resource management and energy production in affected basins around the globe. For instance, climate-induced changes in forests will feed back to the climate by altering the albedo and reducing the carbon sink, which is anticipated to further transform ecosystems in either positive or negative ways (Dale et al., 2001). This is particularly salient for regions of the United States such as the Colorado River basin (CRB) where forest cover is anticipated to be significantly impacted by a higher incidence of wildfire, drought, and pest infestations (Williams et al., 2013).

To date, predictions of future streamflow in forested river basins under future changes in climate and land cover have exhibited wide disagreement as to the strength and even the sign of change. Water yields and peak streamflows in North American river basins are projected to either increase or show no response to changing forest cover (Adams et al., 2012; Schnorbus et al., 2010; Guardiola-Claramonte et al., 2011). Causes of the reported changes have been related to topography (Schnorbus et al., 2010) and climate variability (Allen et al., 2015). Other reported causes are secondary impacts occurring as a result of forest mortality that enhance processes such as radiation (Royer et al., 2011; Varhola et al., 2010), changes in albedo (Winkler et al., 2010), changes in evapotranspiration (Zou et al., 2010; Kang et al., 2006), and soil moisture states (Dale et al., 2001). Snow has been reported to play an important role in changes in streamflow (Solander et al., 2017; Bennett et al., 2015) through increased snow accumulation and snow melt (Bewley et al., 2010; Boon, 2007), and snow cover duration in disturbed forests (Boon, 2009). Some research points to the fact that we have a limited understanding on why or how streamflow changes in the future under land cover shifts (Bonan, 2008; Buma and Livneh, 2015).

Climate impacts studies from the Southwestern USA highlight the strong influence of changing temperature and precipitation on forest distributions (Dale et al., 2001), forest mortality (Allen et al., 2015), and streamflow (McCabe and Wolock, 2007; Nash and Gleick, 1991). Specifically, the strong interaction between forests and water (Anderson et al., 1976) means that forest disturbances will have a large impact on water volume and the timing of streamflow. Streamflow decreases were previously reported for the Upper CRB at Lee's Ferry using the Variable Infiltration Capacity (VIC) hydrologic model (Liang et al., 1994; Liang et al., 1996) forced with future



climate change scenarios alone and reported decreased future projected streamflow (Christensen and Lettenmaier, 2002). A separate study reported that CRB streamflow is highly sensitive to precipitation and temperature shifts, with large reductions in streamflow estimated for small increases in precipitation and temperature (Nash and Gleick, 1991) resulting from enhanced rates of evapotranspiration.

5 The strong link between forests and water, coupled with the emerging threat to future forest viability and integrity, provides a major impetus to study the combined effects of climate and land cover change on streamflow in the CRB. To this end, we apply a hydrologic model that incorporates projected climate changes as well as forest and land cover changes based on recent research. Unlike many other studies, we have included impacts of regrowth of shrubs into our modelling approach since this is the most likely outcome in the CRB (Rother et al.,  
10 2015). We determine the direction and shift in streamflow under different scenarios of climate and land cover changes, as well as identify the physical alterations occurring within watersheds across scales and forest compositions, to reveal the driving mechanisms behind the streamflow changes. Knowledge of the physical mechanisms and dominant triggers for streamflow alterations is critical because it will allow decision makers to make better-informed assessments of near- and long-term water operations in a water-constrained world with a  
15 changing climate (Zou et al., 2010).

The manuscript is arranged as follows: section 1.1 details information on forest disturbances and the impact of climate change on streamflow, section 2 outlines our study site and data, section 3 details our methodology, and section 4, 5 and 6 provide results, discussion, and conclusion, respectively.

## 2 Climate-driven forest disturbances and climate change impacts on streamflow

20 As fire, pests, and drought modify forests in response to climate change, a number of important energy and water fluxes become altered (Adams et al., 2012). Land covers with a low canopy profile and a small crown cover, e.g. shrubs or bare ground, partition water and energy in different manners from forests that have a large crown cover. For example, shrubs retain snow pack later into the melt season thus snow is able to reach the ground and accumulate (Boon, 2007, 2009). This is largely due to reduced canopy cover which results in reduced  
25 interception and sublimation/transpiration (Pomeroy et al., 1998) from this above ground storage reservoir, as well as higher solar radiation and wind speeds through the open areas (Harpold et al., 2014). Snow also melts off more quickly in response to increased shortwave radiation and turbulent heat transfers linked to negative longwave radiative fluxes (Burles and Boon, 2011). The responses have also been linked to decreased albedo due to higher litter and/or darker soils associated with dying trees (Bewley et al., 2010; Winkler et al., 2010). Other



effects of tree mortality on the water balance include changes to soil moisture states, changes in groundwater recharge, and potential feedbacks to the atmosphere (Bonan, 2008).

Variations in the type and magnitude of forest disturbances can also strengthen or dampen some of these effects (Buma and Livneh, 2015). For example, forest wildfires change soil properties and litter depths, which in turn alter infiltration and runoff processes. In addition, mountain pine beetle infestations change canopy conditions but do not change soil properties and moisture states; and drought may affect the forest canopy and the ability of soils to infiltrate water as trees die and soils desiccate (Adams et al., 2012). Effects may also be impacted by the type of post-disturbance regrowth that occurs, such as increases or a difference in shrubs, grasses, forbs coverage or differences in the composition of tree species that replace the forests. For instance, water yield declines have been associated with changes in species compositions of forests in the southern Appalachian Mountains (Caldwell et al., 2016).

Selection of appropriate timescale is an important aspect to consider in vegetation disturbance studies. For example, research focused on forest disturbance over a single summer season, such as post-fire hydrology in New Mexico (Atchley et al., 2017), shows that surface runoff and recharge both rise following fire. However, these studies do not measure the year-long change in water balance, which is critical for snow-driven systems. Examination of water partitioning in disturbed forests within time frames of less than five years may not adequately resolve effects such as forest succession on the hydrologic regime of study basins (Brown et al., 2005), which may result in a significant overestimation of the impacts of changing forests on water yields over the long term (Pugh and Gordon, 2013). Indeed, most studies that encompass forest disturbance monitoring of greater than five years point to increased evapotranspiration from understory regrowth and an associated decline or mitigating effect of the forest cover removal (Brown et al., 2014; Biederman et al., 2014; Guardiola-Claramonte et al., 2011). Other research has shown that vegetation management, such as cutting or thinning of forests in the face of climate change, could be used to ameliorate the impacts of reduced streamflow in the CRB, noting that these effects would only last for a period of approximately ten years (Zou et al., 2010).

Finally, spatial heterogeneity also plays an important role in terms of changing hydroecology in the wake of disturbances. Studies that focus on plot-scale results, where the disturbed forest is the primary cover type, illustrate different responses than locations where impacted trees are only a component of the overall land cover, including grasses, shrubs and non-impacted tree species (Biederman et al., 2014; Pugh and Gordon, 2013). For example, recent studies examining climate change and extreme wildfire on runoff erosion found that peak streamflow sediment yield will increase with climate change and fire severity due to the lack of spatial heterogeneity in land cover types (Gould et al., 2016).



### 3 Methods

#### 3.1 Study Site

To understand the impact of forest disturbances on streamflow under climate changes at different temporal scales and spatial settings, we implement the VIC model for the San Juan River basin, a sub-basin to the CRB, to simulate streamflow responses to future changes in temperature, precipitation and land cover (Fig. 1). The San Juan is a major headwater basin of the Colorado River, accounting for 15% of streamflow and 22% of the area of the Upper CRB. Spanning four states—the Four Corners—the San Juan watershed is also critical for thermoelectric and hydropower generation, substantial oil and gas development, and extensive irrigated agriculture. The San Juan basin captures the diversity present within the Upper CRB. For instance, high elevation (> 4000 m) Colorado mountain ranges and large, snowmelt driven rivers comprise the upper San Juan basin. The lower San Juan basin, located in New Mexico and Arizona, is flat and semi-arid, with intermittent streams that drain into the main tributary of the San Juan during the summer when they are charged by summer monsoonal rains. The San Juan river eventually drains into the Colorado, just below the town of Bluff, Utah (Fig. 1).

#### 3.2 Hydrologic Model

For this work, we used the Variable Infiltration Capacity (VIC) model version 4.2 (Bohn and Vivoni, 2016; Liang et al., 1994) at 1/16 degree (6 km) spatial resolution. In each grid cell, VIC simulates vertical energy and water dynamics at an hourly time step for a mosaic of land cover tiles underlain by a 3-layer soil column. Sub-grid heterogeneity in infiltration is represented by a statistical distribution (the variable infiltration capacity curve). Surface runoff is generated via saturation excess, while sub-surface runoff is characterized by the non-linear baseflow curve of Francini and Pacciani (1991). VIC v4.2 includes fractional canopy coverage derived from normalized difference vegetation index (NDVI) and a spatially varying monthly climatology of leaf area index (LAI), albedo, and canopy fraction (Bohn and Vivoni, 2016). Historical climate data used to run VIC (daily precipitation, minimum and maximum temperature, and wind speed) were obtained from existing gridded data sets for the United States (Livneh et al., 2015). These daily fields were disaggregated to hourly intervals within the VIC model via algorithms as described in Bohn et al. (2013), which also estimated hourly short- and longwave radiation and humidity. Land cover fractional areas were taken from the average of years 2001-2012 of the Moderate Resolution Imaging Spectroradiometer (MODIS) MCD12Q1 Collection 5 Plant Functional Type (PFT) product of Friedl et al. (2010), using the International Geosphere-Biosphere Program (IGBP) classification. Repeating climatological seasonal cycles of vegetation parameters (LAI, canopy fraction, and albedo) were



derived from the MODIS collection 5 MOD15A2, MCD43A3, and MOD13A1 products (Myneni et al., 2002; Schaaf et al., 2002; Huete et al., 2002) over the period 2000–2012, and aggregated spatially over the MCD12Q1 land cover classes within each 1/16<sup>th</sup> degree grid cell.

Soil physical properties (e.g., bulk density, saturated hydraulic conductivity, quartz content) were derived from global datasets such as the United Nations Food and Agriculture Organization (FAO) Digital Soil Map of the World (FAO, 1998). Vegetation structural parameters, such as stomatal and canopy resistances, were taken from Ducoudré et al. (1993). Several other parameters were calibrated empirically:  $D2$  and  $D3$  (the thicknesses of the 2<sup>nd</sup> and 3<sup>rd</sup> soil layers);  $b_{infiltr}$  (the infiltration capacity curve shape parameter);  $Ds$ ,  $Ws$ , and  $Dsmax$  (non-linear baseflow parameters), and  $\alpha_{snow}$  (the albedo of newly-fallen snow). The model was calibrated using an automatic calibration tool (Yapo et al., 1998) to correct streamflow biases against United States Geological Survey (USGS) naturalized gauged streamflow (2006–2010) for the San Juan River basin at Bluff, UT (Fig. 1, Table 1). Our calibration achieved a Nash Sutcliffe efficiency of 0.90 over the calibration period and a 0.93 over the validation period for the San Juan at Bluff, UT USGS naturalized flow data (Table 1).

### 3.3 Climate and Vegetation Change Scenarios

Our study focused on three vegetation projections and four climate projections for a total of 12 different scenarios (Fig. 2). The four climate projections employed to drive VIC were based on Earth System Model (ESM) simulations from the Coupled Model Intercomparison Project, Version 5 (CMIP5; Taylor et al., 2012) climate data including daily temperature, precipitation, and wind speed, downscaled using the Multivariate Adaptive Constructed Analogue (MACA) approach (Fig. 2a and b, Abatzoglou and Brown, 2012) and again disaggregated to hourly intervals via the algorithms described in Bohn et al. (2013). We selected the four ESMs from CMIP5 because they implemented dynamic vegetation processes: HadGEM2-ES (Collins et al., 2011; Cox, 2001), MIROC-ESM (SIEB-DGVM, Watanabe et al., 2011; Sato et al., 2007), MPI-ESM-LR (JSBACH, Giorgetta et al., 2013; Reick et al., 2013), and IPSL-CM5B-LR (ORCHIDEE, Krinner et al., 2005). We used the representative concentration pathway (RCP) 8.5 emissions scenario, which stipulates strongly increasing emissions by 2100 (Van Vuuren et al., 2011) and corroborates current emissions on par with RCP 8.5 (Le Quéré et al., 2015).

The three vegetation projections used in this study are: (1) *climate-only*, which assumes static land cover (i.e., vegetation types do not change), (2) *dynamic*, which uses the dynamic vegetation changes present within the four CMIP5 ESMs, and (3) *disturbed*, which uses disturbance projections based on empirically-based statistical



estimates of forest mortality in the US Southwest (Fig 2c, McDowell et al., 2016). Vegetation classes were aggregated to six dominant cover types from 16 classes in the IGBP vegetation classification and from 9–13 classes in the ESMs (Table 2). Vegetation changes observed in the *dynamic* and *disturbed* scenarios were applied to the historical MODIS vegetation to alter forest coverage for future runs using a simple delta-change approach.

5 Namely, for both the *dynamic* and *disturbed* scenarios, historical forest cover fractions were reduced and concordantly replaced by shrubs in increments of ~10 years (2006–2010, 2010–2020, and so forth); we then ran the projections in 10-year segments, with each segment having a new (constant) forest fraction and starting with the state from the previous time period. Forest cover was reduced by approximately 90% by 2100 for the *disturbed* scenario (McDowell and Allen, 2015; McDowell et al., 2015; McDowell et al., 2016) Fig. 2 c, d).

10 We ran the 12 different scenarios for 1950–2099. We analysed daily, seasonal, and annual streamflow as well as monthly statistics of temperature, precipitation, evapotranspiration, and snow water equivalent (SWE) to understand changes in the water balance in the San Juan River basin. In addition, we investigated the aridity effect upon water availability under forest disturbances using a one-cell analysis. Some studies have suggested an aridity effect, whereby basins with less than 500 mm annual precipitation will see streamflow decrease and vice-  
15 versa (Guardiola-Claramonte et al., 2011; Adams et al., 2012) although this finding is not supported in all work (Brown et al., 2010; Caldwell et al., 2016). For a single cell, we considered climate change by adding +3°C (warm) and +6°C (hot) to the temperature time series and changing precipitation by 20% or -20%. We then changed vegetation characteristics in the single cell to simulate changes in LAI or fractional vegetation spacing (canopy spacing).

## 20 4 Results

### 4.1 Changing climate and land cover

Temperature and precipitation changes are consistent with previous modelling efforts for the region CRB (Gangopadhyay et al., 2011). The four ESMs projected consistent increases in temperatures (4.3°C–7.1°C, mean change of 5.7°C), but variable changes in precipitation (both increases and decreases, -7.2%–7.4%, mean change  
25 of 1.6%) for the San Juan River basin (Fig. 2a, b), by comparing the last 30 years of this century to the last 30 years of the previous century. For both temperature and precipitation, changes are most dramatic and variable after the 2050s concurrent with increasing greenhouse gas emissions. These four ESMs represent the range of warm/hot and wet/dry changes for the San Juan River basin by the 2080s in CMIP5 (Taylor et al., 2012; Brekke et al., 2013). Seasonal variations in temperature and precipitation change indicate important regional process



shifts. For instance, summer and winter differences show that the summer is warming slightly more than the winter (5.8°C compared to 5.6°C). Annual average fall and winter precipitation is projected to increase while spring and summer precipitation is projected to decrease slightly (-1%) with a large range in variability across the basin. We note that the signals of change for both temperature and precipitation differ from the results offered by the Bureau of Reclamation data sets at 1/8<sup>th</sup> of a degree and downscaled using a slightly different technique (Bureau of Reclamation, 2011).

Our study identifies a key challenge in representing land cover: land cover change is represented differently by each of the four ESMs due to the variable representations of land use and the application of different dynamic models within each ESM (Arora, 2002). The *dynamic* trajectory of change is variable depending on the ESM considered (Fig. 2d). The MIROC-ESM model projects increasing forest cover, while the HadGEM2-ES model projects the most amount of change in terms of forest loss. Both IPSL-CM5B-LR and MRI-ESM-LR show only a small amount of change in terms of land cover shifts by the 2080s. The *disturbed* scenario reflects regional changes expected in the Southwestern USA (McDowell et al. 2016) and therefore projects a more severe, and likely realistic, scenario in terms of forest cover change in the San Juan River basin.

#### 4.2 Changing streamflow and water balances

Annually, simulated streamflow in the San Juan River basin under the *climate-only* scenario exhibits differences of -15% to 45%, while the *disturbed* scenario indicates a shift of -21% to 34% for the 2080s as compared with historical streamflow, dependent on the ESM (Fig 3). The *dynamic* scenario changes the streamflow by -16% to 50%, but is generally very similar to the *climate-only* scenario. An exception to this is the MIROC-ESM *dynamic* scenario, which projects an increase in streamflow during winter and a decrease in summer peak flow in response to increasing forest cover and thus decreasing shrubs represented in this model (Fig. 2d). Due to the lack of a large distinction in the vegetation changes under the *dynamic* scenarios and resultant similarities between the *climate-only* and *dynamic* scenarios, we focus on the differences between the *climate-only* and the *disturbed* scenarios for the remainder of this paper.

Seasonal streamflow in the *climate-only* scenario versus historical simulated streamflow illustrates a shift in the timing of peak flow and increased winter streamflow in the San Juan River basin (Fig 3). Peak streamflow occurs approximately one month earlier, owing to earlier snow melt due to a temperature increase. This shift means that winter flows are higher and also indicates more mid-winter warming events. Under *disturbed* forest cover conditions, seasonal streamflow shows a different hydrograph that represents a shift in timing of winter streamflow and a change in the magnitude of both low (Dec–Jan) and high (Apr–May) streamflow compared to



the *climate-only* scenario (Fig. 3). The VIC simulations driven by *disturbed* scenarios project a lower late fall and winter streamflow, with a delay in spring melt and subsequent increase in the pulse of peak streamflow during April–May. Recessional streamflow (May–Jul) is also slightly higher in the *disturbed scenario* than the *climate-only* scenario, resulting in greater water availability in summer (Fig. 3).

5 The mechanisms causing these differences in streamflow response to climate change and forest disturbance are illustrated in Figure 4. In the *climate-only* scenario, the streamflow response is dominated by the impact of temperature on snow pack. Warmer winter temperatures and reduced March snowfall (Fig. 4a) lead to a reduction in snow water equivalent, snow melt, and sublimation from both the pack and the canopy (Fig. 4b-e). Warmer temperatures, earlier snow melt, and greater April rainfall subsequently lead to increases in soil evaporation (Fig. 10 4f) and transpiration (Fig. 4g) in the spring. Warmer temperatures also lead to increased canopy evaporation in late summer (Fig. 4h). However, the higher rates of soil evaporation and transpiration in the spring deplete the middle layer soil moisture (Fig. 4i), which diminishes their rates in the summer (Fig. 4f, g). In the fall, higher rainfall and warmer temperatures lead to greater soil evaporation but only minimal replenishment of soil moisture. Both bottom layer soil moisture (Fig. 4j) and total (surface and subsurface) runoff (Fig. 4k) exhibit 15 earlier and smaller peaks in the spring, reflecting the earlier melting of the reduced snow pack, and lower levels in summer, due to greater ET in the spring. Meanwhile, LAI values are similar between shrubs and forests (Fig. 4l), indicating that the water and energy balance differences are due mainly to snow processes shifts between those two cover types.

Two factors in the *disturbed* scenario further impact the snow pack, partially compensating for the effects of 20 climate change. First, replacement of forest with shrubs leads to an increase (relative to the *climate-only* scenario) in the on-the-ground snow pack accumulation (Fig. 4a), a prominent feature observed in disturbed forests across North America (Boon, 2007; Zou et al., 2010; Biederman et al., 2015; Brown et al., 2014; Harpold et al., 2014). This disturbance-driven increase in on-the-ground snow pack partially compensates for the climate-induced streamflow increase during this time of the year. In addition, the larger on-the-ground snow pack and 25 smaller canopy snow pack are accompanied by proportionally higher and lower rates of sublimation from the ground and canopy snow packs, respectively, relative to *climate-only* (Fig. 4c, d). Second, the increase in snow on the ground caused by the higher shrub coverage leads to higher rates of snow melt (approximately 30%, Fig 4c), releasing water during the late spring and early summer (Apr–June, Fig 4k). The larger snow melt flux leads to substantial increases in transpiration in spring and early summer, not only relative to the *climate-only* scenario, 30 but also relative to historical conditions (Fig. 4f). A similar increase in soil evaporation (Fig. 4f) does not occur due to the upper and middle soil layers already being saturated in the spring in the *climate-only* case (Fig. 4h).



The high shrub transpiration rate in the *disturbed* scenario delays the larger snow melt flux in reaching the bottom soil layer (Fig. 4i), leading to a delayed peak in runoff (Fig. 4j, k).

Differences in streamflow among *disturbed* and *climate-only* scenarios results are most notable at fine scales (100 km<sup>2</sup> to 3,000 km<sup>2</sup>, 72 grid cells in the San Juan) where the forest cover dominates (more than 50%) the land cover (Fig. 5). As basin size increases (smaller circles in Fig. 5), and forest cover becomes sparser with respect to other types of land cover (e.g. mixed forest, deciduous, shrubs and grass covers), the differences between the *climate-only* scenarios and the *disturbed* scenarios begin to decrease, which is corroborated by several other studies (Zhang et al., 2014; Anderegg et al., 2015). The difference between the two scenarios for the densest forest and smallest portion of the basin is approximately 20%, as observed for the larger circles in Fig. 5. This dynamic occurs even under changing precipitation and temperature projections that would otherwise cause increasing or decreasing streamflow (observed in Fig. 5 colour ramps that change from red to blue for precipitation and different symbols for different temperature ranges). Even as the ESMs project increasing precipitation and temperature, we see the variability in responses through the cascade of scale and land cover variability. This finding is consistent with other studies that observed that variability (forest cover composition and topography) in the area and size of forest-shrub conversion can buffer responses of streamflow or ET shifts from climate change (Winkler et al., 2014; Harpold et al., 2014; Caldwell et al., 2016).

The dynamic between temperature, precipitation, and vegetation changes is examined in more detail by using a simple single cell analysis where climate and vegetation is altered in a sensitivity framework. Fig. 6 illustrates the single cell changes in terms of runoff from the grid cell to investigate the aridity effect. We can see that forests and shrubs act similarly in hot/warm and dry environments while the differences are more pronounced between forests and shrubs in wetter environments (Fig. 6a). For the vegetation changes, we see that changing forest structure (e.g., fraction of the canopy that may occur as a result of pest outbreaks) results in small shifts for forests but very large shifts for shrubs (Fig. 6b). For example, shrubs with 50% canopy spacing produce more water than the forests for the majority of runoff conditions (below the 3<sup>rd</sup> quantile, Fig 6b). On the other hand, under high runoff conditions (above the 3<sup>rd</sup> quantile, Fig 6b) shrubs act similarly to regularly spaced shrubs. These changes are on the same order as changes within the climate, generally speaking. Therefore, we hypothesize that larger changes in hydrology under disturbances are more likely to occur not from the forest disturbance itself, but the secondary effects such as regrowth of shrubs, the type of regrowth and the pattern of that regrowth on the landscape. This is why timescale is such an important consideration in studies of this nature, as regrowth patterns are key to understanding how water is partitioned across a disturbed landscape.



## 5 Discussion

Climate change in the CRB is anticipated to cause large impacts to water resources sustainability (Christensen and Lettenmaier, 2007; Rasmussen et al., 2014; Dawadi and Ahmad, 2012). However, to our knowledge few modelling studies have considered the impacts of climate change coupled with changes in vegetation (Buma and Livneh, 2015). In this study, we incorporated changes from the CMIP5 dynamic vegetation models and from an estimate of forest mortality (McDowell et al. 2016) to consider the impacts of both climate and vegetation changes on the water balance in a headwater system to the CRB, the San Juan River basin. We found that failing to consider climate change coupled with vegetation disturbances could result in a ~10% over-estimation of the annual water availability for this basin. For a river system such as the Colorado that is already gravely stressed, 10% less water in the system may lead to significant water management challenges. And, considering seasonality in flows, the changes we illustrated in our monthly and seasonal scenarios indicate that less water will runoff during spring with more water arriving at peak melt. This could lead to water shortages and flooding and lead to planning issues for short term water delivery.

In this work, we considered not only the impacts of changes to streamflow but also the reasons why streamflow is changing. As in other studies, we found that at the size of the basin and the land cover variability can obscure the signal of change (Biederman et al., 2015). When we consider forested regions only, we are able to understand how and why streamflow is projected to change under the disturbed conditions. The main mechanism that is shifting streamflow is the manner in which shrubs impact the water balance during the cold and warm seasons. Snow pack is retained further into the melt season and when snow starts to melt, it melts more quickly and results in higher peak flows. These peak flows, however, occur during a time that the shrubs are using the water, resulting in large transpiration losses. We found differences in canopy evaporation, soil evaporation, and sublimation, but these differences are quite small overall when comparing the overall volume of water held back in the snowpack (SWE) and leaving via shrub transpiration. Overall, the differences in the mechanisms and delivery of the water changed, and this results in a 20% reduction in the amount of water that is available for runoff through the year, compared to the historical streamflow conditions for the forested regions of the San Juan River basin.

We also found that climate and disturbance have opposing influences on snow pack. In the early season, the forest disturbance partially compensates for streamflow impacts caused by warmer temperatures (Fig. 3). We see that the shift towards an earlier snowmelt is compensated in part by the fact that shrubs held on to snow for longer into the snow melt season, releasing snow at a later date. This effect results in overall less water in the



system and peak flows that occur on the same point in the year, but are higher in magnitude which results in lower soil moisture, such that the effect of having a late melting pack may not be beneficial for water resources. However, the snow-on-the-ground could have important consequences for early season energy balances.

The findings presented herein represent a deviation from the more broadly accepted viewpoint that forest disturbances will lead to reduced evapotranspiration and increased streamflow. Previous findings tend to be based on studies carried out over short time periods (first 1–2 year responses, < 5 year studies), pair-basin analyses in watersheds disturbed by clear cuts, and where climate variability may have obscured results (Brown et al., 2005). Studies pointing towards increased streamflow also broadly found evapotranspiration from the canopy decreased, leading to an increase in runoff. However, work based on observations across scales and encompassing multiple disturbances indicates that the regrowth potential for understory, such as shrubs used in this study, is high and that the regrowth is a major control on water availability and direction of change for evapotranspiration and runoff (Caldwell et al., 2016; Biederman et al., 2014; Brown et al., 2014; Biederman et al., 2015). Moreover, ecologists project that global forest covers are expected to decline and be replaced with species and understory compositions that are more water intensive. It is paramount therefore to treat regrowth correctly within models to align these two currently disparate principles and account accurately for changing streamflow. This is a fundamental issue because ESMs rely upon the research principles developed at plot-scale and watershed-scale observational studies and modelling work.

We also investigated the aridity effect upon water availability under forest disturbances. By comparing vegetation changes to climate shifts in a single cell, we highlighted the impacts of changing temperature and precipitation versus the impacts of changing forest cover properties. We found that the greatest differences in our results for forests and shrubs under climate change occurred under wet conditions, and vice versa for dry conditions. This may be due to the large component of water that is leaving via evaporation in arid environments. As well, we find an important control on water availability in the shrub environments was the canopy spacing between the shrubs that led to changes in water balances and water partitioning within the environment. Thus, we believe that not only climate (i.e. aridity) but also vegetation characteristics (i.e. canopy spacing) may play a fundamental role in the impact of disturbances on water availability.

Our study also points to the challenges associated with incorporating spatially variable estimates of changing vegetation patterns. Although CMIP5 contains information from dynamic vegetation models, we find a large disparity between the results from CMIP5 (*dynamic*) and the current estimates and cutting-edge research on forest cover mortality (*disturbed*). The importance of accurately representing the effects that are observed at



catchment scale and in offline models, such as VIC, cannot be understated. ESMs require the capability to incorporate these changes in a meaningful way that can be validated using our current understanding of changes to forests in the Southwestern USA, and globally. This will enable the science community to accurately capture the range of responses and the impacts of changing climate and changing land cover on water resources. ESMs  
5 such as the Department of Energy's ACME model have started down this path with the incorporation and testing of the FATES dynamic vegetation model (Fisher R. et al., 2015).

The differences in future streamflow projections in comparison to historical conditions we observed in this study are notable, and suggest that streamflow will decline by a minimum of ~10% for headwater systems of the Colorado. However, in the San Juan River basin forest cover accounts for only 17% of the area, and only 4% of  
10 these cells have greater than 80% forest cover. Impacts may be much larger for adjacent systems, such as the Gunnison. Future work will investigate the impacts of changing forest cover in the entire Colorado system, using sensitivity analysis to understand the cumulative effect of changing climate and vegetation cover in the Colorado.

## 6 Conclusion

The implications of declining streamflow due to forest disturbances in the San Juan River basin are significant  
15 for the CRB. Given that the San Juan basin is one of the least forested portions of the CRB and assuming similar rates of forest decline will occur nearby (Van Mantgem et al., 2009), other regions of the CRB are likely to experience streamflow declines of much higher severity due to changes in forest cover. Failure to adequately understand the direction and magnitude of changes as we do herein could have catastrophic consequences for those who rely on this resource. The basin—roughly 11% area of the continental United States—directly supports  
20 water supply for more than 30 million people, accounts for approximately 15% of the US's crops and livestock (Cohen et al., 2013), and 53 GW of power generation capacity (Buono and Eckstein, 2014; Cohen et al., 2013). Pressing work includes improvements in spatial representation of changing forest covers across the CRB, further investigations into drivers of change including an understanding of the role of aridity and variability, a stronger link between ecological dynamics and hydrological knowledge that is translated into models, and upward  
25 propagation of these findings into global and earth system models.

## 7 Acknowledgements

We acknowledge the support of John Abatzoglou and Katherine Hedgwick for providing us additional models not in the original MACA data set. We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the climate modelling groups for producing



and making available their model output. For CMIP, the U.S. Department of Energy's Program for Climate Model Diagnosis and Intercomparison provides coordinating support and led development of software infrastructure in partnership with the Global Organization for Earth System Science Portals. Finally, Bennett, Solander, Middleton, McDowell and Xu acknowledge the Los Alamos National Lab's LDRD program for supporting this work. Bohn was supported by Grant 1216037 from the U.S. National Science Foundation (NSF) Science, Engineering and Education for Sustainability (SEES) Post-Doctoral Fellowship program and NSF Grant 1462086, DMUU: Decision Centre for a Desert City III: Transformational Solutions for Urban Water Sustainability Transitions in the Colorado River Basin.



## 8 References

- Abatzoglou, J. T., and Brown, T. J.: A comparison of statistical downscaling methods suited for wildfire applications, *International Journal of Climatology*, 32, 772-780, 2012.
- 5 Adams, H. D., Luce, C. H., Breshears, D. D., Allen, C. D., Weiler, M., Hale, V. C., Smith, A., and Huxman, T. E.: Ecohydrological consequences of drought-and infestation-triggered tree die-off: insights and hypotheses, *Ecohydrology*, 5, 145-159, 2012.
- Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., Rigling, A., Breshears, D. D., and Hogg, E. T.: A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests, *Forest ecology and management*, 259, 660-684, 2010.
- 10 Allen, C. D., Breshears, D. D., and McDowell, N. G.: On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene, *Ecosphere*, 6, 1-55, 2015.
- Anderegg, W. R. L., Kane, J. M., and Anderegg, L. D. L.: Consequences of widespread tree mortality triggered by drought and temperature stress, *Nature Clim. Change*, 3, 30-36, 2013.
- 15 Anderegg, W. R. L., Hicke, J. A., Fisher, R. A., Allen, C. D., Aukema, J., Bentz, B., Hood, S., Lichstein, J. W., Macalady, A. K., McDowell, N., Pan, Y., Raffa, K., Sala, A., Shaw, J. D., Stephenson, N. L., Tague, C., and Zeppel, M.: Tree mortality from drought, insects, and their interactions in a changing climate, *New Phytologist*, 208, 674-683, 10.1111/nph.13477, 2015.
- Anderson, H. W., Hoover, M. D., and Reinhart, K. G.: Forests and water: effects of forest management on floods, sedimentation, and water supply. General Technical Report PSW-018., USDA Forest Service Berkeley CA, 115, 1976.
- 20 Arora, V.: Modeling vegetation as a dynamic component in soil-vegetation-atmosphere transfer schemes and hydrological models, *Reviews of Geophysics*, 40, 2002.
- Atchley, A. L., Kinoshita, A. M., Lopez, S., Trader, L., and Middleton, R. S.: Simulating surface and subsurface water balance changes due to fire severity at the hillslope scale, *Journal of Hydrology* (in review), 2017.
- 25 Bennett, K. E., Cannon, A. J., and Hinzman, L.: Historical trends and extremes in boreal Alaska river basins, *J of Hydro*, 527, 590-607, 2015.
- Bewley, D., Alila, Y., and Varhola, A.: Variability of snow water equivalent and snow energetics across a large catchment subject to Mountain Pine Beetle infestation and rapid salvage logging, *Journal of Hydrology*, 388, 464-479, 2010.
- 30 Biederman, J., Harpold, A., Gochis, D., Ewers, B., Reed, D., Papuga, S., and Brooks, P.: Increased evaporation following widespread tree mortality limits streamflow response, *Water Resources Research*, 50, 5395-5409, 2014.
- Biederman, J. A., Somor, A. J., Harpold, A. A., Gutmann, E. D., Breshears, D. D., Troch, P. A., Gochis, D. J., Scott, R. L., Meddens, A. J., and Brooks, P. D.: Recent tree die-off has little effect on streamflow in contrast to expected increases from historical studies, *Water Resources Research*, 51, 9775-9789, 2015.



- Bohn, T. J., Livneh, B., Oyster, J. W., Running, S. W., Nijssen, B., and Lettenmaier, D. P.: Global evaluation of MTCLIM and related algorithms for forcing of ecological and hydrological models, *Agricultural and forest meteorology*, 176, 38-49, 2013.
- 5 Bohn, T. J., and Vivoni, E. R.: Process-based characterization of evapotranspiration sources over the North American monsoon region, *Water Resources Research*, 52, 358–384, doi:10.1002/2015WR017934., 2016.
- Bonan, G. B.: Forests and Climate Change: Forcings, Feedbacks, and the Climate Benefits of Forests, *Science*, 320, 1444-1449, 10.1126/science.1155121, 2008.
- Boon, S.: Snow accumulation and ablation in a beetle-killed pine stand in Northern Interior British Columbia, *Journal of Ecosystems and Management*, 8, 2007.
- 10 Boon, S.: Snow ablation energy balance in a dead forest stand, *Hydrological Processes*, 23, 2600, 2009.
- Brekke, L., Thrasher, B. L., Maurer, E. P., and Pruitt, T.: Downscaled CMIP3 and CMIP5 Climate and Hydrology Projections: Release of Downscaled CMIP5 Climate Projections, Comparison with preceding Information, and Summary of User Needs, U.S. Department of the Interior, Bureau of Reclamation, Technical Services Center, Denver, Colorado, 47, 2013.
- 15 Brown, A. E., Zhang, L., McMahon, T. A., Western, A. W., and Vertessy, R. A.: A review of paired catchment studies for determining changes in water yield resulting from alterations in vegetation, *Journal of hydrology*, 310, 28-61, 2005.
- Brown, M. G., Black, T. A., Nestic, Z., Foord, V. N., Spittlehouse, D. L., Fredeen, A. L., Bowler, R., Grant, N. J., Burton, P. J., and Trofymow, J.: Evapotranspiration and canopy characteristics of two lodgepole pine stands following mountain pine beetle attack, *Hydrological Processes*, 28, 3326-3340, 2014.
- 20 Brown, R., Derksen, C., and Wang, L.: A multi-data set analysis of variability and change in Arctic spring snow cover extent, 1967–2008, *Journal of Geophysical Research: Atmospheres*, 115, D16111, 10.1029/2010jd013975, 2010.
- Buma, B., and Livneh, B.: Potential Effects of Forest Disturbances and Management on Water Resources in a Warmer Climate, *Forest Science*, 61, 895-903, 2015.
- Buono, R. M., and Eckstein, G.: Minute 319: a cooperative approach to Mexico–US hydro-relations on the Colorado River, 25 *Water International*, 39, 263-276, 2014.
- Burles, K., and Boon, S.: Snowmelt energy balance in a burned forest plot, Crowsnest Pass, Alberta, Canada, *Hydrological Processes*, 25, 3012-3029, 10.1002/hyp.8067, 2011.
- Caldwell, P. V., Miniati, C. F., Elliott, K. J., Swank, W. T., Brantley, S. T., and Laseter, S. H.: Declining water yield from forested mountain watersheds in response to climate change and forest mesophication, *Global change biology*, 2016.
- 30 Christensen, N. S., and Lettenmaier, D. P.: Effects of climate change on the hydrology and water resource of the Colorado River basin, Department of Civil and Environmental Engineering, University of Washington, Seattle, WA, 2002.
- Christensen, N. S., and Lettenmaier, D. P.: A multimodel ensemble approach to assessment of climate change impacts on the hydrology and water resources of the Colorado River Basin, *Hydrology and Earth System Sciences*, 11, 1417-1434, 2007.
- 35 Cohen, M., Christian-Smith, J., and Berggren, J.: Water to supply the land : Irrigated agriculture in the Colorado River basin, Pacific Institute. Oakland, CA, 2013.



- Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C., Sitch, S., Totterdell, I., Wiltshire, A., and Woodward, S.: Development and evaluation of an Earth-System model – HadGEM2, *Geosci. Model Dev.*, 4, 1051-1075, 10.5194/gmd-4-1051-2011, 2011.
- 5 Cox, P. M.: Description of the TRIFFID dynamic global vegetation model, Technical Note 24, Hadley Centre, United Kingdom Meteorological Office, Bracknell, UK, 2001.
- Dale, V. H., Joyce, L. A., McNulty, S., Neilson, R. P., Ayres, M. P., Flannigan, M. D., Hanson, P. J., Irland, L. C., Lugo, A. E., Peterson, C. J., Simberloff, D., Swanson, F. J., Stocks, B. J., and Wotton, B. M.: Climate change and forest disturbances, *BioScience*, 51, 723-734, 2001.
- 10 Dawadi, S., and Ahmad, S.: Changing climatic conditions in the Colorado River Basin: Implications for water resources management, *Journal of Hydrology*, 430–431, 127-141, <http://dx.doi.org/10.1016/j.jhydrol.2012.02.010>, 2012.
- Ducoudré, N. I., Laval, K., and Perrier, A.: SECHIBA, a new set of parameterizations of the hydrologic exchanges at the land-atmosphere interface within the LMD atmospheric general circulation model, *Journal of Climate*, 6, 248-273, 1993.
- 15 FAO: United Nations Food and Agriculture Organization. Digital Soil Map of the World and Derived Soil Properties, Land Water. Digital Media Ser. 1 [CD-ROM], Rome., in, 1998.
- Fisher R., Muszala S., Spessa A., Versteinstein M., Lawrence, P., Xu C., McDowell N., Kluzek E., Koven C., Knox R., Holm J., Rogers B.M., Lawrence D., and G., B.: Taking off the training wheels: the properties of a dynamic vegetation model without climate envelopes., *Geoscientific Model Development*, 8, 3593-3619, 2015.
- 20 Franchini, M., and Pacciani, M.: Comparative analysis of several conceptual rainfall-runoff models, *Journal of Hydrology*, 122, 161-219, 1991.
- Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., and Huang, X.: MODIS Collection 5 global land cover: Algorithm refinements and characterization of new datasets, *Remote Sensing of Environment*, 114, 168-182, 2010.
- 25 Gangopadhyay, S., Pruitt, T., Brekke, L., and Raff, D.: Hydrologic projections for the western United States, *Eos, Transactions American Geophysical Union*, 92, 441-442, 2011.
- Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., and Fieg, K.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, *Journal of Advances in Modeling Earth Systems*, 5, 572-597, 2013.
- 30 Gould, G. K., Liu, M., Barber, M. E., Cherkauer, K. A., Robichaud, P. R., and Adam, J. C.: The effects of climate change and extreme wildfire events on runoff erosion over a mountain watershed, *Journal of Hydrology*, 536, 74-91, 2016.
- Guardiola-Claramonte, M., Troch, P. A., Breshears, D. D., Huxman, T. E., Switanek, M. B., Durcik, M., and Cobb, N. S.: Decreased streamflow in semi-arid basins following drought-induced tree die-off: a counter-intuitive and indirect climate impact on hydrology, *Journal of Hydrology*, 406, 225-233, 2011.
- 35 Harpold, A. A., Biederman, J. A., Condon, K., Merino, M., Korgaonkar, Y., Nan, T., Sloat, L. L., Ross, M., and Brooks, P. D.: Changes in snow accumulation and ablation following the Las Conchas Forest Fire, New Mexico, USA, *Ecology*, 7, 440-452, 10.1002/eco.1363, 2014.



- Huete, A., Didan, K., Miura, T., Rodriguez, E. P., Gao, X., and Ferreira, L. G.: Overview of the radiometric and biophysical performance of the MODIS vegetation indices, *Remote sensing of environment*, 83, 195-213, 2002.
- Kang, S., Kimball, J. S., and Running, S. W.: Simulating effects of fire disturbance and climate change on boreal forest productivity and evapotranspiration, *Science of The Total Environment*, 362, 85-102, 2006.
- 5 <http://dx.doi.org/10.1016/j.scitotenv.2005.11.014>, 2006.
- Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S., and Prentice, I. C.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, *Global Biogeochemical Cycles*, 19, 2005.
- 10 Le Quéré, C., Moriarty, R., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Friedlingstein, P., Peters, G. P., Andres, R. J., and Boden, T.: Global carbon budget 2015, *Earth System Science Data*, 7, 349-396, 2015.
- Liang, X., Lettenmaier, D. P., Wood, E. F., and Burges, S. J.: A simple hydrologically based model of land surface water and energy fluxes for general circulation models, *Journal of Geophysical Research*, 99, 14,415-414,428, 1994.
- Liang, X., Wood, E. F., and Lettenmaier, D. P.: Surface soil moisture parameterization of the VIC-2L model: Evaluation and modification, *Global and Planetary Change*, 13, 195-206, 1996.
- 15 Livneh, B., Bohn, T. J., Pierce, D. W., Munoz-Arriola, F., Nijssen, B., Vose, R., Cayan, D. R., and Brekke, L.: A spatially comprehensive, hydrometeorological data set for Mexico, the U.S., and Southern Canada 1950–2013, *Scientific Data*, 2, 150042, 10.1038/sdata.2015.42, 2015.
- McCabe, G. J., and Wolock, D. M.: Warming may create substantial water supply shortages in the Colorado River basin, *Geophysical Research Letters*, 34, 2007.
- 20 McDowell, N. G., and Allen, C. D.: Darcy's law predicts widespread forest mortality under climate warming, *Nature Climate Change*, 5, 669-672, 2015.
- McDowell, N. G., Williams, A., Xu, C., Pockman, W., Dickman, L., Sevanto, S., Pangle, R., Limousin, J., Plaut, J., and Mackay, D.: Multi-scale predictions of massive conifer mortality due to chronic temperature rise, *Nature Climate Change*, 2015.
- 25 McDowell, N. G., Williams, A. P., Xu, C., Pockman, W. T., Dickman, L. T., Sevanto, S., Pangle, R., Limousin, J., Plaut, J., Mackay, D. S., Ogee, J., Domec, J. C., Allen, C. D., Fisher, R. A., Jiang, X., Muss, J. D., Breshears, D. D., Rauscher, S. A., and Koven, C.: Multi-scale predictions of massive conifer mortality due to chronic temperature rise, *Nature Clim. Change*, 6, 295-300, 10.1038/nclimate2873, 2016.
- 30 Myneni, R. B., Hoffman, S., Knyazikhin, Y., Privette, J. L., Glassy, J., Tian, Y., Wang, Y., Song, X., Zhang, Y., Smith, G. R., Lotsch, A., Friedl, M., Morisette, J. T., Votava, P., Nemani, R. R., and Running, S. W.: Global products of vegetation leaf area and fraction absorbed PAR from year one of MODIS data, *Remote Sensing of Environment*, 83, 214-231, [http://dx.doi.org/10.1016/S0034-4257\(02\)00074-3](http://dx.doi.org/10.1016/S0034-4257(02)00074-3), 2002.
- Nash, L. L., and Gleick, P. H.: Sensitivity of streamflow in the Colorado basin to climatic changes, *Journal of Hydrology*, 125, 221-241, 1991.



- Pomeroy, J. W., Parviainen, J., Hedstrom, N., and Gray, D. M.: Coupled modelling of forest snow interception and sublimation, *Hydrological Processes*, 12, 2317-2337, 10.1002/(SICI)1099-1085(199812)12:15<2317::AID-HYP799>3.0.CO;2-X, 1998.
- 5 Pugh, E., and Gordon, E.: A conceptual model of water yield effects from beetle-induced tree death in snow-dominated lodgepole pine forests, *Hydrological Processes*, 27, 2048-2060, 10.1002/hyp.9312, 2013.
- Rasmussen, R., Ikeda, K., Liu, C., Gochis, D., Clark, M., Dai, A., Gutmann, E., Dudhia, J., Chen, F., and Barlage, M.: Climate change impacts on the water balance of the Colorado Headwaters: High-resolution regional climate model simulations, *Journal of Hydrometeorology*, 15, 1091-1116, 2014.
- 10 Reclamation, B. o.: West-Wide Climate Risk Assessments: Bias-Corrected and Spatially Downscaled Surface Water Projections. Technical Memorandum No. 86-68210-2011-01, U.S. Department of the Interior, Bureau of Reclamation, Technical Services Center, Denver, Colorado, 138, 2011.
- Reick, C., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and anthropogenic land cover change in MPI-ESM, *Journal of Advances in Modeling Earth Systems*, 5, 459-482, 2013.
- 15 Rother, M. T., Veblen, T. T., and Furman, L. G.: A field experiment informs expected patterns of conifer regeneration after disturbance under changing climate conditions, *Canadian Journal of Forest Research*, 45, 1607-1616, 2015.
- Royer, P. D., Cobb, N. S., Clifford, M. J., Huang, C. Y., Breshears, D. D., Adams, H. D., and Villegas, J. C.: Extreme climatic event-triggered overstorey vegetation loss increases understorey solar input regionally: primary and secondary ecological implications, *Journal of Ecology*, 99, 714-723, 2011.
- 20 Sato, H., Itoh, A., and Kohyama, T.: SEIB-DGVM: A new Dynamic Global Vegetation Model using a spatially explicit individual-based approach, *Ecological Modelling*, 200, 279-307, <http://dx.doi.org/10.1016/j.ecolmodel.2006.09.006>, 2007.
- Schaaf, C. B., Gao, F., Strahler, A. H., Lucht, W., Li, X., Tsang, T., Strugnell, N. C., Zhang, X., Jin, Y., and Muller, J.-P.: First operational BRDF, albedo nadir reflectance products from MODIS, *Remote sensing of Environment*, 83, 135-148, 2002.
- 25 Schnorbus, M., Bennett, K. E., and Werner, A. T.: Quantifying the Water Resources Impacts of Mountain Pine Beetle and Associated Salvage Harvest Operations Across a Range of Watershed Scales: Hydrologic Modelling of the Fraser River Basin. Information Report BC-X-423, Canadian Forest Service, Pacific Forestry Centre, Victoria, 64, 2010.
- Solander, K. C., K.E. Bennett, and R.S. Middleton: Historical shifts in streamflow extremes in the Colorado River Basin, *Journal of Hydrology: Regional Studies*, (in review). 2017.
- 30 Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bulletin of the American Meteorological Society*, 93, 485-498, 2012.
- Van Mantgem, P. J., Stephenson, N. L., Byrne, J. C., Daniels, L. D., Franklin, J. F., Fulé, P. Z., Harmon, M. E., Larson, A. J., Smith, J. M., and Taylor, A. H.: Widespread increase of tree mortality rates in the western United States, *Science*, 323, 521-524, 2009.
- 35 Van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., and Lamarque, J.-F.: The representative concentration pathways: an overview, *Climatic change*, 109, 5-31, 2011.



- Varhola, A., Coops, N. C., Bater, C. W., Teti, P., Boon, S., and Weiler, M.: The influence of ground- and lidar-derived forest structure metrics on snow accumulation and ablation in disturbed forests, *Canadian Journal of Forest Research*, 40, 812-821, 2010.
- 5 Watanabe, S., Hajima, T., Sudo, K., Nagashima, T., Takemura, T., Okajima, H., Nozawa, T., Kawase, H., Abe, M., and Yokohata, T.: MIROC-ESM 2010: model description and basic results of CMIP 5-20 c 3 m experiments, *Geoscientific Model Development*, 4, 845-872, 2011.
- Williams, A. P., Allen, C. D., Macalady, A. K., Griffin, D., Woodhouse, C. A., Meko, D. M., Swetnam, T. W., Rauscher, S. A., Seager, R., and Grissino-Mayer, H. D.: Temperature as a potent driver of regional forest drought stress and tree mortality, *Nature climate change*, 3, 292-297, 2013.
- 10 Winkler, R., Boon, S., Zimonick, B., and Baleshta, K.: Assessing the effects of post-pine beetle forest litter on snow albedo, *Hydrological Processes*, 24, 803-812, 2010.
- Winkler, R., Boon, S., Zimonick, B., and Spittlehouse, D.: Snow accumulation and ablation response to changes in forest structure and snow surface albedo after attack by mountain pine beetle, *Hydrological Processes*, 28, 197-209, 2014.
- 15 Yapo, P. O., Gupta, H. V., and Sorooshian, S.: Multi-objective global optimization for hydrologic models, *Journal of Hydrology*, 204, 83 - 97, 1998.
- Zhang, L., Podlasly, C., Ren, Y., Feger, K.-H., Wang, Y., and Schwärzel, K.: Separating the effects of changes in land management and climatic conditions on long-term streamflow trends analyzed for a small catchment in the Loess Plateau region, NW China, *Hydrological Processes*, 28, 1284-1293, 10.1002/hyp.9663, 2014.
- 20 Zou, C. B., Ffolliott, P. F., and Wine, M.: Streamflow responses to vegetation manipulations along a gradient of precipitation in the Colorado River Basin, *Forest Ecology and Management*, 259, 1268-1276, 2010.



1 **Tables**

2 **Table 1. Calibration, validation and period of record statistics for San Juan at Archuleta and San**  
 3 **Juan at Bluff, UT.**

	<b>Objective Function</b>	<b>San Juan at Archuleta</b>	<b>San Juan at Bluff, UT</b>
Calibration (2006-2010)	Nash Sutcliffe	0.91	0.90
	Nash Sutcliffe log	0.78	0.76
	Volume Bias	0.77	0.75
Validation (2001-2005)	Nash Sutcliffe	0.93	0.93
	Nash Sutcliffe log	0.83	0.59
	Volume Bias	0.77	0.43
Entire Period (2001-2010)	Nash Sutcliffe	0.90	0.91
	Nash Sutcliffe log	0.81	0.67
	Volume Bias	0.78	0.58




1 **Table 2. IGBP classification names and codes from MODIS, our new classification, and remapped values (RM). Values for each ESM follows, MIROC-ESM, MPI-ESM-**  
 2 **LR, HadGEM2-ES, and IPSL-CM5A-LR. For each ESM, the original code and the remapped (RM) values are given.**

IGBP Classification	Code	Our Class	RM	MIROC-ESM	Code	RM	MPI-ESM-LR	Code	RM	HadGEM2-ES	Code	RM	IPSL-CM5B-LR	Code	RM
Evergreen Needleleaf forest	1	Evergreen forests	1	Tropical Forests	1	1	glacier	1	nc	broadleaf trees	1	2	Bare soil	1	6
Evergreen Broadleaf forest	2	Evergreen forests	1	Temperate/boreal evergreen	2/4	1	Tropical broadleaf evergreen	2	1	needleleaf trees	2	1	Tropical broad-leaved evergreen	2	1
Deciduous Needleleaf forest	3	Deciduous Forests	2	Temperate/boreal deciduous	3/5	2	Tropical broadleaf deciduous	3	2	C3/C4 grass	3/4	3	Tropical broad-leaved raingreen	3	2
Deciduous Broadleaf forest	4	Deciduous Forests	2	C3/C4 grass	6/7	3	Extra-tropical evergreen	4	1	shrubs	5	4	Temperate/boreal needleleaf evergreen	4/7	1
Mixed forest	5	Evergreen forests	1	Crop	8	5	extra-tropical deciduous	5	2	Urban/inland water/bare/ice	6-9	6	Temperate broad-leaved evergreen	5	1
Closed Shrublands	6	Shrubs	4	Pasture	9	3	raingreen shrubs	6	4				Boreal needleleaf summergreen	9	2
Open Shrublands	7	Shrubs	4	Bare ground/Residual	10/11	6	deciduous shrubs	6	4				Temperate/boreal broad-leaved summergreen	6/8	2
Woody Savannas (40-60% tree cover)	8	Grass	3				C3/C4 grass	8/9	3				C3/C4 grass	10/11	3
Savannas (10-40% tree cover)	9	Grass	3				C3/C4 pasture	10/11	3				C3/C4 agriculture	12/13	5
Grasslands	10	Grass	3				C3/C4 crops	12/13	5						
Permanent wetlands (assuming like grass)	11	Grass	3												
Croplands (corn)	12	Crops	5												
Urban and built-up (assuming sparse grass)	13	Urban/Water/Bare	6												
Cropland/Natural vegetation mosaic	14	Crops	5												
Snow and ice (assuming sparse grass)	15	Urban/Water/Bare	6												
Barren or sparsely vegetated (assuming sparse grass)	16	Urban/Water/Bare	6												




## 1 Figures




2

3 Figure 1. San Juan River basin ( $61,560 \text{ km}^2$ ) in the Colorado River watershed ( $634,150 \text{ km}^2$ , inset), one of the most  
4 important water resources in the western US. The San Juan River basin (outline of HUC8 watersheds shown in red)  
5 spans the Four Corners region (CO, UT, AZ and NM) of the US, and supports multiple energy and water projects.  
6 Naturalized flow gage sites (USBR) at San Juan at Bluff, UT and San Juan at Archuleta, NM are shown with closed  
7 blue triangles.




1

2 Figure 2. Temperature (a), precipitation (b) and land cover (c and d)  
 3 changes to the 2080s (2070-2099) for the four  
 4 ESMS as a difference from the historical climatology (1970-1999), averaged over the ESMS and spatially over the  
 5 San Juan River basin (61,560 km<sup>2</sup> or 1570 grid cells). Vegetation disturbance scenarios based on McDowell et al.  
 6 2016 (c) illustrate declining forest covers and increased shrub covers, with strong declines (>50% forest loss) while  
 7 CMIP5 forest disturbances illustrate more moderate change with HadGEM2-ES projecting the largest decline and  
 MIROC-ESM projecting an increase in forest cover.




1

2 Figure 3. Monthly average future streamflows (2070-2099) compared to historical (1970-1999) for the San Juan  
3 River basin at Bluff, UT. The range of responses for each GCM is represented by the semi-transparent envelope  
4 around the lines. The historical is shown in black (single line, no envelope), the climate scenarios are shown in grey  
5 (solid dark grey line), and the disturbance scenarios are shown in green (dashed dark green line).




1

2 Figure 4. Monthly water balance panel plot for forested (greater than 50%) regions in the San Juan river basin.  
 3 Panels include a) precipitation (mm) and temperature (C) for historical (1970-1999, black) and future (2070-2099,  
 4 grey), snow water equivalent (SWE, b), snow melt (c), sublimation from snow (d), sublimation from canopy (e), soil  
 5 evaporation (f), transpiration (g), canopy evaporation (h), soil moisture from second soil later (L2, i), soil moisture  
 6 from the third soil layer (L3, j), leaf area index (LAI, k), and runoff (l), for historical (black line), climate-only (grey  
 7 line) and disturbed scenarios (green line). April and June are indicated with vertical grey lines.



1  
2 Figure 5. Runoff differences for the *climate-only* scenario (x-axis) plotted against disturbed scenario (y-axis). All  
3 changes are shown as projected differences from the historical (in % change).



1

2 Figure 6. Cumulative density distributions are used here to illustrate the variations in (a) climate and (b) vegetation  
3 parameterizations for a single cell in the San Juan River basin. Climate alterations are shown for a hot-wet, hot-dry,  
4 warm-wet and warm-dry scenarios for both forest and shrubs. For the vegetation parameterization, forests, forests  
5 with two times leaf area index (LAI) and canopy fraction of 0.5, shrubs, and shrubs with fraction of 0.5 and 1.0. A  
6 shrub fraction of 1 indicates that there are no spaces between the plants. The third quantile in the average of the  
7 distributions is indicated in each plot with intersecting lines.