

Data assimilation of GRACE terrestrial water storage estimates into a regional hydrological model of the Rhine River basin

N. Tangdamrongsub¹, S. C. Steele-Dunne², B. C. Gunter^{1,3}, P. G. Ditmar¹, and A. H. Weerts^{4,5}

[1]Department of Geoscience and Remote Sensing, Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, The Netherlands

[2]Department of Water Resources, Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, The Netherlands

[3]School of Aerospace Engineering, Georgia Institute of Technology, Atlanta, The United States of America

[4]Operational Water Management, Deltares, Delft, The Netherlands

[5]Hydrology and Quantitative Water Management Group, Department of Environmental Sciences, Wageningen University, The Netherlands

Correspondence to: N. Tangdamrongsub (n.tangdamrongsub@tudelft.nl)

Abstract

The ability to estimate Terrestrial Water Storage (TWS) realistically is essential for understanding past hydrological events and predicting future changes in the hydrological cycle. Inadequacies in model physics, uncertainty in model land parameters, and uncertainties in meteorological data commonly limit the accuracy of hydrological models in simulating TWS. In an effort to improve model performance, this study investigated the benefits of assimilating TWS estimates derived from the Gravity Recovery And Climate Experiment (GRACE) data into the OpenStreams [wflow_hbv](#) model using an Ensemble Kalman Filter (EnKF) approach. The study area chosen was the Rhine River basin, which has both well-calibrated model parameters and high-quality forcing data that were used for experimentation and comparison. Four different case studies were examined which were designed to evaluate

29 different levels of forcing data quality and resolution including those typical of other less
30 well-monitored river basins. The results were validated using in situ groundwater and stream
31 gauge data. The analysis showed a noticeable improvement in groundwater estimates when
32 GRACE data were assimilated, with a best-case improvement of 71% in correlation
33 coefficient (from 0.31 to 0.53) and 35% in RMS error (from 8.4 to 5.4 cm) compared to the
34 reference (ensemble open-loop) case. The correlation and RMSE improvements in
35 groundwater estimates for the data-sparse case were up to 33% and 35%, respectively, while
36 the average improvements for all four cases evaluated were 13 % and 14%, respectively. Only
37 a slight overall improvement was observed in streamflow estimates when GRACE data were
38 assimilated. Further analysis suggested that this is likely due to sporadic short-term, but
39 sizeable, errors in the forcing data and the lack of sufficient constraints on the soil moisture
40 component. Overall, the results highlight the benefit of assimilating GRACE data into
41 hydrological models, particularly in data-sparse regions, while also providing insight on
42 future refinements of the methodology.

43

44 **1 Introduction**

45 Terrestrial Water Storage (TWS) is the integrated sum of all surface water, soil moisture,
46 snow water, and groundwater availability, and is a metric critical for monitoring the water
47 supply for domestic, industrial, and agricultural sectors. The ability to estimate TWS is useful
48 for understanding past events and predicting future changes in the hydrological cycle,
49 streamflow and water availability, as well as their impact on the occurrence of droughts, heat
50 waves, and floods (Hirschi, et al., 2007). The individual components of TWS influence the
51 climate system in different ways. Soil moisture is a major source of water for the atmosphere
52 in the terrestrial water cycle (Jung et al, 2010) and plays a particularly important role in the
53 climate system (Seneviratne et al., 2010). Soil moisture estimates are also useful for seasonal
54 predictions, and have been shown to improve predictions of air temperature in North America
55 (Koster et al., 2010) and Europe (van den Hurk et al., 2012). Similarly, realistic estimation of
56 the snowpack can improve the prediction of near surface temperature at high latitude regions
57 at 15-30 day scales (Orsolini et al., 2013). Finally, groundwater variability influences soil
58 moisture and evapotranspiration, and is related to long-term water availability and climate
59 changes (Bierkens and van den Hurk, 2007; Green et al., 2011).

60 Despite the importance of having reliable estimates of TWS, knowledge about the spatial and
61 temporal variations of TWS and its components is generally lacking. This is particularly true
62 at large scales, due to the absence of global monitoring systems. Ground-based
63 measurements, while very accurate, only provide point-wise estimates (Dorigo et al., 2011;
64 Lettenmaier and Famiglietti, 2006). Large spatial coverage can be achieved using satellite
65 remote sensing observations, but these often measure only one component of the total storage
66 and suffer from additional limitations. For example, in the case of soil moisture, satellite
67 observations are limited to the top few centimetres of the soil column and to areas free from
68 dense vegetation cover (e.g., de Jeu et al., 2008; Entekhabi et al., 2010; Kerr et al., 2012).
69 Variations in surface water can be observed with satellite altimetry but this technique is
70 currently limited to large target areas (>10 km) (Phan et al., 2012; Schwatke et al., 2013,
71 [Kleinherenbrink et al., 2014](#)).

72 Since measurements alone are not sufficient to comprehensively monitor all components of
73 TWS, hydrological models are often employed. A strong point of hydrological models is their
74 ability to obtain spatially distributed estimates, differentiate TWS components, and simulate
75 changing boundary conditions. Many hydrological models are available, which vary in terms
76 of process description, temporal resolution, spatial resolution, and the detail in process
77 representation (Koster et al., 2000; Rodell et al., 2004). Models vary in terms of which TWS
78 components are included in the model, and how they are represented. The performance of
79 hydrological models is also influenced by the accuracy of the input forcing data and the
80 quality of the model calibration. The existence of model uncertainties motivates the need to
81 combine the model with independent observations to obtain a better representation of the
82 system's behaviour.

83 Changes in TWS can also be estimated by observing variations of the regional gravity field
84 over time. The idea is that changes in water storage, including those deep underground,
85 induce a gravitational signature proportional to the amount of (water) mass redistribution.
86 Since 2002, these variations have been measured by the Gravity Recovery and Climate
87 Experiment (GRACE) satellite mission (Tapley et al., 2004). GRACE allows temporal
88 variations of Earth's gravity field to be observed at spatial scales ranging in the hundreds of
89 kilometres, and at time scales as short as one month. As part of the GRACE data processing,
90 atmospheric and ocean related time-variable gravity effects are removed from the data,
91 leaving the remaining gravity signal over the continents mostly representing changes in TWS

92 (in some areas, additional removal of other nuisance signals is needed, such as those due to
93 glacier melting, glacial isostatic adjustment, and megathrust earthquakes). The GRACE
94 mission has enabled the first direct observations of large-scale TWS, and studies to date have
95 shown high correlation with modelled TWS in terms of seasonal dynamics and regional
96 spatial patterns (Syed et al., 2008; Becker et al., 2011; Longuevergne et al., 2013). A unique
97 feature of satellite gravimetry is that it observes the total column of mass variations (including
98 groundwater) while other remote sensing techniques can only penetrate to a very limited
99 depth, often just a few centimetres. In contrast to hydrological modelling, it is not possible to
100 identify which layer the inferred mass variations can be attributed (Rodell et al., 2009).

101 Several earlier studies have employed data assimilation to combine the strengths of
102 hydrological modelling and GRACE observations and to mitigate their respective weaknesses
103 (Zaitchik et al., 2008; Su et al., 2010; Houborg et al., 2012; Li et al., 2012; Forman et al.,
104 2012). In data assimilation, the model states are constrained by observations, taking into
105 account the estimated uncertainties for both the model states and the observations (Evensen,
106 2003; Reichle, 2008). Employing data assimilation provides a mechanism to downscale the
107 coarse GRACE TWS variations to the temporal and spatial resolution of the model as well as
108 providing insight from the hydrological model into the distribution of TWS between the
109 individual storage terms. Zaitchik et al. (2008) assimilated GRACE into the Catchment Land
110 Surface Model to estimate the TWS over the Mississippi River Basin. Houborg et al. (2012)
111 and Li et al. (2012) applied a similar strategy to improve the drought indicator over North
112 America and Europe, respectively. Su et al. (2010) and Forman et al. (2012), extended the
113 work of Zaitchik et al. (2008) to improve the estimated snow water equivalent over North
114 America and northwestern Canada, respectively. All results from earlier studies reported that
115 assimilating GRACE improved, or at least did not degrade, the hydrology model's
116 performance. In particular, good agreements between estimated state variables, e.g.,
117 groundwater and streamflow, and the in situ measurements were observed. This study adds to
118 these prior works by examining how GRACE assimilation performs when the hydrological
119 model is not well calibrated or when unreliable meteorological data are used to force the
120 model. This focus of the study is on the Rhine River basin (Fig. 1), which is significantly
121 smaller than the large basin or continent scale studies of these prior works, so the analysis
122 presented here provides new insight into the performance of GRACE assimilation over
123 smaller basins. And, while previous data assimilation studies have been performed in the
124 Rhine and neighbouring basins (e.g. Weerts and Serafy, 2006; Rakovec et al., 2012), this

125 study is the first to incorporate GRACE observations in the assimilation scheme for this
126 region.

127 The primary goal of this study was to understand the impact of GRACE assimilation on the
128 estimated TWS, groundwater (GW) variations and streamflow in the Rhine basin. The second
129 goal was to investigate the potential value of assimilating GRACE observations in data-sparse
130 regions. Four scenarios were considered in which the model parameters used were either
131 calibrated (high quality) or basin-averaged (poor quality) values, and the forcing data were
132 obtained from either local (high quality) or global (poorer quality) datasets. In this context,
133 comparison of the four scenarios provide insight into how GRACE can be used to constrain
134 hydrological models when limited data are available.

135

136 **2 Hydrological modelling**

137 The hydrological model employed in this study is the OpenStreams [wflow_hbv](#) model
138 ([Schellekens, 2014](#)). This is a distributed version of the HBV-96 model, named after the
139 Hydrologiska Byråns Vattenbalansavdelning (Hydrological Bureau Waterbalance-section).
140 The HBV model was originally developed at this former section of the Swedish
141 Meteorological and Hydrological Institute (SMHI) in the early 1970's. Since then, the HBV
142 model has been used in over 40 countries. In 1996, a comprehensive re-evaluation of the
143 HBV model routines was carried out ([Lindström et al., 1997](#)), which resulted in the HBV-96
144 version. The OpenStreams [wflow_hbv](#) model is a variant of this model, programmed in the
145 PCRaster-Python environment ([Karszenberg et al., 2009](#)), but using a kinematic wave for
146 hydrological routing. It is publicly available through the OpenStreams project
147 (<https://code.google.com/p/wflow/>, last access 18 January 2015). The defined grid resolution
148 used in this study was 1 km. A schematic representation of OpenStreams [wflow_hbv](#) is given
149 in Fig. 2 (a) and the key parameters of the soil moisture and runoff response routines are listed
150 and described in Table 1.

151 [OpenStreams wflow_hbv](#) consists of three main routines: (i) precipitation and snow, (ii) soil
152 moisture, and (iii) runoff. The water from either precipitation or snow first enters the
153 interception storage and snow routine. The remaining liquid water (from rainfall and snow
154 melt) after the snow routine infiltrates into the soil. The soil moisture storage term (SM in
155 [mm]), which includes both surface and root zone soil moisture is controlled by three main
156 parameters f_c , l_p , and β (see also Table 1). When the amount of water exceeds the maximum

157 capacity (f_c), the excess water becomes available for direct runoff. Within the soil layer,
158 seepage is generated and controlled by an empirical parameter β . The volume of water
159 available for runoff (direct runoff and seepage) is transferred to the runoff response routine.
160 Additionally, some percentage of the soil moisture evaporates, which is controlled by a
161 defined threshold ($f_c \times l_p$).

162 Two linear reservoirs are defined in the runoff routine, namely the upper and lower zones (UZ
163 and LZ). The excess water from SM recharges the upper zone, and some of the water in UZ
164 percolates to LZ, as determined by the perc parameter. At the same time, capillary flow from
165 UZ to SM also occurs, controlled by cflux. The runoff generation in UZ is controlled mainly
166 by two main parameters, the recession constant (khq) and the non-linearity parameter (α). LZ
167 contributes the water to the base flow through the recession constant (k_4). The amount of base
168 flow is simply the multiplication between k_4 and the amount of LZ. Runoff from UZ and LZ
169 then enters the routing model to determine the streamflow.

170 For reference, TWS is defined here as the sum of SM, UZ and LZ. Groundwater storage
171 (GW) is defined as the sum of UZ and LZ. These storage terms are calculated in the soil
172 moisture and runoff response routines. Fig. 2 (b) shows the simulated SM, UZ and LZ from a
173 nominal model run (i.e. using the calibrated parameters and local forcing data). The main
174 source of TWS variation in this model is SM, with the variations in LZ and UZ an order of
175 magnitude smaller. Extraction of groundwater for irrigation is considered to be small over our
176 study region. It accounts for less than $1 \text{ km}^3/\text{year}$. Industry is the largest user (Wada et al.
177 (2014). However, The net removal is small as only 10% of the total water withdrawal over the
178 Rhine is from groundwater and the water is re-introduced to the system after being used for
179 industry. This is markedly different to the extraction of groundwater for irrigated agriculture
180 observed in India (Ferrant et al., 2014). Therefore, this impact on TWS is not considered in
181 this study.

182 The OpenStreams `wflow_hbv` model was calibrated for the Rhine river basin using
183 observations from in situ streamflow gauges (Mülders et al., 1999; Eberle et al., 2002; 2005;
184 Photiadou et al., 2011). The spatial distribution of the calibrated model parameters is shown
185 in Fig. 3.

186 In data-sparse regions, a lack of in situ (meteorological and streamflow) data makes it
187 difficult to calibrate hydrological models (Sivapalan et al., 2003; Hrachowitz et al., 2013).
188 Therefore, we decided to add “non-calibrated” cases to our simulations. In those cases, we

189 defined the non-calibrated parameters as the areally-averaged values of the calibrated
190 parameters in the entire basin, and used these for every grid cell in the basin.

191

192 **3 Datasets**

193 **3.1 GRACE observation**

194 The most recent release (RL05) of the GRACE gravity model product, generated by the
195 University of Texas at Austin's Center of Space Research (CSR: Bettadpur, 2012), was used
196 in the analysis. The CSR RL05 models represent a time-series of Stokes coefficients up to a
197 maximum spherical harmonic degree and order of 60, and are provided monthly. Following
198 the GRACE conventional processing steps, degree-1 coefficients provided by Swenson et al.
199 (2008) were added, and the degree-2 coefficients were replaced by the values estimated from
200 satellite laser ranging (Cheng and Tapley, 2004). Variations in the gravity field were
201 computed by removing the long-term mean (computed over the entire study period, see Sect.
202 5) from each monthly solution. The TWS variations over the Rhine basin were then produced
203 using the approach described by Wahr et al. (1998). Because of strong noise artefacts present
204 in the high degree coefficients, a de-stripping filter similar to that described in Swenson and
205 Wahr (2006) was applied to each monthly solution. The filter used a 5th degree polynomial
206 (Savitsky-Golay) over a 5-point window to remove the correlations, and orders below 8
207 remained unchanged. Further, an additional 250-km radius Gaussian smoothing (Jekeli, 1981)
208 was applied. While this process helps to mitigate noise in the solution, it also attenuates
209 genuine signal, so a scale factor is often applied in an effort to restore some of the signal that
210 gets "leaked" out of the basin due to the spatial filtering. To that end, scale factors using the
211 Global Land Data Assimilation System (GLDAS) hydrological model (Rodell et al., 2004)
212 were computed following the method described by Landerer and Swenson (2012). The sum of
213 four soil moisture layers (0 to 2 m) and a snow water equivalent layer from a monthly
214 GLDAS NOAH Version 1 model was defined as the TWS. We (least squares) fitted the time
215 series between the original and filter GLDAS at every grid node over the Rhine using only
216 one scale factor. The estimated filtering scale factors varied between 0.98 and 1.02 over the
217 Rhine River basin. The correction for glacial isotactic adjustment, which has been shown in
218 other regions to affect the interpretation of long-term trends (Peltier, 2004), was determined to
219 be small in our study, so the corresponding correction was not applied.

220 **3.2 Forcing data**

221 The forcing data required to drive the OpenStreams [wflow_hbv](#) model are precipitation,
222 temperature and potential evapotranspiration (PET). Two types of forcing data were used in
223 this study. “Local” forcing data indicates the best available data, and “global” forcing data
224 indicates a lower quality dataset but one which is available globally or nearly globally.

225 In this study of the Rhine basin, local forcing data refer to meteorological data from the
226 network of local weather stations, providing higher spatial and temporal resolution. Local
227 precipitation and temperature data were retrieved from the European Climate Assessment &
228 Data set (ECA&D) and ENSEMBLE project, known as E-OBS data (Haylock et al., 2008).
229 Data collected from several hundred ground stations were combined to produce a daily grid of
230 precipitation and mean surface temperature at a 0.25-degree spatial resolution. Local PET
231 data were derived from climatological data obtained from the Commission for the Hydrology
232 of the Rhine basin (CHR) and the German Meteorological Service (DWD) (Weerts et al.,
233 2008). The daily local PET was interpolated from a monthly mean value with a fixed annual
234 cycle and was available at a 1-km spatial resolution (Weerts et al., 2008; Photiadou et al.,
235 2011).

236 Global precipitation and temperature data were obtained from Sheffield et al. (2005). These
237 data are constructed based on the long-term near-surface meteorological variables from the
238 National Centers for Environmental Prediction–National Center for Atmospheric Research
239 (NCEP/NCAR) reanalysis product. The daily global precipitation and temperature data were
240 provided at a spatial resolution of 0.5-degree. For global PET, the 1-degree daily product
241 generated by Senay et al. (2008) was used.

242 Fig. 4 shows a comparison between mean daily precipitation, temperature and PET in 2006
243 from the local and global forcing datasets. For the mean temperature, aside from the
244 resolution difference, the spatial distribution and magnitude is very similar between the two
245 datasets. On the other hand, significant differences can be seen between the local and global
246 precipitation data, especially over the High Rhine. Differences are also observed in the PET
247 products, with the global dataset having generally higher values than the local one, in addition
248 to the much coarser spatial resolution of the global product.

249 3.3 Validation data

250 Groundwater and streamflow measurements from various networks are used to validate our
251 estimated results.

252 3.3.1 Groundwater data

253 In situ groundwater measurements were obtained from 3 different networks:

254 1) Ministerium für Klimaschutz, Umwelt, Landwirtschaft, Natur- und Verbraucherschutz
255 des Landes Nordrhein-Westfalen (<http://www.elwasweb.nrw.de>, last access: 5 March
256 2014)

257 2) Bayerisches Landesamt für Umwelt (<http://www.gkd.bayern.de>, last access: 5 March
258 2014)

259 3) Portail national d'Accès aux Données sur les Eaux Souterraines (ADES,
260 <http://www.adeseaufrance.fr>, last access: 17 March 2014)

261 Measurements that did not exhibit seasonal variations were flagged as belonging to confined
262 aquifers, and were excluded. Data from stations with weekly measurements (e.g., ADES)
263 were interpolated to daily intervals. A total of eighteen wells were used for validation. Their
264 locations are shown in Fig. 1, and their names are provided in Table A1.

265 The in situ groundwater measurements were provided in the form of piezometric head. The
266 variations in piezometric head can be related to variations in groundwater storage if the
267 specific yield is known (Rodell et al., 2007). As the latter data were unavailable, the
268 piezometric head was scaled to the units of GW storage based on other GW data. Previous
269 studies have demonstrated that subtracting SM derived from GLDAS from GRACE was able
270 to extract the groundwater component from GRACE in several regions e.g., North America
271 (Rodell et al., 2006; 2007), Australia (Tregoning et al., 2012), the Middle East (Longuevergne
272 et al., 2013), etc. We adopt a similar idea by using the relationship between $\Delta TWS - \Delta SM$
273 (TWS variation from GRACE minus SM variation) and the observed head to scale the
274 observed head. Ideally, we would prefer to use in-situ soil moisture data to represent the SM
275 term, but they are not available at the well locations, and the nearest station from the
276 International Soil Moisture Network (ISMN: Dorigo et al., 2011) does not have data covering
277 the GRACE observation period. The soil moisture estimated from remote sensing was also
278 not appropriate because the penetration depth depends on frequency and would not be the
279 same as that in OpenStreams wflow_hbv. Therefore, we decided to use GLDAS-derived SM

280 in this study. The SM variation from GLDAS (ΔSM_{GLDAS}) was computed by removing its
 281 long-term mean value. The long-term mean value was produced from all GLDAS SM data
 282 over the same period as the GRACE observations (see Sect. 5). The groundwater variations
 283 from GRACE (ΔGW_{GRACE}) were obtained by removing ΔSM_{GLDAS} from the GRACE
 284 observations every month. ΔGW_{GRACE} was interpolated to daily values in order to compare it
 285 to the daily head variations Δh . The comparison was done using the following relationship:

$$286 \quad \Delta GW_{GRACE} + e = a + b \cdot \Delta h \quad (1)$$

287 where e indicates the observation error. The two parameters a and b were estimated by least-
 288 squares regression. The scaled in situ GW variation ($\Delta GW_{in\ situ}$) were then obtained from the
 289 observed variations in piezometric head using:

$$290 \quad \Delta GW_{in-situ} = \hat{a} + \hat{b} \cdot \Delta h \quad (2)$$

291 where \hat{a}, \hat{b} are the parameters estimated from Eq. (1).

292 3.3.2 Streamflow data

293 Streamflow was validated using observations from the thirteen in situ gauges indicated in Fig.
 294 1. Time-series were provided by the Hydrological Modelling Basis in the Rhine Basin
 295 (HYMOG; Bader et al., 2013). The hourly data were aggregated to daily data for this study.

296

297 4 Data assimilation

298 4.1 Ensemble Kalman Filter

299 The Ensemble Kalman Filter (EnKF) is used here to assimilate GRACE TWS into the
 300 OpenStreams `wflow_hbv` model. The EnKF uses a Monte Carlo approach: an ensemble of
 301 model states is integrated forward in time using the forward model. The update equation from
 302 the classical Kalman filter is used to update the model estimate, where the Kalman gain is
 303 determined using the error covariances calculated from the ensemble (Evensen, 1994). The
 304 EnKF and its variants are widely used because they are efficient, easy to implement and allow
 305 great flexibility in terms of model uncertainty (Evensen, 2003). In this study, we implement a
 306 so-called 1D-EnKF (De Lannoy et al., 2009) in which each grid cell is updated individually.
 307 The state equation in discrete form is given as:

$$308 \quad \psi(t + 1) = f(\psi(t), u(t + 1), \alpha, w(t)) \quad (3)$$

309 where f is the model operator, ψ is the state variables, u is the forcings, α is the model
 310 parameters, and w is the model error. In this paper, the state variables (ψ) are an $n \times 1$ vector of
 311 TWS from OpenStreams `wflow_hbv`. The observations available at a measurement time t are
 312 gathered in a vector of observations d (TWS from GRACE):

$$313 \quad d(t) = H\psi(t) + \epsilon; \epsilon \sim \mathcal{N}(0, R) \quad (4)$$

314 where d is an $m \times 1$ vector containing the observations, H is measurement operator which
 315 relates the true state $\psi(t)$ to the measured variables $d(t)$. In this study, the observation and the
 316 state vector are TWS, so $n=m=1$ and H is the unit matrix. The uncertainties in the
 317 observations are given in the random error ϵ , which is assumed to have zero mean and
 318 covariance matrix R . In the initialization phase, the EnKF is initialized by generating an
 319 ensemble (i) of N realizations of the state vector $\psi_i(t)$, $i=1, \dots, N$ around a nominal $\psi(t)$. This
 320 reflects the prior knowledge of the state at the initial time. The EnKF moves sequentially from
 321 one observation time to the next and works in two steps, a forecast step and an update step. At
 322 the updated time t (when the observation is available), an ensemble of perturbed observations,
 323 $d_i(t)$ is generated as:

$$324 \quad d_i(t) = d(t) + \epsilon_i(t), \quad (5)$$

325 where ϵ_i denotes the perturbation of the error of each ensemble member i . If the ensembles of
 326 the variables are stored in a matrix $A = (\psi_1, \psi_2, \psi_3, \dots, \psi_N)$, the ensemble perturbation matrix
 327 can be defined as $A' = A - \bar{A}$ where \bar{A} is the mean computed from all ensemble members.
 328 Similarly, the ensemble members of the observation and perturbations are gathered into the
 329 matrices $D = (d_1, d_2, d_3, \dots, d_N)$ and $\gamma = (\epsilon_1, \epsilon_2, \epsilon_3, \dots, \epsilon_N)$. The update equation can be
 330 expressed as (Evensen, 2003):

$$331 \quad A^a(t) = A(t) + A'(t)A'^T(t)H^T(HA'(t)A'^T(t)H^T + \gamma\gamma^T)^{-1}(D(t) - HA(t)) \quad (6)$$

332

333 **4.2 Assimilating GRACE observations**

334 Several steps must be taken before GRACE TWS can be assimilated into OpenStreams
 335 `wflow_hbv`. GRACE observations represent average TWS variations over one month, while
 336 the OpenStreams `wflow_hbv` model has a daily time step. In this study, it is assumed that the
 337 average TWS corresponds to the middle of the month. Then, spline interpolation between
 338 consecutive months is used to generate a time series of GRACE TWS variations at five-day

339 intervals. The five-day interval was chosen through trial-and-error to be a good compromise
340 between allowing the ensemble to grow between updates and avoiding implausible jumps. As
341 in any land surface assimilation application, the update results in discontinuities as mass is
342 added or removed from the state but these are not large enough to be obvious when a five-day
343 interval is used (see Sect. 5.1). If the update took place at larger time interval (e.g., once a
344 month) and the entire increment was applied on one day, more significant artefacts or
345 temporal discontinuities would occur (Widiastuti, 2009). In order to convert GRACE
346 variations to absolute values the mean TWS in the study period was calculated from the
347 nominal OpenStreams `wflow_hbv` run and added to the GRACE time series.

348 GRACE observes total TWS, some components of which can be neglected (e.g., nominal
349 OpenStreams `wflow_hbv` simulations indicate that surface water and interception storage
350 contributed by less than 1 % to the estimated TWS). Dry snow is also small averaged over
351 the study area (approximately 2% to the estimated TWS in winter). Only over the Alps (see
352 Fig. 1) is the snow contribution greater (approximately 7%). Therefore, we decided to exclude
353 the dry snow from the state vector. To reconcile GRACE to OpenStreams `wflow_hbv` TWS,
354 we then removed the dry snow component estimated from the nominal run from the GRACE
355 prior to assimilation. Note that in catchments where the dry snow component is more
356 significant, it should not be excluded from the state vector.

357 In the EnKF, the GRACE TWS are calculated and assimilated at each 1-km model grid cell
358 every five days. Because the updated model state $A^a(t)$ was an integrated value of TWS, the
359 increment ($\Delta A(t) = A^a(t) - A(t)$) for every ensemble member needed to be disseminated
360 among the three stores, SM, UZ, and LZ. The information about the distribution of the
361 increment among the different model compartments could be obtained directly from the
362 Kalman filter. However, we chose to carry out the vertical distribution in the way consistent
363 with the OpenStreams `wflow_hbv` model (Fig. 2). While the SM and LZ stores have upper
364 bounds determined by model parameters, UZ does not. As a result, allowing it to update
365 freely in the EnKF runs the risk that it becomes excessively large, which would also have a
366 detrimental effect on runoff. Therefore, the increment is used to adjust the SM first, subject to
367 the upper and lower limits of zero and f_c . Any remaining increment is applied in turn to LZ,
368 up to its upper limit, and only then to UZ.

369 The GRACE observation error is assumed to be 20 mm and horizontal observation error
370 correlations are not considered. The 20 mm value is considered realistic as it was suggested

371 by several independent assessments e.g., Klees et al. (2008), Wahr et al. (2006), Schmidt et al.
372 (2008) and it also had been applied in previous GRACE assimilation studies (Zaitchik et al.,
373 2008; Houborg et al., 2012). Our philosophy was to set the GRACE errors to realistic values
374 determined from independent studies, so that the solutions were not guided towards any
375 particular outcome.

376

377 **4.3 Uncertainty in model forcing data and parameters**

378 In the EnKF, stochastic noise can be included in model forcing data and parameters to
379 account for model uncertainty. An earlier sensitivity study (Widiastuti, 2009) was conducted
380 to identify the parameters of the OpenStreams `wflow_hbv` model that had a significant impact
381 on TWS. Six such parameters, which include fc , lp , β , $cflux$, khq , and $perc$ were found.
382 Therefore, the soil moisture routine parameters, fc , lp and β , as well as the runoff routine
383 parameters, $cflux$, khq and $perc$, were perturbed. For the “calibrated” case, the calibrated
384 model parameters in each grid cell were perturbed using additive Gaussian noise, with a mean
385 of zero and a standard deviation equal to 10% of the range of values that occurred over the
386 whole Rhine basin. In the “non-calibrated” case, the mean parameter value in each grid cell
387 was set to the average calibrated value across the whole basin, and the standard deviation was
388 set to that of the calibrated parameter across the whole basin. This was considered as a proxy
389 for assigning approximate values based on the land cover type, topography, and climatology
390 from the globally available databases. Averaging each parameter across the entire Rhine basin
391 is intended merely to reflect this kind of first-order assumption. Though not all OpenStreams
392 `wflow_hbv` parameters can be gleaned from such global databases, we the averaged values
393 could be compared to those in the Food and Agriculture Organization of the United Nations
394 (FAO) database (<http://www.fao.org/geonetwork/srv/en/main.home>, last access: 5 December
395 2014). The areally averaged parameter values over the Rhine were found to be within the
396 range the provided by FAO. For example, the areally averaged soil moisture field capacity
397 over the Rhine FAO provided is mostly between 150 and 200 mm, while the areally averaged
398 value of approximately 180 mm is used as a mean in this study with a standard deviation of
399 33 cm. The meteorological forcing data were also varied, with the temperature data being
400 perturbed with additive Gaussian noise, and the precipitation and PET being perturbed with
401 additive lognormal noise. In the “local forcing data” case, noise with standard deviation based
402 on 10 % of the nominal value was added to precipitation while 15 % noise was added to

403 temperature and PET. For the “global forcing data” case, we assumed that the local forcing
404 data were accurate and reliable, and the differences between the local and global forcing data
405 represent the [errors](#) of global forcing data. The errors were assumed to be spatially correlated,
406 so an exponential correlation function was applied to the covariance matrix for each variable.
407 The correlation lengths for precipitation, temperature and PET were determined using
408 variogram analysis (Widiastuti, 2009) and found to be 21 km, 21 km, and 59 km, respectively.
409 Recall from Sect. 1 and Sect. 3.2, that four cases are considered in this study: 1) calibrated
410 parameters with local forcing data (CL), 2) calibrated parameters with global forcing data
411 (CG), 3) non-calibrated parameters with local forcing data (NCL), and 4) non-calibrated
412 parameters with global forcing data (NCG). Comparison of the four scenarios provides insight
413 into the benefit of GRACE assimilation under different degrees of uncertainty. The lowest
414 and highest levels of uncertainty are associated with the CL and the NCG cases.

415

416 **5 Results and discussion**

417 Using the EnKF approach described above, GRACE observations were assimilated into the
418 OpenStreams [wflow_hbv](#) model. An ensemble of 100 model states was propagated forward
419 from 1 Jan 2001 to 30 Nov 2003 to spin up the model. The ensemble state at the end of the
420 spin-up period provided the initial state for the assimilation. The study period is from 1 Dec
421 2003 to 31 Oct 2007 because the observed streamflow was only available until Autumn 2007.

422 **5.1 Impact of GRACE assimilation on TWS estimates**

423 First, the impact of assimilating GRACE on the temporal and spatial patterns of the estimated
424 TWS is considered. For the temporal pattern, the areal mean of the estimated TWS over the
425 entire Rhine River basin was computed. The time series of TWS variations from the ensemble
426 open loop (EnOL, ensemble run without GRACE assimilation), EnKF, and GRACE
427 observations are shown in Fig. 5.

428 As expected, there is a seasonal cycle in the TWS estimates, which varies between ± 75 mm.
429 The high frequency variations in TWS in the CL and NCL that are not apparent in CG and
430 NCG are due to the coarser spatial resolution of the global precipitation product. [Lower
431 spatial variability of the global data causes smoother averaged TWS presented in the CG and
432 NCG time series.](#) During the summer of 2006 (June, July, August: JJA), the areal mean global

433 and local precipitation and temperature products agree. However, the global PET product
434 estimates an areal mean PET of 4.10 mm/day while the local PET data suggest it was 2.89
435 mm/day. As the result, the minimum TWS in the CL and NCL cases in the EnOL is -69 mm
436 while CG and NCG are close to -90 mm. In this period, GRACE assimilation has little impact
437 on CL and NCL, but results in a significant (25 mm) update in TWS in the CG and NCG
438 cases. The largest difference between the EnOL and EnKF occurs when TWS is increasing
439 (for example, October 2005). This is apparent in all cases, but is greatest in the two non-
440 calibrated cases. In all cases, Fig. 5 shows that assimilation draws the TWS estimate toward
441 the GRACE observation.

442 The impact of GRACE assimilation also varies within the basin. Fig. 6 shows the spatial
443 distribution of the average increment (posterior minus prior) in TWS during winter
444 (December, January, February: DJF, 2005-2006) and summer (JJA) of 2006. During the
445 winter (left), the EnKF estimated wetter conditions over entire Rhine River basin when the
446 local forcing data were used. In the Alps, the global precipitation product is approximately
447 35% higher than the local precipitation product. Therefore, GRACE assimilation reduced the
448 TWS estimate over the Alps in the CG and NCG cases. During the summer (right), GRACE
449 assimilation reduced the TWS estimate over the Alps and Neckar basin when local forcing
450 data were applied, but adds moisture in the global data case. In this period, the local PET
451 product is 66% lower than the global product over the Alps and 44% lower over the Lahn
452 basin. This is consistent with the increase in areal averaged TWS observed in the CG and
453 NCG cases in Fig. 5. Since the local precipitation data are generally considered to be more
454 accurate, the adjustment of the TWS estimates towards those produced by the local product is
455 an excellent example of the benefit of GRACE assimilation, particularly in data sparse areas.

456 In the Regnitz basin (east of domain), GRACE assimilation leads to a significant increase in
457 TWS in both calibrated cases during the winter months. In this basin, the upper zone
458 recession coefficient (khq) is 0.52 in the calibrated case, compared to 0.3 in the non-calibrated
459 case. This results in almost twice as much fast runoff in the calibrated case, which depletes
460 the terrestrial water storage in the winter months. GRACE assimilation adds moisture to the
461 UZ and LZ stores, drawing the TWS closer to the GRACE observations.

462 In the summer, an average of 0.7 and 1.07 mm was removed in each update from the southern
463 part of Moselle basin in the CL and CG cases, respectively (Fig. 6(b) and (d)), compared to
464 0.74 mm and 1.25 mm added per update in the NCL and NCG cases. In the two calibrated

465 cases, the evaporation threshold value (the product of fc and lp) is approximately 11 % less
466 than that in the non-calibrated cases. This leads to less soil evaporation and higher soil
467 moisture in the calibrated cases. GRACE assimilation reduces the SM in the calibrated cases,
468 and increases it in the non-calibrated cases to draw the TWS closer to the GRACE
469 observations in all cases.

470 **5.2 Impact of GRACE assimilation on GW estimates**

471 The TWS and GW variations from OpenStreams [wflow_hbv](#) were computed at every grid
472 cell. The estimates at the Sundern and A319C wells are shown in Fig. 7 and 8. The two
473 stations represent the behaviour of the other 16 stations (detailed below). For example,
474 stations 2, 3, 4, 6, 9, 10, 11, 13, and 18 have similar behaviour to Sundern, while the rest have
475 similar behaviour to A319C station. Recall that GW is defined as the sum of UZ and LZ, so
476 the difference between the left and right columns is the SM term. GRACE measures monthly
477 variations, so the monthly mean of TWS, GW estimates and the in situ data are shown.
478 Similar to the areal mean values, the TWS from the EnKF in the individual grid cells (left
479 column) is generally between the values from the EnOL and those observed by GRACE.

480 At Sundern (Fig. 7) in the CG and NCG cases, the impact of the forcing data was seen in the
481 summer of every year. Table 2 shows that the precipitation, temperature and PET at Sundern
482 were higher in the global forcing data than in the local data. Fig. 7(c) and 7(g) suggest that
483 this leads to a more negative estimate of TWS in the EnOL for the CG and NCG cases. In the
484 EnKF results, these TWS estimates are drawn towards the GRACE observations. The
485 corresponding updates in terms of GW are larger in the global forcing data case than in the
486 local forcing data cases - assimilation added approximately 5-10 mm of water to GW in the
487 global data cases. Similar behaviour was also seen in CL and NCL cases in summer 2005.

488 At Sundern, the estimated GW in the CL case agrees quite well with the in situ values,
489 suggesting that the distribution between the SM and GW components is reasonable in the
490 calibrated cases. The fact that a good estimate of TWS does not result in an improved GW
491 estimate indicates that the non-calibrated parameters are leading to an incorrect distribution of
492 the TWS between the different stores. In the NCL and NCG cases, fc is just 179 mm
493 compared to the calibrated value of 239 mm. So, for the same TWS value, the non-calibrated
494 cases have more water in GW than the calibrated cases. As a result, despite the agreement in
495 TWS in the winter months, the GW variation is considerably overestimated.

496 In every case at the A319C well location (Fig. 8), the EnOL estimated lower TWS in the first
497 half of 2004 and 2006, and higher in the second half of the same years. Assimilation updated
498 the TWS toward GRACE observation in these periods and resulted in better agreement
499 between the assimilated and observed GW. In late-2005, the estimated TWS from the EnOL
500 and EnKF are very close to the GRACE observations. However, the estimated GW in both
501 cases is a lot lower than that observed in situ. As discussed, the difference between the two is
502 soil moisture. The model is predicting a significant [increase](#) in soil moisture in all four cases.
503 However, given there is little to improve in terms of TWS, the GW estimate from the EnKF is
504 as bad as that from the EnOL.

505 The impact of the forcing data used is also presented. In CG and NCG cases, on 3 Oct and 23
506 Oct 2006, underestimated global precipitation caused the underestimated GW. GRACE could
507 not correct such a high frequency event due to the limitation of its temporal resolution.

508 The choice of the parameters plays a role in the estimated GW magnitude (as seen in Fig. 7),
509 but now the non-calibrated parameters (compared to the calibrated ones) provided closer
510 values to the in situ data (Fig. 8(f) and (h)). Higher non-calibrated f_c parameter (see Table 3
511 for the values) was responsible for smaller GW estimates.

512 Tables 4 and 5 show the correlation coefficient and RMS error (RMSE) between the
513 estimated and in situ GW for all eighteen well locations indicated on Fig. 1. These were
514 calculated based on the monthly mean, but similar results were obtained using the daily
515 values. In most cases, assimilation leads to an increase in correlation coefficient and a
516 reduction in RMSE.

517 The results varied across the wells. The highest correlation coefficients in the EnOL
518 simulations were typically found in the CL case, followed by the NCL. Clearly, using the
519 local forcing data has a significant impact in resolving features at a single grid cell. An
520 exception is the Main basin (wells 5, 7-10) where the global forcing data produce TWS more
521 consistently with the GRACE observations and hence result in a better agreement with the
522 GW. The highest correlation coefficients in the EnKF cases are also found in the two local
523 data cases. The improvements in correlation coefficient are seen in all four cases. The CL and
524 NCL cases also yield the lowest RMSE values in the EnOL case, and the results with the
525 EnKF are very mixed.

526 It is important to note that at many wells, the NCL and NCG cases yield higher correlation
527 coefficients than the CL and CG cases, respectively. Recall that the model is calibrated using

528 streamflow, not groundwater data. So, while assimilation draws the modelled TWS towards
529 the GRACE observations, the model parameters have a significant impact on whether or not
530 this translates to an improvement in GW estimate.

531 One of the objectives was to examine the potential value of GRACE assimilation in data-
532 sparse regions. In the NCG case, it is encouraging that GRACE assimilation consistently
533 leads to an increase in correlation coefficient (up to 33 %) and reduction in RMSE (up to 35
534 %). In other scenarios, assimilation of GRACE observations also leads into an increase in
535 correlation coefficient (up to 71%, at station 11 in the CG case) and a decrease in RMSE (up
536 to 35 %, at station 1 in the NCG case). In average, correlation and RMSE improvements in
537 groundwater estimates for all cases evaluated are 13 % and 14 %, respectively.

538 **5.3 Impact of GRACE assimilation on streamflow estimates**

539 The estimated and observed streamflows at Maxau (upstream) and Wessel (downstream)
540 gauge stations are shown in Fig. 9 and 10. Accurate forcing data, particularly precipitation,
541 are essential for reproducing the observed streamflow. The high frequency variations in
542 streamflow associated with fast response to local precipitation are often reproduced
543 reasonably well in the CL case, but not in the CG case (compare Fig. 9(a) to 9(b) and 10(a) to
544 10(b)).

545 Use of the global data frequently underestimates the streamflow. This is clear on 5 Jun 2004,
546 24 Aug 2005, 6 Oct 2006, and 10 Aug 2007 in Fig. 9(b) and 9(d). Comparing Fig. 9(a) to
547 9(b), it is clear that the larger peaks in streamflow are poorly estimated when the global data
548 are used. Because GRACE observations describe monthly variations over a larger area, they
549 can do little to capture these individual streamflow events. By correcting TWS, GRACE
550 assimilation mainly influences the longer term variations. The difference between EnOL and
551 EnKF is very small in the CL case. The largest differences are observed in the CG and NCG
552 cases, where TWS is updated to correct for errors in forcing data (e.g., summer 2004 and
553 2006 in Fig. 9).

554 Fig. 11 shows the impact of GRACE assimilation on the correlation coefficient, Nash-
555 Sutcliffe coefficient (NS) (Nash and Sutcliffe, 1970), and RMSE in streamflow. Results are
556 shown for four gauge stations along the main channel, as well as the average value across all
557 thirteen stations. These results underscore the importance of forcing data and calibration for
558 estimating streamflow. By far, the highest correlation coefficients and Nash-Sutcliffe

559 coefficients and lowest RMSEs are obtained when local forcing data are used. Use of global
560 forcing data leads to a significant loss in performance. For example, using global rather than
561 local forcing data with the calibrated model results in a decrease in correlation coefficient
562 from 0.89 to 0.65, a decrease in Nash-Sutcliffe coefficient from 0.76 to 0.35 and an increase
563 in RMSE of 71 % in the EnKF results. Using the non-calibrated model rather than the
564 calibrated model also leads to poorer performance, though to a lesser degree. For example,
565 using the non-calibrated rather than calibrated model with the local forcing data results in a
566 decrease in correlation coefficient from 0.89 to 0.88, a decrease in Nash-Sutcliffe coefficient
567 from 0.76 to 0.65 and an increase in RMSE of 23 % in the EnKF results.

568 Compared to the differences due to forcing data and calibration, GRACE assimilation leads to
569 a relatively modest improvement in streamflow estimates. In terms of correlation coefficient,
570 the largest improvements on average (Avg column) are found when the global forcing data
571 are used. The correlation coefficient increased from 0.64 to 0.65 in the CG case, and 0.65 to
572 0.66 in the NCG case. The largest improvement at an individual station was found at Maxau
573 where assimilation resulted in an increase in correlation coefficient from 0.54 to 0.59 in the
574 NCG case.

575 Similarly, GRACE assimilation leads to a modest improvement in terms of NS coefficient.
576 The largest average improvement was from 0.62 to 0.65 in the NCL case. GRACE
577 assimilation slightly reduced the RMSE in all 4 cases. The greatest reduction is 4 % in the
578 NCL case.

579 Though it is encouraging that GRACE assimilation improved the estimated streamflow, these
580 results demonstrate that it clearly cannot replace high quality forcing data or good model
581 calibration.

582

583 **6 Conclusions**

584 The first goal of this study was to investigate the impact of assimilating GRACE into the
585 OpenStreams [wflow_hbv](#) model on the estimated terrestrial water storage, groundwater
586 storage and streamflow in the Rhine river basin. GRACE observations were assimilated into
587 each grid cell of the model with an EnKF to update the soil moisture and upper and lower
588 zone storage terms of the model. In general, assimilation drew the EnOL estimated TWS
589 closer to the GRACE observations. In the absence of independent TWS observations, a

590 qualitative analysis of the increments in TWS indicated that GRACE assimilation could
591 partially correct the TWS estimate for the influence of errors in the meteorological forcing
592 data and model parameters. As result, an improvement in groundwater estimate after
593 assimilating GRACE data was noticeable, with an overall improvement up to 71%
594 (correlation coefficient) and 35% (RMS error) over the EnOL case. However, it is found that
595 the improvement in TWS estimates did not always translate to an improved agreement
596 between the estimated and observed groundwater storage variation at certain well locations.
597 The differences may be due to the OpenStreams [wflow_hbv](#) parameters: if the upper limit on
598 soil moisture storage is too high (low), then the groundwater variations could be under (over)-
599 estimated. This is particularly relevant in the type of model where the calibration is per sub-
600 basin. This does not allow for local differences on the order of single or a few grid cells. The
601 issue of scale is also significant because GRACE observes monthly variations on the order of
602 hundreds of kilometres. Groundwater variations can be influenced by local features at finer
603 scales. [When the basin average is considered](#), validation against a denser network of well data
604 or an independent groundwater model could be used to determine if an improvement occurs at
605 the scale of the entire basin.

606 Furthermore, the considered model was used to simulate runoff. The groundwater terms, UZ
607 and LZ, primarily serve as reservoirs for quick and base runoff generation. Due to the coarse
608 resolution of the observations, GRACE assimilation resulted in only a modest improvement in
609 streamflow estimates. Correlation coefficients increased by up to 2 %, Nash Sutcliffe
610 coefficients increased by up to 4 % and RMSE was reduced by up to 4 %.

611 The second goal of this study was to investigate the potential value of assimilating GRACE
612 observations in data-sparse regions. Results from four scenarios were compared in which the
613 ensemble mean model parameters were either calibrated values, or basin average values and
614 the meteorological forcing data were either local (high quality) data or global (poorer quality)
615 data. By comparing the four cases, it was shown that GRACE assimilation could correct for
616 errors in model forcing data and parameter calibration by drawing the estimated TWS toward
617 that observed by GRACE. This also resulted in drawing the estimated groundwater storage
618 closer to the in situ measurement. [Given that the most significant improvements were
619 observed in the NCG case, this suggests that GRACE observations are most valuable in data
620 sparse regions. In these regions any additional observations, even those at coarse spatial
621 and/or temporal resolution, are welcome. GRACE can provide essential independent](#)

622 observations for validation, and serves as a constraint for TWS within the assimilation
623 process. In terms of streamflow, a comparison of the four scenarios demonstrates that the
624 ability to capture high flow events is determined largely by the quality of the forcing data and
625 the model parameters. The improvements in streamflow estimates after assimilation are
626 modest. Nevertheless, we consider the obtained results as promising, particularly in data-
627 sparse scenarios, e.g., the NCG case. They indicate that GRACE contains information that can
628 be useful for streamflow estimation. Whether updating TWS is the best way to use this
629 information is an open question. An alternative strategy could be to use GRACE assimilation
630 for parameter estimation at a sub-basin or basin scale and constrain the rainfall-runoff model
631 through assimilation of soil moisture observations.

632 In conclusion, GRACE assimilation is beneficial, and the largest improvements are generally
633 observed in the NCG (i.e. “data-sparse”) cases. In addition to providing a modest
634 improvement to the estimated streamflow, it may result in a noticeable improvement in TWS
635 estimates, yielding an extra insight into the behaviour of the hydrological model, its forcing
636 data and parameters. Further research will combine assimilation of GRACE and a soil
637 moisture remote sensing product to constrain the SM estimate storage term, and ensure that
638 improved TWS would lead to more consistently improved estimates of groundwater storage
639 variations. Further research will also explore the value of assimilating GRACE into a
640 groundwater model in which the primary processes of interest vary on temporal and spatial
641 scales similar to those of GRACE. In addition, recent studies have explored the effect of
642 spatial aggregation of GRACE TWS prior to assimilation (Forman and Reichle, 2013) as well
643 as inclusion of the full GRACE error structure (Eicker et al., 2014). Combining the advances
644 made in those studies with the assimilation framework presented here is expected to yield
645 even more realistic estimates. As shown by De Lannoy et al. (2009), working with a spatially
646 distributed state vector (3D-EnKF) can lead to an improved estimate. Given the coarse
647 resolution of GRACE, we expect that implementing a 3D-EnKF within the assimilation
648 framework would lead to an improved performance. This could be particularly important in
649 small basins like the Rhine, and can be used to account for the fact that the GRACE
650 overpasses are infrequent and may not be sensitive to TWS variations in response to specific
651 events.

652

653

654 **Appendix A: Names of well locations**

655 Table A1: Names of the well locations used in this paper.

Location number	Name	Source
1	Sundern	Ministerium für Klimaschutz, Umwelt,
2	GEW KOELN 557	Landwirtschaft, Natur- und
3	SHELL GODORF GW I	Verbraucherschutz des
4	LGD BN-BEUEL	LandesNordrhein-Westfalen(http://www.elwasweb.nrw.de)
5	Stetten S1	
6	Dietersdorf	
7	Haßfurt Q2	Bayerisches Landesamt für
8	Limbach Q1	Umwelt(http://www.gkd.bayern.de)
9	Rattelsdorf 136	
10	Faulbach	
11	01373X0130/A25	
12	02303X0065/P	
13	02307X0281/S	
14	01995X0030/563	Portail national d'Accès aux
15	02344X0082/326E	Données sur les Eaux
16	02344X0055/319	Souterraines(http://www.adeseaufrance.fr)
17	02348X0009/319C (called A319C in this paper)	
18	03426X0197/136	

656

657

658

659 **Acknowledgements**

660 This research was funded by The Netherlands Organisation for Scientific Research, NWO
661 (project number 842.00.006). The work of S. C. Steele-Dunne was supported by the NWO
662 Veni Grant Program (ALW 863.09.015). We would like to thank Eric Sprokkereef Secretary
663 of the International Commission for the Hydrology of the Rhine basin for granting permission
664 for the use of hydrologic data. Jaap Schellekens (Deltares) is acknowledged for help with the
665 OpenStreams model of the Rhine. The authors also want to thank Edwin Sutanudjaja for
666 productive discussions about the groundwater data.

667

668 **References**

- 669 Bader, S., Belz, J. U., Jakob, A., Mathis, C., Morf-Graf, S., Mürlebach, M., and Schürch, M.:
670 International commission for the hydrology of the Rhine basin (CHR): Annual CHR
671 report 2012, Sprokkereef, E, Annual CHR report 2012, Lelystad, the Netherlands, 2013.
- 672 Becker, M., Meyssignac, B., Xavier, L., Cazenave, A., Alkama, R., and Decharm, B.: Past
673 terrestrial water storage (1980–2008) in the Amazon Basin reconstructed from GRACE
674 and in situ river gauging data, *Hydrol. Earth Syst. Sci.*, 15, 533–546, doi:10.5194/hess-15-
675 533-2011, 2011.
- 676 Bettadpur, S.: Insights into the Earth System mass variability from CSR-RL05 GRACE
677 gravity fields, EGU Meeting, abstract #EGU2012-6409, Vienna, Austria, 2012.
- 678 Bierkens, M. F. P, and van den Hurk, B. J. J. M.: Groundwater convergence as a possible
679 mechanism for multi-year persistence in rainfall, *Geophys. Res. Lett.*, 34, L02402,
680 doi:10.1029/2006GL028396, 2007.
- 681 Cheng, M., and Tapley, B.: Variations in the Earth's oblateness during the past 28 years, *J.*
682 *Geophys. Res.*, 109(B09402), doi:10.1029/2004JB003028, 2004.
- 683 de Jeu, R. A. M., Wagner, W., Holmes, T. R. H., Dolman, A. J., van de Giesen, N. C.,
684 Friesen, J.: Global Soil Moisture Patterns Observed by Space Borne Microwave
685 Radiometers and Scatterometers, *Surv. Geophys.*, 28, 399-420 doi 10.1007/s10712-008-
686 9044-0, 2008.
- 687 De Lannoy, G. J. M., Reichle, R. H., Houser, P. R., Arsenault, K. R., Verhoest, N. E. C., and
688 Pauwels, R. N.: Satellite-Scale Snow Water Equivalent Assimilation into a High-
689 Resolution Land Surface Model, *J. Hydrometeor.*, 11, 352–369,
690 doi:http://dx.doi.org/10.1175/2009JHM1192.1, 2009.
- 691 Dorigo, W. A., Wagner, W., Hohensinn, R., Hahn, S., Paulik, C., Xaver, A., Gruber, A.,
692 Drusch, M., Mecklenburg, S., van Oevelen, P., Robock, A., and Jackson, T.: The
693 International Soil Moisture Network: a data hosting facility for global in situ soil moisture
694 measurements, *Hydrol. Earth Syst. Sci.*, 15, 1675-1698, 2011.
- 695 Eberle, M., Buiteveld, H., Beersma, J., Krahe, P., and Wilke, K.: Estimation of extreme
696 floods in the river Rhine basin by combining precipitation-runoff modelling and a rainfall
697 generator, in: Proceedings International Conference on Flood Estimation, Berne, CHR

698 report II-17, International Commission for the Hydrology of the Rhine basin (CHR),
699 IJsselstein, The Netherlands, 2002.

700 Eberle, M., Buiteveld, H., Krahe, P., and Wilke, K.: Hydrological Modelling in the River
701 Rhine Basin, Part III: Daily HBV model for the Rhine basin, Report 1451, Bundesanstalt
702 für Gewässerkunde (BFG), Koblenz, Germany, 2005.

703 Entekhabi, D., Njoku, E. G., O'Neill, P. E., Kellogg, K. H., Crow, W. T., Edelstein, W. N.,
704 Entin, J. K., Goodman, S. D., Jackson, T. J., Johnson, J., Kimball, J., Piepmeier, J. R.,
705 Koster, R. D., Martin, N., McDonald, K. C., Moghaddam, M., Moran, S., Reichle, R., Shi,
706 J. C., Spencer, M. W., Thurman, S. W., Leung T., van Zyl, J.: The Soil Moisture Active
707 Passive (SMAP) Mission, Proceedings of the IEEE, 98, 704–716,
708 doi:10.1109/JPROC.2010.2043918, 2010.

709 [Eicker, A., Schumacher, M., Kusche, J., Döll, P., Müller Schmied H.: Calibration data
710 assimilation approach for integrating GRACE data into the WaterGAP Global Hydrology
711 Model \(WGHM\) using an Ensemble Kalman Filter: First Results, Surv. Geophys., 35\(6\),
712 1285-1309, doi:10.1007/s10712-014-9309-8, 2014.](#)

713 Evensen, G.: Sequential data assimilation with a non-linear quasi-geostrophic model using
714 Monte Carlo methods to forecast error statistics, J. Geophys. Res., 99(C5), 10143–10162,
715 1994.

716 Evensen, G.: The ensemble Kalman filter: Theoretical formulation and practical
717 implementation, Ocean Dyn., 53(4), 343-367, doi:10.1007/S10236-003-0036-9, 2003.

718 [Ferrant, S., Caballero, Y., Perrin, J., Gascoin, S., Dewandel, B., Aulong, S., Dazin, F.,
719 Ahmed, S., and Maréchal J.-P.: Projected impacts of climate change on farmers'
720 extraction of groundwater from crystalline aquifers in South India, Nature Sci. Rep., 4,
721 3697, doi:10.1038/srep03697, 2014.](#)

722 Forman, B. A., Reichle, R. H., and Rodell, M.: Assimilation of terrestrial water storage from
723 GRACE in snow-dominated basin, Water Resour. Res., 48, W01507,
724 doi:10.1029/2011WR011239, 2012.

725 [Forman, B. A. and Reichle, R.: The spatial scale of model errors and assimilated retrievals in
726 a terrestrial water storage assimilation system. Water Resour. Res., 49, 7457–7468.
727 doi:10.1002/2012WR012885, 2013.](#)

728 Green, T. R., Taniguchi, M., Kooi, H., Gurdak, J. J., Allen, D. M., Hiscock, K. M., Treidel,
729 H., and Aureli, A.: Beneath the surface of global change: Impacts of climate change on
730 groundwater, *J. Hydrol.*, 405, 532-560, doi:10.1016/j.jhydrol.2011.05.002, 2011.

731 Güntner, A.: Improvement of global hydrological models using GRACE data, *Surv.*
732 *Geophys.*, 29, 375-397, doi:10.1007/s10712-008-9038-y, 2008.

733 Haylock, M. R., Hofstra, N., Klein Tank, A. M. G., Klok, E. J., Jones, P. D., and New, M.: A
734 European daily high-resolution gridded dataset of surface temperature and precipitation, *J.*
735 *Geophys. Res.*, 113, D20119, doi:10.1029/2008JD010201, 2008.

736 Hirschi, M., Seneviratne, S. I., Hagemann, S., and Schär, C.: Analysis of seasonal terrestrial
737 water storage variations in regional climate simulations over Europe, *J. Geophys. Res.*,
738 112, D22109, doi: 10.1029/2006JD008338, 2007.

739 Houborg, R., Rodell, M., Li, B., Reichle, R., and Zaitchik, B. F.: Drought indicators based on
740 model-assimilated Gravity Recovery and Climate Experiment (GRACE) terrestrial water
741 storage observations, *Water Resour. Res.*, 48, W07525, doi:10.1029/2011WR011291,
742 2012.

743 Hrachowitz, M., Savenije, H. H. G., Blöschl, G., McDonnell, J. J., Sivapalan, M., Pomeroy,
744 J. W., Arheimer, B., Blume, T., Clark, M. P., Ehret, U., Fenicia, F., Freer, J. E., Gelfan,
745 A., Gupta, H. V., Hughes, D. A., Hut, R. W., Montanari, A., Pande, S., Tetzlaff, D.,
746 Troch, P. A., Uhlenbrook, S., Wagener, T., Winsemius, H. C., Woods, R. A., Zehe, E.,
747 and Cudennec, C.: A decade of Predictions in Ungauged Basins (PUB) - A review.
748 *Hydrolog. Sci. J.*, 58(6), 1198-1255, doi: 10.1080/02626667.2013.803183, 2013.

749 Jekeli, C.: Alternative methods to smooth the Earth's gravity field, *Rep.*, 327, Dept. of Geod.
750 *Sci. and Surv.*, Ohio State Univ., Columbus, 1981.

751 Jung, M., Reichstein, M., Ciais, P., Seneviratne, S. I., Sheffield, J., Goulden, M. L., Bonan,
752 G., Cescatti, A., Chen, J., de Jeu, R., Dolman, A. J., Eugster, W., Gerten, D., Gianelle, D.,
753 Gobron, N., Heinke, J., Kimball, J., Law, B. E., Montagnani, L., Mu, Q., Mueller, B.,
754 Oleson, K., Papale, D., Richardson, A. D., Rouspard, O., Running, S., Tomelleri, E.,
755 Viovy, N., Weber, U., Williams, C., Wood, E., Zaehle, S., and Zhang, K.: Recent decline
756 in the global land evapotranspiration trend due to limited moisture supply, *Nature*, 467,
757 951–954, doi:10.1038/nature09396, 2010.

758 Karszenberg, D., Schmitz, O., Salamon, P., De Jong, K., and Bierkens, M. F. P.: A software
759 framework for construction of process-based stochastic spatio-temporal models and data
760 assimilation, *Environ. Modell. Softw.*, 25, 489-502, 2009.

761 Kerr, Y. H., Waldteufel, P., Richaume, P., Wigneron, J. P., Ferrazzoli, P., Mahmoodi, A.,
762 Bitar, A. A., Cabot, F., Gruhier, C., Juglea, S. E., Leroux, D., Mialon, A., and Delwart, S.:
763 The SMOS Soil Moisture Retrieval Algorithm, *IEEE Trans. Geosci. Remote Sens.*, 50,
764 1384-1403, doi:10.1109/TGRS.2012.2184548, 2012.

765 Klees, R., Liu, X., Wittwe, T., Gunter, B. C., Revtova, E. A., Tenzer, R., Ditmar, P.,
766 Winsemius, H. C., and Savenije, H. H. G.: A Comparison of Global and Regional GRACE
767 Models for Land Hydrology, *Surv. Geophys.*, 29, 335-359, doi:10.1007/s10712-008-9049-
768 8, 2008.

769 [Kleinherenbrink, M., Ditmar, P. G., and Lindenbergh, R. C.: Retracking Cryosat data in the
770 SARIn mode and robust lake level extraction, *Remote Sens. Environ.*, 152, 38-50,
771 doi:10.1016/j.rse.2014.05.014, 2014.](#)

772 Koster, R. D., Suarez, M. J., Ducharne, A., Stieglitz, M., and Kumar, P.: A catchment-based
773 approach to modeling land surface processes in a general circulation model. 1. Model
774 structure, *J. Geophys. Res.*, 105, 24809-24822, 2000.

775 Koster, R. D., Manhanama, S. P. P., Yamada, T. J., Balsamo, G., Berg, A. A., Boisserie, M.,
776 Dirmeyer, P. A., Doblas-Reyes, F. J., Drewitt, G., Gordon, C. T., Guo, Z., Jeong, J.-H.,
777 Lawrence, D. M., Lee, W.-S., Li, Z., Luo, L., Malyshev, S., Merryfield, W. J.,
778 Seneviratne, S. I., Stanelle, T., van den Hurk, B. J. J. M., Vitart, F., and Wood, E. F.:
779 Contribution of land surface initialization to subseasonal forecast skill: First results from a
780 multi-model experiment, *Geophys. Res. Lett.*, 37, L02402, doi:10.1029/2009GL041677,
781 2010.

782 Landerer, F. W., and Swenson, S. C.: Accuracy of scaled GRACE terrestrial water storage
783 estimates, *Water Resour. Res.*, 48, W04531, , doi:10.1029/2011WR011453, 2012.

784 Lettenmaier, D. P., and Famiglietti, J. S.: Water from on high, *Nature*, 444, 562–563, 2006.

785 Li, B., Rodell, M., Zaitchik, B. F., Reichle, R. H., Koster, R. D., and van Dam, T. M.:
786 Assimilation of GRACE terrestrial water storage into a land surface model: Evaluation
787 and potential value for drought monitoring in western and central Europe, *J. Hydrol.*, 446-
788 447, 103-115, 2012.

789 Lindström, G., Johansson, B., Persson, M., Gardelin, M., and Bergström, S.: Development
790 and test of the distributed HBV-96 hydrological model, *J. Hydrol.*, 201(1-4), 272-288,
791 1997.

792 Longuevergne, L., Wilson, C. R., Scanlon, B. R., and Crétaux, J. F.: GRACE water storage
793 estimates for the Middle East and other regions with significant reservoir and lake storage,
794 *Hydrol. Earth Syst. Sci.*, 17, 4817-4830, doi:10.5194/hess-17-4817-2013, 2013.

795 Mülders, R., Parmet, B., and Wilke, K.: Hydrological Modelling in the river Rhine basin,
796 final report, Report 1215, Bundesanstalt für Gewässerkunde (BFG), Koblenz, Germany,
797 1999.

798 Nash, J. E., and Sutcliffe, J. V.: River flow forecasting through conceptual models part I — A
799 discussion of principles, *J. Hydrol.*, 10(3), 282–290, 1970.

800 Orsolini, Y. J., Senan, R., Balsamo, G., Doblas-Reyes, F. J., Vitart, F., Weisheimer, A.,
801 Carrasco, A., and Benestad, R. E.: Impact of snow initialization on sub-seasonal forecasts,
802 *Clim. Dyn.*, 41, 1969-1982, doi:10.1007/s00382-013-1782-0, 2013.

803 Peltier, W. R.: Global glacial isostasy and the surface of the ice-age Earth: The ICE-5G
804 (VM2) model and GRACE, *Annu. Rev. Earth Planet. Sci.*, 32, 111-149, 2004.

805 [Phan, V. H., Lindenbergh, R. C., and Menenti, M.: ICESat derived elevation changes of
806 Tibetan lakes between 2003 and 2009, *Int. J. Appl. Earth Obs. Geoinf.*, 17, 12-22,
807 doi:10.1016/j.jag.2011.09.015, 2012.](#)

808 Photiadou, C. S., Weerts, A. H., van den Hurk, B. J. J. M.: Evaluation of two precipitation
809 data sets for the Rhine River using streamflow simulations, *Hydrol. Earth Syst. Sci.*, 15,
810 3355–3366, doi:10.5194/hess-15-3355-2011, 2011.

811 Rakovec, O., Weerts, A. H., Hazenberg, P., Torfs, P. J. J. F., and Uijlenhoet, R.: State
812 updating of a distributed hydrological model with Ensemble Kalman Filtering: effects of
813 updating frequency and observation network density on forecast accuracy, *Hydrol. Earth
814 Syst. Sci.*, 16, 3435-3449, doi:10.5194/hess-16-3435-2012 , 2012.

815 Reichle, R. H.: Data assimilation methods in the Earth sciences, *Adv. Water Resour.*, 31,
816 1411-1418, doi:10.1016/j.advwatres.2008.01.001, 2008.

817 Rodell, M., Houser, P. R., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C. J., Arsenault,
818 K., Cosgrove, B., Radakovich, J., Bosilovich, M., Entin, J. K., Walker, J. P., Lohmann, D.,

819 and Toll, D.: The global land data assimilation system, *Bull. Amer. Meteor. Soc.*, 85(3),
820 381–394, 2004.

821 Rodell, M., Chen, J., Kato, H., Famiglietti, J. S., Nigro, J., and Wilson, C. R.: Estimating
822 groundwater storage changes in the Mississippi River basin (USA) using GRACE,
823 *Hydrogeol. J.*, 15, 159-166, 2007.

824 Rodell, M., Velicogna, I., and Famiglietti, J. S.: Satellite-based estimates of groundwater
825 depletion in India, *Nature*, 460, 999-1002, 2009.

826 Schellekens, J.: [OpenStream wflow documentation release 1.0RC1](http://schj.home.xs4all.nl/html), Deltares,
827 <http://schj.home.xs4all.nl/html> (last access: 5 January 2015), 2014.

828 Senay, G. B., Verdin, J. P., Lietzow, R., and Melesse, A. M.: Global daily reference
829 evapotranspiration modeling and evaluation, *J. Am. Water Resour. Assoc.*, 44(4), 969-
830 979, doi:10.1111/j.1752-1688.2008.00195.x, 2008.

831 Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B.,
832 and Teuling, A.: Investigating soil moisture-climate interactions in a changing climate: A
833 review, *Earth-Sci. Rev.*, 99, 125-161, doi:10.1016/j.earscirev.2010.02.004, 2010.

834 Schwatke, C., Bosch, W., and Dettmering D.: [A new Database of Water Level Time Series
835 for Lakes, Rivers, and Wetlands from Multi-Mission Satellite Altimetry](#), IAG Scientific
836 [Assembly 2013, Potsdam, Germany, 2013-09-01/06](#), 2013.

837 Sheffield, J., Goteti, G., and Wood, E. F.: Development of a 50-yr high-resolution global
838 dataset of meteorological forcings for land surface modeling, *J. Climate*, 19 (13), 3088-
839 3111, 2005.

840 Sivapalan, M., Takeuchi, K., Franks, S. W., Gupta, V. K., Karambiri, H., Lakshmi, V., Liang,
841 X., McDonnell, J. J., Mendiola, E. M., O’Connell, P. E., Oki, T., Pomeroy, J. W.,
842 Schertzer, D., Uhlenbrook, S., and Zehe, E.: IAHS decade on predictions in ungauged
843 basins (PUB) 2003–2012: shaping an exciting future for the hydrological sciences,
844 *Hydrolog. Sci. J.*, 48,857–880, 2003.

845 Su, H., Yang, Z.-L., Dickinson, R. E., Wilson, C. R., and Niu, G.-Y.: Multisensor snow data
846 assimilation at the continental scale: The value of Gravity Recovery and Climate
847 Experiment terrestrial water storage information, *J. Geophys. Res.*, 15, D10104,
848 doi:10.1029/2009JD013035, 2010.

- 849 Swenson, S., and Wahr, J.: Post-processing removal of correlated errors in GRACE data,
850 Geophys. Res. Lett., 33(L08402), doi:10.1029/2005GL025285, 2006.
- 851 Swenson, S., Chambers, D., and Wahr, J.: Estimating geocenter variations from a
852 combination of GRACE and ocean model output, J. Geophys. Res., 113(B08410),
853 doi:10.1029/2007JB005338, 2008.
- 854 Syed, T. H., Famiglietti, J. S., and Chambers, D.: GRACE-based estimates of terrestrial
855 freshwater discharge from basin to continental scales, J. Micrometeorol., 10, 22-40,
856 doi:10.1175/2008JHM993.1, 2008.
- 857 Tapley, B. D., Bettadpur, S., Ries, J. C., Thompson, P. F., and Watkins, M.: GRACE
858 Measurements of Mass Variability in the Earth System, Science, 305 (5683), 503-505,
859 2004.
- 860 Tregoning, P., McClusky, S., van Dijk, A. I. J. M., Crosbie, R. S., and Peña-Arancibia, J. L.:
861 Assessment of GRACE Satellites for Groundwater Estimation in Australia, National
862 Water Commission, Canberra, 82, 2012.
- 863 van den Hurk, B., Doblas-Reyes, F., Balsamo, G., Koster, R. D., Seneviratne, S. I., and
864 Camargo Jr, H.: Soil moisture effects on seasonal temperature and precipitation forecast
865 scores in Europe, Clim. Dyn., 28, 349-362, 2012.
- 866 [Wada, Y., Wisser, D., and Bierkens, M. F. P.: Global modeling of withdrawal, allocation and
867 consumptive use of surface water and groundwater resources, Earth Syst. Dynam., 5, 15-
868 40, doi:10.5194/esd-5-15-2014, 2014.](#)
- 869 Wahr, J., Molenaar, M., and Bryan, F.: Time variability of the Earth's gravity field:
870 Hydrological and oceanic effects and their possible detection using GRACE, J. Geophys.
871 Res., 103(B12), 30205–30229, 1998.
- 872 Wahr, J., Swenson, S., and Velicogna, I.: Accuracy of GRACE mass estimates, Geophys.
873 Res. Lett., 33, L06401, doi:10.1029/2005GL025305, 2006.
- 874 Weerts, A. H., and Serafy G. Y. H.: Particle filtering and ensemble Kalman filtering for state
875 updating with hydrological conceptual rainfall-runoff models, Water Resour. Res., 42,
876 W09403, doi:10.1029/2005WR004093, 2006.
- 877 Weerts, A. H., Meißner, D., and Rademacher, S.: Input data rainfall-runoff model operational
878 systems FEWS-NL & FEWS-DE, Tech. rep., Deltares, 2008.

- 879 Widiastuti, E.: Data assimilation of GRACE terrestrial water storage data into a hydrological
880 model using the ensemble Kalman smoother: A case study of the Rhine river basin, MSc
881 Thesis, TU Delft, 2009.
- 882 Zaitchik, B. F., Rodell, M., and Reichle, E. H.: Assimilation of GRACE terrestrial water
883 storage data into a land surface model: Results for the Mississippi basin, Amer. Meteor.
884 Soc., *J. Hydrometeor*, 9, 535–548, 2008.
- 885

886 Table 1. Parameters of the soil moisture and runoff routines in the OpenStreams [wflow_hbv](#)
 887 model.

Parameter	Description	Unit
fc	Maximum soil moisture storage	mm
β	Empirical based parameter determines the relative contribution to runoff from soil moisture storage	-
$cflux$	Maximum value of capillary rise from upper zone storage to soil moisture storage	mm/day
khq	Recession constant of upper zone storage, determines the amount of quick runoff from upper zone storage	1/day
$perc$	Maximum percolation value (from upper to lower zone storage)	mm/day
lp	Soil moisture fraction above which actual evapotranspiration (ET) equals potential ET	-
$k4$	Recession constant of lower zone storage, determines the amount of baseflow from lower zone storage	1/day
α	Non-linearity parameter of upper zone storage	-

888

889 Table 2. Daily mean values of the forcing data at Sundern during summer (JJA) months.

	Precipitation (mm)		Temperature (°C)		PET (mm)	
	Local	Global	Local	Global	Local	Global
2004	5.48	5.21	16.10	17.65	2.47	3.30
2005	5.25	4.68	16.12	17.46	2.47	3.55
2006	4.96	4.39	17.64	19.25	2.47	3.90
2007	9.97	7.08	16.09	17.52	2.47	3.25

890

891 Table 3. Ensemble mean parameter values at the Sundern, and A319C well locations for the
892 calibrated and non-calibrated simulations.

Parameter	Sundern		A319C	
	Calibrated	Non-calibrated	Calibrated	Non-calibrated
<i>fc</i>	239.03	179.12	130.95	181.98
<i>β</i>	2.06	1.65	1.95	1.68
<i>cflux</i>	0.06	0.27	0.41	0.30
<i>khq</i>	0.10	0.12	0.06	0.09
<i>perc</i>	0.67	1.15	0.43	1.09
<i>lp</i>	0.88	0.75	0.67	0.72
<i>k4</i>	0.63	0.03	0.01	0.03

893


894 Table 4. Correlation coefficient computed between monthly mean estimated GW variation
 895 and monthly mean in situ variation. Names of the stations (first column) are provided in
 896 Appendix A.

	CL		CG		NCL		NCG	
	EnOL	EnKF	EnOL	EnKF	EnOL	EnKF	EnOL	EnKF
1	0.85	0.83	0.71	0.74	0.79	0.85	0.70	0.78
2	0.57	0.68	0.32	0.52	0.43	0.65	0.38	0.45
3	0.69	0.81	0.46	0.68	0.51	0.73	0.46	0.61
4	0.60	0.67	0.42	0.58	0.50	0.75	0.60	0.67
5	0.71	0.68	0.67	0.76	0.71	0.72	0.72	0.78
6	0.57	0.64	0.58	0.66	0.74	0.78	0.66	0.72
7	0.77	0.80	0.67	0.71	0.80	0.83	0.64	0.72
8	0.75	0.80	0.65	0.78	0.81	0.83	0.62	0.74
9	0.50	0.64	0.54	0.70	0.72	0.78	0.65	0.80
10	0.56	0.58	0.50	0.55	0.66	0.70	0.42	0.46
11	0.41	0.55	0.31	0.53	0.71	0.73	0.72	0.74
12	0.71	0.80	0.64	0.71	0.76	0.85	0.74	0.78
13	0.77	0.80	0.50	0.56	0.72	0.84	0.59	0.66
14	0.71	0.70	0.73	0.74	0.33	0.47	0.51	0.56
15	0.82	0.85	0.67	0.69	0.72	0.83	0.65	0.71
16	0.68	0.80	0.55	0.64	0.77	0.88	0.63	0.71
17	0.67	0.79	0.55	0.60	0.70	0.82	0.57	0.66
18	0.65	0.66	0.64	0.65	0.45	0.54	0.59	0.63
Mean	0.67	0.73	0.56	0.66	0.66	0.75	0.60	0.68

897

898 Table 5. RMSE [mm] computed between monthly mean estimated GW variation and monthly
 899 mean in situ variation. Names of the stations (first column) are provided in Appendix A.

	CL		CG		NCL		NCG	
	EnOL	EnKF	EnOL	EnKF	EnOL	EnKF	EnOL	EnKF
1	4.16	3.84	5.63	4.02	7.00	6.99	8.37	5.40
2	5.34	4.91	6.66	5.96	6.36	5.73	10.78	8.14
3	3.62	3.06	5.04	4.35	5.96	4.85	10.64	8.13
4	3.79	3.65	4.41	3.55	5.83	5.03	9.00	7.56
5	9.72	8.30	6.89	5.49	8.43	7.83	6.06	5.03
6	6.19	5.19	5.47	5.26	7.56	6.31	5.25	4.29
7	8.30	6.75	7.48	6.99	6.45	5.88	7.36	6.82
8	8.76	6.59	6.63	4.96	5.21	5.20	5.71	4.67
9	5.95	5.38	5.16	5.09	7.33	6.43	5.43	3.91
10	8.95	7.64	6.44	5.66	8.92	8.54	7.62	6.21
11	6.03	5.10	6.21	4.89	9.88	8.32	11.43	8.30
12	7.17	6.37	7.33	7.42	6.24	5.01	6.58	5.95
13	6.25	5.34	6.80	5.91	7.90	7.55	8.84	8.53
14	12.67	10.16	11.43	9.01	9.34	7.97	9.29	7.21
15	8.83	8.28	10.28	10.08	9.78	8.31	10.08	9.89
16	12.74	9.58	13.20	10.60	9.76	8.32	10.44	9.78
17	12.01	9.17	11.10	9.54	7.59	6.14	7.90	6.38
18	7.23	7.50	8.62	7.94	9.59	8.42	9.51	7.88
Mean	7.65	6.49	7.49	6.48	7.73	6.82	8.35	6.89




901

902 Figure 1. River gauge (circle) and well (triangle) locations over the Rhine River basin used in
 903 this paper. Red triangles indicate Sundern (1) and A319C locations (17). Names of all well
 904 locations are given in Table A1.


905

906 (a)



907

908 (b)




909

910 Figure 2. (a) OpenStreams [wflow_hbv](#) model structure, adapted from Schellekens (2014).


911 Sample results of the nominal run related to soil moisture (SM), upper groundwater zone (UZ),

912 and lower groundwater zone (LZ) storages averaged over Rhine River basin.



913


914 Figure 3. Calibrated parameters of the soil moisture and runoff response routines of the
 915 OpenStreams *wflow_hbv* model.



916

917 Figure 4. Mean daily precipitation, temperature, and potential evapotranspiration in 2006


918 from the local (left) and global (right) forcing datasets.



919

920 Figure 5. Area-averaged mean terrestrial water storage (TWS) over the Rhine River basin
 921 from the EnOL, EnKF and GRACE observations in 4 different scenarios (CL: calibrated
 922 parameters with local forcing data, CG: calibrated parameters with global forcing data, NCL:
 923 non-calibrated parameters with local forcing data, NCG: non-calibrated parameters with
 924 global forcing data).


925



926


927 Figure 6. Averaged increment (posterior minus prior) of TWS in mm during the winter 2005-
 928 2006 (left) and summer of 2006 (right) in 4 different scenarios (CL: calibrated parameters
 929 with local forcing data, CG: calibrated parameters with global forcing data, NCL: non-

930 calibrated parameters with local forcing data, NCG: non-calibrated parameters with global
931 forcing data). The polygons in the right column define the southern part of Moselle basin.




932

933 Figure 7. TWS variation (left) and GW variation (right) at the Sudern well location in 4
 934 different scenarios (CL: calibrated parameters with local forcing data, CG: calibrated
 935 parameters with global forcing data, NCL: non-calibrated parameters with local forcing data,
 936 NCG: non-calibrated parameters with global forcing data).




937

938 Figure 8. TWS variation (left) and GW variation (right) at the A319C well location in 4
 939 different scenarios (CL: calibrated parameters with local forcing data, CG: calibrated
 940 parameters with global forcing data, NCL: non-calibrated parameters with local forcing data,
 941 NCG: non-calibrated parameters with global forcing data).




942

943 Figure 9. Estimated and observed streamflow at the Maxau gauge station in 4 different
 944 scenarios (CL: calibrated parameters with local forcing data, CG: calibrated parameters with
 945 global forcing data, NCL: non-calibrated parameters with local forcing data, NCG: non-
 946 calibrated parameters with global forcing data).



947

948 Figure 10. Estimated and observed streamflow at the Wesel gauge station in 4 different
 949 scenarios (CL: calibrated parameters with local forcing data, CG: calibrated parameters with
 950 global forcing data, NCL: non-calibrated parameters with local forcing data, NCG: non-
 951 calibrated parameters with global forcing data).



952

953 Figure 11. The correlation coefficient (left), Nash-Sutcliffe coefficient (middle) and RMS
 954 error (right) computed between estimated streamflows and gauge measurements in 4 different
 955 scenarios (CL: calibrated parameters with local forcing data, CG: calibrated parameters with
 956 global forcing data, NCL: non-calibrated parameters with local forcing data, NCG: non-
 957 calibrated parameters with global forcing data). Results are shown for the Maxau (Max),
 958 Mainz (Mai), Andernach (And), and Wesel (Wes) gauge stations. Average values (Avg)
 959 calculated across all 13 gauge locations are shown in the rightmost bar of each histogram,
 960 with the standard deviations indicated by error bars.