

Dear Professor Gentine,

Please find in the document our responses. We formatted the referees' original comments in blue italic, while our responses are formatted in black. We indicate text modifications with quotation marks. Furthermore, we attached a marked-up manuscript after the responses to the reviewer.

Best regards,

Sebastian Gebler, on behalf of all authors

We appreciate the constructive comments of the reviewers on our manuscript and addressed all recommendations in the revised version of our paper.

The main changes made are:

- We added motivation to the introduction part of the revised version. More emphasis was put on scientific merit and novelty by pointing out better the differences to previous studies.
- A clarification of the link between evapotranspiration differences and grass cover height differences for the study site field on one hand and the lysimeter on the other hand was included as requested by reviewer #1.
- We pointed out that a wind-shielded precipitation device commonly decreases the underestimation of solid precipitation, but was not available for the study. Additionally, we added a precipitation comparison between lysimeter measurements and precipitation from corrected tipping bucket records according to the method of Richter (1998).
- It was also requested by reviewer #2 to add information about the cumulative drainage flux and soil water storage. This is now added in the manuscript.
- A concern of reviewer #2 was also the justification of the chosen time window for the calculation of energy balance deficit and evaporative fraction. In the revised version of the manuscript, we give additional explanation to clarify this.
- As suggested by reviewer #2, we computed evapotranspiration with the full combination equation for the lysimeters. This was done on the basis of the reconstructed lysimeter grass length. The results showed an underestimation of evapotranspiration by the Kc method and improvement of the relationships between differences in grass length (between lysimeter and field) and evapotranspiration differences for the different locations. Therefore, we replaced the empirical Kc evapotranspiration calculation with the outcome of the full-form Penman-Monteith equation in the revised manuscript.

We think that these new results underpin the scientific merit of the paper.

Please note that the recalculation of ET with the full-combination Penman-Monteith equation as well as additional comparison with wind-corrected precipitation resulted in larger modifications to the sections 2 and 3. We can only address the most relevant changes of text passages given reviewer comments. The line numbers indicate the position in the revised manuscript. The complete revisions including modified figure and table labeling, abstract and conclusions can be tracked in the attached marked-up manuscript.

Comments of anonymous referee #1

General Comments:

In the introduction part, the authors reviewed some literature on the topics of comparison between EC method and LYS method. The findings of previous literature include (1) A strong underestimation of EC-ET_a compared to LYS-ET_a is probably due to strong advection and vegetation status; (2) Errors of precipitation measurements by tipping buckets of rain gauges are caused by wind and different precipitation types (rime, dew, fog, drizzle, snow, sleet, etc.) The current study draws the similar conclusions as those finding in previous literature. Thus the novelty and scientific merit of the current paper need more justification.

We added additional motivation to the introduction part in the revised version. The relevant changes in the manuscript are:

“Moreover, measurement methods (e.g. condensation plates, optical methods) to estimate the contribution of rime, dew and fog to the total precipitation, exhibit a high uncertainty (Jacobs et al., 2006). A short term lysimeter case study by Meissner et al. (2007) and a long term investigation with a surface energy budget model calibrated with micro-lysimeters by Jacobs et al. (2006) show that rime, fog and dew contribute up to 5 % to the annual precipitation at a humid grassland site, and are usually not captured by a standard precipitation gauge.”

(line 64-70)

“In this work, a long term investigation to precipitation estimation with a lysimeter is presented. One of the points of attention in the study is the contribution of dew and rime to the total precipitation amount. The novelty compared to the work by Meissner et al. (2007) is the length of the study and the fact that a series of six lysimeters is used. Our work allows corroborating results from Jacobs et al. (2006), which used in their long term study a different, more uncertain measurement method.”

(line 111-116)

“Whereas above mentioned studies conclude that deviations between ET_a measurements are related to vegetation differences, the EC footprint and the ability to close the energy balance gap, the uncertainties of lysimeter measurements in this context are hardly investigated. Lysimeter ET_a estimations often rely on relatively low temporal resolution due to challenges in noise reduction, which impedes a simultaneous estimation of both P and ET_a, by lysimeters. Furthermore, studies with cost and maintenance intensive lysimeters are either with a few or without redundant devices, so that measurement uncertainty cannot be addressed well.”

(line 138-145)

“(4) analysed the variability of the measurements by the six lysimeters under typical field conditions with identical configuration and management.”

(line 165-167)

Minor Comments:

Table 3. Two columns should be better for presenting Sum and Mean.

We changed Table 3 as suggested by the reviewer.

Page 10, Line 12. The meaning of $S_{res,i}$ in equation (1) and $S_{dat,i}$ in equation (2) should be explained.

Thank you, we corrected this error:

“The average residual $s_{res,i}$ of measured and predicted values (Eq. 1) and the standard deviation of measured values $s_{dat,i}$ (Eq. 2) lead to the quotient B_i , which gives information about the explained variance of the fit and is related to the coefficient of determination (R^2).”
(line 246-249)

Page 12, Line 16-Line 19. “For the analysis of P and ETa, we compared the estimations of the TB and the eddy covariance method with the mean of six redundant lysimeter devices (unless specified otherwise) assuming that the lysimeter average is the most representative for estimating precipitation and actual evapotranspiration”. This sentence is confusing for readers. My understanding is that the author wants to first compare precipitation derived from lysimeter and from tipping bucket and then compare evapotranspiration derived from lysimeter and from eddy covariance method. I suggest the author to rewrite this sentence (maybe separate into 2 sentences) and clarify two objectives clearly.

The text was modified for clarification:

“In this study precipitation measured by lysimeter and TB are compared, as well as evapotranspiration measured by lysimeter and eddy covariance. The precipitation or ETa averaged over the six redundant lysimeters are used in this comparison. We assume that the lysimeter average of six redundant lysimeter devices is the most representative estimation for the lysimeter precipitation and actual evapotranspiration (unless specified otherwise).”
(line 290-295)

Page 19, Line 14-16. A comma is needed before “the relationship : : :” And a table showing the values of wind speed and the precipitation differences or a figure showing the relationship is preferred.

We corrected this and included a new figure to the manuscript showing the relationship between wind speed and precipitation differences.

Page 21, Line 1. Can the authors explain why evapotranspiration was limited by energy not by water according to the result that ETa-EC is close to ETc-FAO? The explanations on physical mechanisms should be elaborated.

For better explanation we added:

“This indicates that in general over the year 2012 evapotranspiration was limited by energy and not by water, as actual evapotranspiration was close to a theoretical maximum value for well watered conditions as estimated by ET_{PM} . This also implies that our assumption of a

stomatal resistance corresponding to well-watered conditions was justified. Water stress conditions would lead to decreased plant transpiration rates and increased stomatal resistance.” **(line 544-549)**

Page 23, Line 5. “positiv” should be “positive”.

Thank you, we corrected this error.

Fig.7. The grass height evolution trends for lysimeter field and EC station are different from July to Sep. Will this cause differences of measured evapotranspiration by the two methods and how?

With the help of the video surveillance system and the maintenance protocols we were able to reconstruct the grass length at the lysimeter. We added the grass length at the lysimeter to the figure of grass length evolution for clarity. We further added an explanation how grass lengths affect the (evapo)transpiration and cause therefore differences between the different measurement locations.

“The grass length is related to the LAI, which impacts water vapor flow at the leaf surface. Under well-watered conditions more surface for plant transpiration leads in general to higher transpiration rates by decreasing the bulk surface resistance.”
(line 592-594).

“It is assumed, on the basis of information from the video surveillance system, that grass heights generally are in good agreement between lysimeters (lysimeter site) and the surrounding field (lysimeter field), which allows a reconstruction of the grass length illustrated in Fig. 9. However, the grass harvesting dates of lysimeters and surrounding field deviate in August and September and are given for the lysimeters in Fig. 9.”
(line 596-601)

In Page 23, Line 13-16. The author mentions that the evapotranspiration differences between ET_a -EC and ET_c -LYS and grass length differences show a good correlation ($R^2=0.52$) during the period from May 24 to June 24. From Fig. 7, we can only see that the grass height evolution trend is the same from May 24 to June 24. Can the authors present a plot with the evapotranspiration difference as y-axis and grass length difference as x-axis?

We changed this according to the suggestions of the reviewer to clarify this. We added a figure to the manuscript and extended the period from May 21st to July 3rd.

“For the period from the 21st of May to the 3rd of July, a period with high grass length differences (Fig. 9) between the lysimeter site and the field behind the EC-station, ET_a differences (ET_a -EC - ET_a -LYS) and grass length differences show a good correlation ($R^2=0.58$), which is illustrated in Fig. 11.”
(line 607-611)

“The differences between ET_a -EC and ET_{PM} do not show such a significant correlation with grass heights, although the relationship in August is in correspondence with the differences of ET_a -EC and ET_a -LYS. This could be related to the EC-footprint, because the EC station is centrally located in between the two investigated fields with different grass lengths. The EC-footprint might also include other surrounding fields with different grass heights.”

(line 612-616)

In Figure 5, I would like to see the differences between P-LYS and P-TB rather than the absolute value P-LYS and P-TB.

The differences are already plotted in the figure (see upper part of figure).

Comments of anonymous referee #2

Major comments

A weighing rain gauge with wind shield (such as the Geonor one) is usually recommended to measure solid and liquid precipitation, often in conjunction with snow pillows and snow height measurements. The underestimation of solid precipitation could be decreased by this system, it should be pointed out in the document. I guess using a combination of those instruments (which are easier to install than a lysimeter and have similar measurements footprints) could lead to a difference in total rainfall of the same order as that of the total evapotranspiration. Did you try classical wind correction algorithms for raingauge systems (even if you acknowledge that the error residual do not correlate well with wind)?

The device available for our study was not equipped with a wind shield. We acknowledge this by adding a comparison of the corrected tipping bucket precipitation according to the method of Richter. Please note that we adapted the original method for hourly data to compare with the lysimeter data. The method is described in detail in the new section 2.2.4. We furthermore emphasized in the introduction and methods sections that the “standard precipitation gauge” (tipping) we used for comparison is without a wind shield and referred to literature which shows the potential measurement improvement of such a device.

“Intercomparison studies between different rain gauge designs of the World Meteorological Organization (WMO) indicated that shielded devices can considerably reduce this undercatch compared to unshielded gauges, in particular for snow and mixed precipitation (Goodison et al., 1997).”

(line 59 – 62)

“For our study, we (1) compared precipitation measurements by lysimeters and a (unshielded) standard tipping bucket device and interpreted the differences.”

(line 160 – 161)

“The unshielded gauge was temporary heated during winter time to avoid freezing of the instrument.”

(line 221 – 222)

“It was checked whether correcting the tipping bucket data (TB_{corr}) according to the method of Richter (1995) could reduce the precipitation difference between lysimeter and TB. The total precipitation sum after correction is 996.9 mm for 2012, only 3% smaller than the yearly lysimeter average and within the range of the individual lysimeters. The correction of TB data in general decreased the differences in the winter period (January – March, November - December). However, for the summer period the monthly precipitation sum of TB_{corr} mainly overestimated precipitation and tended to slightly increase the precipitation differences.”

(line 502 – 508)

“An additional comparison with corrected tipping bucket precipitation measurements according to the method of Richter (1995) shows in general a decrease of the monthly and yearly difference, which was 3 % after correction.”

(line 667 – 669)

What is the difference between the 6 lysimeters with respect with the other components of the water balance (drainage, integrated soil moisture storage) ? (it could be useful to show cumulative differences between the 6 instruments and those 2 fluxes)

We added a figure (Figure 5) showing drainage and changes in soil water storage to the revised manuscript. From drainage measurements we calculated the soil water storage term with the water balance. The changes in section 3.1 are:

“In order to further address the lysimeter uncertainty, we calculated the average cumulative drainage and soil water storage with minimum and maximum ranges for the individual lysimeters (Fig. 5). The soil water storage was determined by the remaining term of the water balance on a daily basis. The total drainage, averaged over the six lysimeters was 411.2 mm for 2012 with a variation between 385.5 and 440.4 mm. The soil moisture storage change over the year varies between -5.1 mm to 28.3 mm with an average of +11.2 mm. The assessment of drainage volumes and changes in soil water storage was somewhat hampered by erroneous data related to drainage leakage (January) or system wide shut down due to freezing. However, the uncertainty in the water balance during those periods should have a minor effect on the short term calculations of lysimeter P and ET_a .”

(line 449-458)

Minor comments

P13808: error in relating eq. 6 and 7 and the methods to derive P and ETa (lines 4-6)

We corrected this link to:

“Supposing that no evapotranspiration occurs during a precipitation event and assuming a fixed water density of 1000 kg m^{-3} , precipitation (P) [M T^{-1}] can be derived from the lysimeter water balance (Eq. 7) as:”

(line 279 – 282)

P13809L17 and L27: why 3h ? why 7 days ? Those 2 figures sounds fairly large to me, please justify; moisture status can change a lot in 7 days.

We added an explanation of the selected time windows to section 2.2.2:

“The moving window of three hours is a compromise between two sources of error. First, it guarantees a relatively small impact of random sampling errors and therefore increases the reliability of the EBD calculation. Second, the relatively short interval ensures that the calculations are not too much affected by non-stationary conditions.”

(line 324 – 327)

“Kessomkiat et al. (2013) investigated the impact of the time window on the calculation of the EF and found that a moving average over seven days gives good results, whereas a too short time window of one day gives unstable, unreliable results.”

(line 335 – 338)

P13810L7: I don't understand how EBD3h(EF) is computed.

We added some explanation below equation (10):

“The EBD is added to the uncorrected LE according to the partitioning of heat fluxes in the EF. Further details on the EBD correction method can be found in Kessomkiat et al. (2013).“
(line 344 – 346)

P13818L23: why didn't you compute Eta with the full Combination Equation instead of the empirical Kc method ? (using actual roughness length derived from vegetation height for instance, esp. for such a well known grass cover)

The Kc method is often used as a standard method for ET calculations, but probably underestimated ET for our specific grass cover conditions. We used this simple estimation of ET as the original idea of the paper was to use ET Kc solely as reference to compare with ET-EC and ET-LYS. For the revised version we considered the full-combination Penman-Monteith equation for the ET calculation. We used aerodynamic and stomatal resistance calculated on the basis of the reconstructed lysimeter grass length measurements for this approach. From our point of view the results of these calculations strengthen the conclusions regarding the role of the differences in grass height.

Therefore, we replaced the Kc-based ET estimation with the new results (ET_{PM}). The revision includes a description of the used equations for ET, aerodynamic and stomatal resistances in the methods section. New results are reported in the results section. In general, ET_{PM} was found to be slightly larger than measured ET by lysimeters and eddy covariance method.

8 Abstract

9 This study compares actual evapotranspiration (ET_a) measurements by a set of six weighable
10 lysimeters, ET_a estimates obtained with the eddy covariance (EC) method, and ~~potential crop~~
11 evapotranspiration ~~according to FAO~~ (ET_e -FAO) calculated with the full-form Penman-Monteith
12 equation (ET_{PM}) for the Rollesbroich site in the Eifel (Western Germany). The comparison of ET_a
13 measured by EC (including correction of the energy balance deficit) and by lysimeters is rarely
14 reported in literature and allows more insight into the performance of both methods. An
15 evaluation of ET_a for the two methods for the year 2012 shows a good agreement with a total
16 difference of 3.8 % (19 mm) between the ET_a estimates. The highest agreement and smallest
17 relative differences (< 8 %) on monthly basis between both methods are found in summer. ET_a
18 was close to ~~ET_e -FAO~~ ET_{PM} , indicating that ET was energy limited and not limited by water
19 availability. ET_a differences between lysimeter, ~~ET_e -FAO~~, and EC were mainly related to
20 differences in grass height caused by ~~harvesting management~~ harvest and the EC footprint. The
21 lysimeter data were also used to estimate precipitation amounts in combination with a filter
22 algorithm for high precision lysimeters recently introduced by Peters et al. (2014). The estimated
23 precipitation amounts from the lysimeter data ~~show significant differences compared to the differ~~
24 significantly from precipitation amounts recorded with a standard rain gauge at the Rollesbroich
25 test site. For the complete year 2012 the lysimeter records show a 16 % higher precipitation
26 amount than the tipping bucket. After a correction of the tipping bucket measurements by the
27 method of Richter (1995) this amount was reduced to 3 %. With the help of an on-site camera the
28 precipitation measurements of the lysimeters were analyzed in more detail. It was found that the
29 lysimeters record more precipitation than the tipping bucket in part related to the detection of
30 rime and dew, which contributes 17 % to the yearly difference between both methods. In
31 addition, fog and drizzle explain an additional 5.5 % of the total difference. Larger differences
32 are also recorded for snow and sleet situations. During snowfall, the tipping bucket device
33 underestimated precipitation severely and these situations contributed also 7.9 % to the total
34 difference. However, 36% of the total yearly difference was associated to snow cover without
35 apparent snowfall and under these conditions snow bridges and snow drift seem to explain the
36 strong ~~underestimation~~ overestimation of precipitation by the lysimeter. The remaining
37 precipitation difference (about 33 %) could not be explained, and did not show a clear relation
38 with wind speed. The ~~variations~~ variation of the individual lysimeters devices compared to the

39 | lysimeter mean ~~of 2012~~ are small showing variations up to 3 % for precipitation and 8 % for
40 | evapotranspiration.

41 1. Introduction

42 ~~Precipitation and actual evapotranspiration measurements have a quite long tradition. Precise~~
43 ~~estimates of precipitation and actual evapotranspiration are important for an improved~~
44 ~~understanding of water and energy exchange processes between land and atmosphere relevant for~~
45 ~~many scientific disciplines and agricultural management. Information about measurement errors~~
46 ~~and uncertainties is essential for improving measurement methods and correction techniques as~~
47 ~~well as for dealing with uncertainty during calibration and validation of model simulations.~~
48 Although first devices for modern scientific purposes were developed in Europe during the 17th
49 century (Kohnke et al., 1940; Strangeways, 2010). ~~However,~~ the accurate estimation of
50 precipitation (P) and actual evapotranspiration (ET_a) is still a challenge. Common precipitation
51 measurement methods exhibit systematic and random errors depending on the device locations
52 and climatic conditions. Legates and DeLiberty (1993) concluded from their long-term study of
53 precipitation biases in the United States that Hellman type gauges (US standard) undercatch
54 precipitation amounts. Undercatch is larger in case of snowfall and larger wind speeds. Wind-
55 induced loss is seen as the main source of error (Sevruk, 1981 & 1996; Yang et al., 1998;
56 Chvíla et al., 2005; Brutsaert, 2010). Precipitation gauges are commonly installed above ground
57 to avoid negative impact on the measurements by splash water, hail, and snow drift. However,
58 this common gauge setup causes wind distortion and promotes the development of eddies around
59 the device. Wind tunnel experiments with Hellman type gauges (Nešpor and Sevruk, 1999) have
60 shown precipitation losses of 2 – 10 % for rain and 20 – 50 % for snow compared to the preset
61 precipitation amount. In general, wind-induced loss increases with installation height of the
62 device and wind speed and decreases with precipitation intensity (Sevruk, 1989). Intercomparison
63 studies between different rain gauge designs of the World Meteorological Organization (WMO)
64 indicated that shielded devices can considerably reduce this undercatch compared to unshielded
65 gauges, in particular for snow and mixed precipitation (Goodison et al., 1997). Further
66 precipitation losses, which affect the rain gauge measurement, are evaporation of water from the
67 gauge surface and recording mechanisms (Sevruk, 1981; Michelson, 2004). Moreover,
68 measurement methods (e.g. condensation plates, optical methods) to estimate the contribution of
69 rime, fog dew and dew, which fog to the total precipitation, exhibit a high uncertainty (Jacobs et
70 al., 2006). A short term lysimeter case study by Meissner et al. (2007) and a long term
71 investigation with a surface energy budget model calibrated with micro-lysimeters by Jacobs et
72 al. (2006) show that rime, fog and dew contribute up to 5 % to the annual precipitation at a humid

73 | grassland site (~~Jacobs et al., 2006, Meissner et al., 2007~~), and are usually not captured by a
74 | standard precipitation gauge.

75 | The eddy covariance (EC) method is one of the most established techniques to determine the
76 | exchange of water, energy and trace gases between the land surface and the atmosphere. On the
77 | basis of the covariance between vertical wind speed and water vapor density, the EC method
78 | calculates the vertical moisture flux (and therefore ET) in high spatial and temporal resolution
79 | with relatively low operational costs. The size and shape of the measurement area (EC footprint)
80 | varies strongly with time (Finnigan, 2004). Under conditions of limited mechanical and thermal
81 | turbulence the EC method tends to underestimate fluxes (Wilson et al., 2001; Li et al., 2008).
82 | Energy balance deficits are on average found to be between 20 and 25% (Wilson et al., 2001;
83 | Hendricks Franssen et al., 2010) and therefore latent heat flux or actual evapotranspiration
84 | estimated from EC data shows potentially a strong underestimation. The energy balance closure
85 | problem can be corrected by closure procedures using the Bowen ratio. However, this is
86 | controversially discussed, especially because not only the underestimation of the land surface
87 | fluxes, but also other factors like the underestimation of energy storage in the canopy might play
88 | a role (Twine et al., 2000; Foken et al., 2011).

89 | As an alternative to classical rain gauges and the eddy covariance method, state-of-the-art high
90 | precision weighing lysimeters are able to capture the fluxes at the interface of soil, vegetation and
91 | atmosphere (Unold and Fank, 2008). A high weighing accuracy and a controlled lower boundary
92 | condition permit high temporal resolution precipitation measurements at ground level, including
93 | dew, fog, rime, and snow. Additionally, ET_a can be estimated with the help of the lysimeter water
94 | balance. However, the high acquisition and operational costs are a disadvantage of lysimeters.
95 | Moreover, the accuracy of lysimeter measurements is affected by several error sources.
96 | Differences in the thermal, wind and radiation regime between a lysimeter device and its
97 | surroundings (oasis effect) (Zenker, 2003) as well as lysimeter management (e.g., inaccuracies in
98 | biomass determination) can affect the measurements. Wind or animal induced mechanical
99 | vibrations can influence the weighing system, but can be handled by accurate data processing
100 | using filtering and smoothing algorithms (Schrader et al., 2013; Peters et al., 2014). Vaughan and
101 | Ayars (2009) examined lysimeter measurement noise for ~~minutely resolved~~ data at a temporal
102 | resolution of one minute, caused by wind loading. They presented noise reduction techniques that
103 | rely on Savitzky-Golay (Savitzky and Golay, 1964) smoothing. Schrader et al. (2013) evaluated

104 the different filter and smoothing strategies for lysimeter data processing on the basis of synthetic
105 and real measurement data. They pointed out, that the adequate filter method for lysimeter
106 measurements is still a challenge, especially at high temporal resolution, due the fact that noise of
107 lysimeter measurements varies strongly with weather conditions and mass balance dynamics.
108 Peters et al. (2014) recently introduced a filter algorithm for high precision lysimeters, which
109 combines a variable smoothing time window with a noise dependent threshold filter that accounts
110 for the factors mentioned above. They showed that their “Adaptive Window
111 and Adaptive Threshold Filter” (AWAT) improves actual evapotranspiration and precipitation
112 estimates from noisy lysimeter measurements compared to smoothing methods for lysimeter data
113 using the Savitzky-Golay filter or simple moving averages used in other lysimeter studies (e.g.,
114 Vaughan and Ayars, 2009; Huang et al., 2012; Nolz et al., 2013; Schrader et al., 2013).

~~115 The eddy covariance (EC) method is one of the most established techniques to determine the
116 exchange of water, energy and trace gases between the land surface and the atmosphere. On the
117 basis of the covariance between vertical wind speed and water vapor density, the EC method
118 calculates the vertical moisture flux (and therefore ET) in high spatial and temporal resolution
119 with relatively low operational costs. The size and shape of the measurement area (EC footprint)
120 varies strongly with time (Finnigan, 2004). Under conditions of limited mechanical and thermal
121 turbulence the EC method tends to underestimate fluxes (Wilson et al., 2001; Li et al., 2008).
122 Energy balance deficits are on average found to be between 20 and 25% (Wilson et al., 2001;
123 Hendricks Franssen et al., 2010) and therefore latent heat flux or actual evapotranspiration
124 estimated from EC data shows potentially a strong underestimation. The energy balance closure
125 problem can be corrected by closure procedures using the Bowen ratio. However, this is
126 controversially discussed, especially because not only the underestimation of the land surface
127 fluxes, but also other factors like the underestimation of energy storage in the canopy might play
128 a role (Twine et al., 2000; Foken et al., 2011).~~

129 In this work, a long term investigation to precipitation estimation with a lysimeter is presented.
130 One of the points of attention in the study is the contribution of dew and rime to the total
131 precipitation amount. The novelty compared to the work by Meissner et al. (2007) is the length of
132 the study and the fact that a series of six lysimeters is used. Our work allows corroborating results
133 from Jacobs et al. (2006), which used in their long term study a different, more uncertain
134 measurement method.

135 In the literature we find several comparisons between lysimeter measurements and standard ET
136 calculations. López-Urrea et al. (2006) found a good agreement of FAO-56 Penman-Monteith
137 with lysimeter data on an hourly basis. Vaughan et al. (2007) also reported a good accordance of
138 hourly lysimeter measurements with a Penman-Monteith approach of the California Irrigation
139 Management Information System. Wegehenkel and Gerke (2013) compared lysimeter ET with
140 reference ET and ET estimated by a numerical plant growth model. They found that lysimeter ET
141 overestimated actual ET, the cause being an oasis effect. On the other hand, also ET estimated by
142 EC measurements and water budget calculations are compared in literature. Scott (2010) found
143 that the EC-method underestimated evapotranspiration for a grassland site related to the energy
144 balance deficit. However, only a few comparisons between ET estimated by EC and lysimeter
145 data were found in literature. Chavez et al. (2009) evaluated actual evapotranspiration determined
146 by lysimeters and EC in the growing season for a cotton field site. They found a good agreement
147 of both methods after correcting the energy balance deficit and they suggested to consider also
148 the footprint area for EC calculations. Ding et al. (2010) found a lack of energy balance closure
149 and underestimation of ET_a by the EC-method for maize fields. An energy balance closure based
150 on the Bowen-Ratio method was able to reduce the ET-underestimation. Alfieri et al. (2012)
151 provided two possible explanations for a strong underestimation of EC- ET_a compared to
152 lysimeter ET_a . First, the energy balance deficit of the EC data, especially for those cases where
153 EC-measurements are affected by strong advection. Second, deviations between the vegetation
154 status of the lysimeter and the surrounding field. Evett et al. (2012) found an 18 %
155 underestimation of corrected EC- ET_a compared to ET_a estimated by lysimeter and attributed the
156 difference to differences in vegetation growth. Whereas above mentioned studies conclude that
157 deviations between ET_a measurements are related to vegetation differences, the EC footprint and
158 the ability to close the energy balance gap, the uncertainties of lysimeter measurements in this
159 context are hardly investigated. Lysimeter ET_a estimations often rely on relatively low temporal
160 resolution due to challenges in noise reduction, which impedes a simultaneous estimation of both
161 P and ET_a , by lysimeters. Furthermore, studies with cost and maintenance intensive lysimeters
162 are either with a few or without redundant devices, so that measurement uncertainty cannot be
163 addressed well.

164 The Terrestrial Environmental Observatories (TERENO) offer the possibility of detailed long-
165 term investigations of the water cycle components at a high spatio-temporal resolution (Zacharias

166 et al., 2011). This study compares precipitation and evapotranspiration estimates calculated with
167 a set of six weighing lysimeters (LYS) with nearby eddy covariance and precipitation
168 measurements for the TERENO grassland site Rollesbroich. Additional soil moisture, soil
169 temperature and meteorological measurements at this TERENO test site enable a detailed
170 analysis of differences between the different measurement techniques. The lysimeter data (ET_a -
171 LYS) are processed with the AWAT filter (Peters et al., 2014), which allows a simultaneous
172 estimation of P and ET_a in a high temporal resolution and the comparison is carried out with
173 energy balance corrected EC data (ET_a -EC). Actual ET estimates are additionally compared to
174 ~~FAO standard grass reference evapotranspiration (ET_0 -FAO) and potential crop~~
175 ~~evapotranspiration (ET_c -FAO) calculated according to the FAO crop approach for grassland full-~~
176 ~~form Penman-Monteith equation (Allen, 2000). et al., 1998) accounting for the effects of variable~~
177 grass cover height. Precipitation measurements by a classical Hellmann type tipping bucket, with
178 and without accounting for wind and evaporation induced loss (Richter correction) were
179 compared with lysimeter data for one year (2012).

180 For our study, we (1) compared precipitation measurements by lysimeters and a (unshielded)
181 standard tipping bucket device and interpreted the differences. For example, the vegetated high
182 precision lysimeters potentially allow better estimates of precipitation accounting for dew, rime
183 and fog; (2) compared eddy covariance and lysimeter ET estimates and tried to explain
184 differences in estimated values; (3) tested whether a correction of the energy balance deficit for
185 the EC-method results in an ET_a estimate which is close to the lysimeter method; (4) analysed the
186 variability of the measurements by the six lysimeters under typical field conditions with identical
187 configuration and management.

188 2. Material and Methods

189 2.1 Study Site and Measurement Setup

190 The Rollesbroich study site (50° 37' 27" N, 6° 18' 17" E) is located in the TERENO Eifel low
191 mountains range/Lower Rhine Valley Observatory (Germany). This sub-catchment of the river
192 Rur has an area of 31 ha with an altitude ranging from 474 m to 518 m a.s.l.. The vegetation of
193 the extensively managed grassland site is dominated by ryegrass and smooth meadow grass. The
194 annual mean precipitation is 1033 mm and the annual mean temperature 7.7 °C (period 1981-
195 2001); these data are obtained from a meteorological station operated by the North Rhine-
196 Westphalian State Environment Agency (LUA NRW) at a distance of 4 km from the study site.
197 **Figure** **Fig.** 1 shows a map of the study site and gives an overview of the installed measurement
198 devices.

199 In 2010 a set of six lysimeters (TERENO-SoilCan project, UMS GmbH, Munich, Germany) was
200 arranged in a hexagonal design around the centrally placed service unit, which hosts the
201 measurement equipment and data recording devices. Each lysimeter contains silty-clay soil
202 profiles from the Rollesbroich site and is covered with grass. The conditions at the lysimeters
203 therefore closely resemble the ones in the direct surroundings (Fig. 2). Additionally, the spatial
204 gap between lysimeter and surrounding soil was minimized to prevent thermal regimes which
205 differ between the lysimeter and the surrounding field (oasis effect). Every lysimeter device has a
206 surface of 1 m², a depth of 1.5 m and is equipped with a 50 l weighted leachate tank connected
207 via a bidirectional pump to a suction rake in the bottom of each lysimeter. To reproduce the field
208 soil water regime, the lower boundary conditions are controlled by tensiometers (TS1, UMS
209 GmbH, Munich, Germany) monitoring the soil matric potential inside the lysimeter bottom and
210 the surrounding field. Matric potential differences between field and lysimeter are compensated
211 by suction rakes (SIC 40, UMS GmbH, Munich, Germany) injecting leachate tank water into the
212 lysimeter monolith during capillary rise or removing water during drainage conditions. The
213 weighing precision is 100 g for the soil monolith and 10 g for the leachate tank accounting for
214 long-term temperature variations and load alternation hysteresis effects. For short term signal
215 processing the relative accuracy for accumulated mass changes of soil monolith and leachate is
216 10 g. For the year 2012 measurements were made each 5 s and averaged to get minute values. In
217 the winter season a connection between the snow lying on the lysimeter and the surrounding

218 | snow layer potentially disturbs the weighing system. ~~A snow separation system~~A mechanical
219 | vibration plate is engaged at all lysimeter devices to prevent this situation ~~by a mechanical~~
220 | ~~vibration plate, which, and~~ is activated once in 5 s between two measurements. The lysimeters
221 | are also equipped with soil moisture, matric potential and temperature sensors at different depths
222 | (10, 30, 50 and 140 cm). Amongst others, soil temperature is determined in 10, 30 and 50 cm
223 | depth with PT-100 sensors integrated in TS1-tensiometers (UMS GmbH, Munich, Germany). A
224 | schematic overview of the lysimeter device (Fig. 3) shows the ~~installing~~installation locations and
225 | the different sensor types. The lysimeter site was kept under video surveillance by a camera
226 | taking a photo of the lysimeter status every hour. Further technical specifications can be found in
227 | Unold and Fank (2008).

228 | Latent and sensible heat fluxes were measured by an eddy covariance station at a distance of
229 | approximately 30 m from the lysimeters. The EC-station (50° 37' 19" N, 6° 18' 15" E,
230 | 514 m a.s.l.) is equipped with a sonic anemometer (CSAT3, Campbell Scientific, Inc., Logan,
231 | USA) at 2.6 m height to measure wind components. The open path device of the gas analyzer
232 | (LI7500, LI-COR Inc., Lincoln, NE, USA) is mounted along with the anemometer at 2.6 m above
233 | the ground surface and measures H₂O content of the air. Air pressure is measured at the
234 | processing unit of the gas ~~analyser~~analyzer in a height of 0.57 m. Air humidity and temperature
235 | were measured by HMP45C, Vaisala Inc., Helsinki, Finland (at 2.58 m above the ground
236 | surface). Radiation was determined by a four-component net radiometer (NR01, Hukseflux
237 | Thermal Sensors, Delft, Netherlands). Soil heat flux was determined at 0.08 m depth by a pair of
238 | two HFP01 (Hukseflux Thermal Sensors, Delft, Netherlands).

239 | Precipitation measurements are made by a standard Hellmann type tipping bucket balance (TB)
240 | rain gauge (ecoTech GmbH, Bonn, Germany) with a resolution of 0.1 mm and a measurement
241 | interval of 10 minutes. The measurement altitude of 1 m above ground is in accordance with
242 | recommendations of the German weather service (DWD, 1993) for areas with an elevation
243 | > 500 m a.s.l. and occasional heavy snowfall (WMO standard is 0.5 m). The unshielded gauge
244 | was temporary heated during winter time to avoid freezing of the instrument.

245 | Additional soil moisture and soil temperature measurements were carried out with a wireless
246 | sensor network (SoilNet) installed at the study site (BogenaQu et al., 20102013). The 179 sensor
247 | locations at the Rollesbroich site contain six SPADE sensors (model 3.04, sceme.de GmbH i.G.,

248 Horn-Bad Meinberg, Germany) with two redundant sensors at 5, 20 and 50 cm depth. Further
249 technical details can be found in Qu et al. (2013). Soil water content and temperature were also
250 measured by two sensor devices installed nearby the lysimeter site.

251 2.2 Data Processing

252 2.2.1 Lysimeter

253 The lysimeter weighing data were processed in three steps:

254 1. Elimination of outliers by an automated threshold filter

255 2. Smoothing of measurement signal with the AWAT filter routine on ~~minutely basis~~the basis of
256 data at a temporal resolution of one minute

257 3. Estimation of hourly precipitation and evapotranspiration on the basis of the smoothed signal

258 Outliers were removed from the data by limiting the maximum weight difference between two
259 succeeding measurements for the soil column to 5 kg and for the leachate weight to 0.1 kg. The
260 lysimeter readings are affected by large random fluctuations caused by wind and other factors
261 that influence the measurement. Therefore, the AWAT filter (Peters et al., 2014) in a second
262 correction step was applied on the minute-wise summed leachate and on the weights for each
263 individual lysimeter. First, the AWAT routine gathers information about signal strength and data
264 noise by fitting a polynomial to each data point within an interval of 31 minutes. The optimal
265 order (k) of the polynomial is determined by testing different polynomial orders for the given
266 interval (i.e. k : 1-6) and selecting the optimal k according Akaike's information criterion (Akaike,
267 1974, Hurvich and Tsai, 1989). The maximum order of k is limited to six for the AWAT filter
268 preventing an erroneous fit caused by ~~eventual~~ outliers. ~~Measures of~~ The average residual $s_{res,i}$
269 of measured and predicted values (Eq. 1) and the standard deviation of measured values
270 $s_{dat,i}$ (Eq. 2) lead to the quotient B_i , which gives information about the explained variance of the
271 fit and is related to the coefficient of determination (R^2):

$$s_{res,i} = \sqrt{\frac{1}{r} \sum_{j=1}^r [y_j - \hat{y}_j]^2} \quad (1)$$

$$s_{dat,i} = \sqrt{\frac{1}{r} \sum_{j=1}^r [y_j - \bar{y}]^2} \quad (2)$$

$$B_i = \frac{s_{\text{res},i}}{s_{\text{dat},i}} = \sqrt{1 - R_i^2} \quad (3)$$

272 | where y_j [M] is the measured data, \hat{y}_j [M] the fitted value at each time interval-time j , \bar{y} [M] the
 273 | mean of the measurements and r the number of measurements within the given interval of data
 274 | point i . $B_i = 0$ indicates that the polynomial totally reproduces the range of data variation in
 275 | contrast to $B_i = 1$ ~~showing that~~where nothing of the variation in the data is explained by the
 276 | fitted polynomial. Second, AWAT smoothes the data using a moving average for an adaptive
 277 | window width w_i [T], which is a time dependent linear function of B_i (Eq. 4):

$$w_i(B_i) = \max(w_{\text{min}}, B_i w_{\text{max}}) \quad (4)$$

278 |
 279 | where w_{max} [T] and w_{min} [T] are maximum and minimum provided window width. For our
 280 | study w_{min} was set to 11 min, w_{max} was 61 min. A low B_i requires less smoothing and therefore
 281 | small time windows, whereas a B_i close to one requires a smoothing interval close to the
 282 | allowed w_{max} . Third, AWAT applies an adaptive threshold δ_i (Eq. 5) to the data at each time step
 283 | to distinguish between noise and signal ~~due~~related to the dynamics of mechanical disturbances:

$$\delta_i = s_{\text{res},i} \cdot t_{97.5,r} \quad \text{for } \delta_{\text{min}} < s_{\text{res},i} \cdot t_{97.5,r} < \delta_{\text{max}} \quad (5)$$

284 | where δ_i [M] is a function of the interval residuals ($s_{\text{res},i}$) [M] (see Eq. 1) and the Student t value
 285 | ($t_{97.5,r}$) for the 95 % confidence level at each time step, δ_{min} [M] is the minimum and δ_{max} [M]
 286 | is the maximum provided threshold for the mass change. The product of Student t and $s_{\text{res},i}$ is a
 287 | measure for the significance level of mass changes during flux calculation. Hence, the δ_i value
 288 | indicates the range ($\pm s_{\text{res},i} \cdot t_{97.5,r}$), where the interval data points differ not significantly from
 289 | the fitted polynomial at the 95 % confidence level. Mass changes above the adaptive threshold
 290 | δ_i are significant and interpreted as signal, whereas weight differences below δ_i are interpreted as
 291 | noise. The adaptive threshold is limited by δ_{min} and δ_{max} to guarantee that (1) mass changes
 292 | smaller than the lysimeter measurement accuracy are understood as remaining noise and
 293 | therefore not considered for the flux calculation and (2) noise is not interpreted as signal during
 294 | weather conditions, which produce noisy lysimeter readings (i.e. thunderstorms with strong wind
 295 | gusts). Lysimeter calibration tests with standard weights at the study site indicate a system scale
 296 | resolution of 0.05 kg. We chose a slightly higher threshold ($\delta_{\text{min}} = 0.055$ kg) with an adequate

297 tolerance for our TERENO lysimeter devices. For the upper threshold $\delta_{\max} = 0.24$ kg was taken,
298 similar to the example presented by Peters et al. (2014).

299 For the separation of precipitation and actual evapotranspiration (ET_a) AWAT assumes that
300 increases of ~~minutely mean~~ lysimeter and leachate weights (averaged over a period of one
301 minute) are exclusively related to precipitation and negative differences ~~are due~~ to ET_a [$M T^{-1}$].
302 Supposing that no evapotranspiration occurs during a precipitation event and assuming a fixed
303 water density of 1000 kg m^{-3} , precipitation (P) [$M T^{-1}$] can be derived from the lysimeter water
304 balance (Eq. 67) as:

$$ET_a = P - L - \frac{dS_s}{dt} \quad (6)$$

$$P = L + \frac{dS_s}{dt} \quad (7)$$

305
306 where L is the amount of leachate water [$M T^{-1}$] and dS_s/dt is the change of soil water storage
307 [$M T^{-1}$] with time. After smoothing the ~~minutely~~ fluxes at one minute resolution were cumulated
308 to hourly sums of P and ET_a .

309 Although the six lysimeters have a similar soil profile, technical configuration and management
310 (i.e. grass cut, maintenance), differences in measured values between lysimeters are not
311 exclusively related to random errors. Systematic weight variations may for example be caused by
312 soil heterogeneity, mice infestation and differences in plant dynamics. ~~For the analysis of P In~~
313 this study precipitation measured by lysimeter and ET_a we TB are compared the estimations of the
314 TB, as well as evapotranspiration measured by lysimeter and the eddy covariance method with
315 the mean. The precipitation or ET_a averaged over the six redundant lysimeters are used in this
316 comparison. We assume that the lysimeter average of six redundant lysimeter devices (unless
317 specified otherwise) assuming that the lysimeter average is the most representative estimation for
318 estimating the lysimeter precipitation and actual evapotranspiration. (unless specified otherwise).

319 2.2.2 Eddy Covariance Data

320 Eddy covariance raw measurements were taken with a frequency of 20 Hz and fluxes of sensible
321 heat (H) and latent heat (LE) were subsequently calculated for intervals of 30 minutes by using
322 the TK3.1 software package (Mauder and Foken, 2011). The complete post-processing was in
323 line with the standardized strategy for EC data calculation and quality assurance presented by
324 Mauder et al. (2013). It includes the application of site specific plausibility limits and a spike
325 removal algorithm based on median absolute deviation ~~on~~of raw measurements, a time lag
326 correction for vertical wind speed with temperature and water vapor concentration based on
327 maximizing cross-correlations between the measurements of the used sensors, a planar fit
328 coordinate rotation (Wilczak et al., 2001), corrections for high frequency spectral losses (Moore
329 1986), the conversion of sonic temperature to air temperature (Schotanus et al., 1983) and the
330 correction for density fluctuations (Webb et al., 1980). Processed half hourly fluxes and statistics
331 were applied to a three-class quality flagging scheme, based on stationarity and integral
332 turbulence tests (Foken and Wichura, 1996) and classified as high, moderate and low quality
333 data. For this analysis only high and moderate quality data were used, while low quality data
334 were treated as missing values. To assign half hourly fluxes with its source area the footprint
335 model of Korman and Meixner (2001) was applied.

336 Almost every eddy covariance site shows an unclosed energy balance, which means that the
337 available energy (net radiation minus ground heat flux) is found to be larger than the sum of the
338 turbulent fluxes (sensible plus latent heat flux) (Foken, 2008); Foken et al., 2011). In this study
339 the energy balance deficit (EBD) was determined using a 3-h moving window around the
340 measurements (Kessomkiat et al., 2013):

$$341 \quad \text{EBD}_{3\text{h}} = R_{n-3\text{h}} - (G_{3\text{h}} + \text{LE}_{3\text{h}} + H_{3\text{h}} + S_{3\text{h}}) \quad (8)$$

342

343 where $R_{n-3\text{h}}$ is average net radiation [M T^{-3}], $G_{3\text{h}}$ is average soil heat flux [M T^{-3}], $\text{LE}_{3\text{h}}$ is
344 average latent heat flux [M T^{-3}], $H_{3\text{h}}$ is average sensible heat flux [M T^{-3}], and $S_{3\text{h}}$ is average heat
345 storage (canopy air space, biomass and upper soil layer above ground heat flux plate) [M T^{-3}]. All
346 these averages are obtained over a three hour period around a particular 30 min ~~EC-~~
347 measurement. EC-measurement. The moving window of three hours is a compromise between
348 two sources of error. First, it guarantees a relatively small impact of random sampling errors and

349 therefore increases the reliability of the EBD calculation. Second, the relatively short interval
 350 ensures that the calculations are not too much affected by non-stationary conditions. It was
 351 assumed that the energy balance deficit is caused by an underestimation of the turbulent fluxes
 352 and therefore the turbulent fluxes are corrected according to the evaporative fraction. The
 353 evaporative fraction (EF) was determined for a time window of seven days:

$$EF = \frac{\overline{LE}_{7d}}{\overline{LE}_{7d} + \overline{H}_{7d}} \quad (9)$$

356 where \overline{LE}_{7d} and \overline{H}_{7d} [$M T^{-3}$] are the latent and sensible heat fluxes averaged over seven
 357 days. The chosen time period increases the reliability for EF calculation compared to single days.
 358 Dark days with small fluxes may not give meaningful results. Kessomkiat et al. (2013)
 359 investigated the impact of the time window on the calculation of the EF and found that a moving
 360 average over seven days gives good results, whereas a too short time window of one day gives
 361 unstable, unreliable results.

362 The energy balance corrected latent heat flux was determined by redistribution of the latent heat
 363 on the basis of the calculated evaporative fraction:

$$LE_{0.5h}^* = LE_{0.5h} + EBD_{3h}(EF) \quad (10)$$

366 where $LE_{0.5h}^*$ is the latent heat flux (for a certain measurement point in time; i.e. a 30
 367 minutes period for our EC data) after the correction of energy balance deficit (EBD). The EBD
 368 is added to the uncorrected LE according to the partitioning of heat fluxes in the EF. Further
 369 details on the EBD correction method can be found in Kessomkiat et al. (2013).

370 In this study, also the evapotranspiration ($ET_a - EC$) calculated with the original latent heat flux
 371 (not corrected for energy balance closure) will be presented for comparison. Furthermore, the
 372 most extreme case would be that the complete EBD is linked to an underestimation of the latent
 373 heat flux. Some authors argue (Ingwersen et al., 2011) that the EBD could be more related to
 374 underestimation of one of the two turbulent fluxes than the other turbulent flux. Therefore, as an
 375 extreme scenario the complete EBD is assigned to underestimation of the latent heat flux.

376

377 ET_a -EC is calculated from the latent heat flux according to:

378
$$ET_a = \frac{LE_h^*}{L(T_h)_{H_2O} * \rho_{H_2O}} \quad (11)$$

379

380 where ET_a is ET_a -EC [$L T^{-1}$], LE_h^* is latent heat flux [$M T^{-3}$], ρ is the density of water [$M L^{-3}$] and
381 $L(T_h)_{H_2O}$ is the vaporization energy [$L^2 T^{-2}$] at a given temperature.

382 The lysimeters are thought to be representative for the EC footprint, although size and shape of
383 the EC footprint are strongly temporally variable. However, the EC footprint is almost
384 exclusively constrained to the grassland and the lysimeters are also covered by grass.

385 2.2.3 Grass Reference Evapotranspiration

386 The measurements of ET_a by the EC-method and lysimeters were in this study compared with
 387 ~~hourly grass reference~~ evapotranspiration ~~that was~~ calculated ~~according to the single crop FAO-~~
 388 ~~method (Food and Agriculture Organization), based on the~~ with full-form Penman-Monteith
 389 equation ~~(as presented by Allen, 2000):~~ et al. (1998). This approach accounts for vegetation and
 390 ground cover conditions during crop stage considering bulk surface and aerodynamic resistances
 391 for water vapor flow. The calculations were adapted for hourly intervals according to Eq. 12:

$$\begin{aligned}
 & ET_{0_h} \\
 &= \frac{0.408\Delta(R_n - G) + \gamma \frac{37}{T_h + 273} u_2 (e^\circ(T_h) - e_a)}{\Delta + \gamma(1 + 0.34u_2)} ET_{PM} \\
 & \text{where} \quad = \frac{0.408\Delta(R_n - G) + \gamma \frac{3600\varepsilon}{T_{vh}R(r_a u_2)} u_2 (e^\circ(T_h) - e_a)}{\Delta + \gamma(1 + \frac{r_s}{r_a})} \tag{12}
 \end{aligned}$$

396 $ET_{0_h}ET_{PM}$ is the hourly ~~reference~~ Penman-Monteith evapotranspiration [$L T^{-1}$], R_n is net radiation
 397 at the grass surface [$M T^{-3}$], G is soil heat flux density [$M T^{-3}$], T_{vh} is mean hourly virtual
 398 temperature [θ], R is the specific gas constant for dry air [$L^2 T^{-2} \theta^{-1}$], r_a is the aerodynamic
 399 resistance [$T L^{-1}$], r_s is the (bulk) surface resistance [$T L^{-1}$], ε is the ratio molecular weight of
 400 water vapour (dry air) [-], T_h is mean hourly air temperature (θ), Δ slope of the saturated vapour
 401 pressure curve at T_h [$M L^{-1} T^{-2} \theta^{-1}$], γ is psychrometric constant [$M L^{-1} T^{-2} \theta^{-1}$], $e^\circ(T_h)$ is
 402 saturation vapour pressure for the given air temperature [$M L^{-1} T^{-2}$], e_a is average hourly actual
 403 vapour pressure [$M L^{-1} T^{-2}$], and u_2 is average hourly wind speed [$L T^{-1}$] at 2 m height. All
 404 required meteorological input parameters for calculating ~~the reference evapotranspiration~~ ET_{PM}
 405 were taken from the EC station. The wind speed data were corrected to ~~the~~ 2 m using the FAO-
 406 standard ~~for ET_0 calculations using the~~ wind profile relationship ~~according to~~ Allen (2000). ~~For~~
 407 ~~our ET_0 calculations we assume furthermore a fixed standard surface resistance of $70 s m^{-1}$ and a~~
 408 ~~crop height of 0.12 m.~~ et al. (1998).

409 ~~According to Allen (2000) the reference ET ($ET_e - FAO$) for a specific crop can be obtained~~
 410 ~~invoking a crop specific coefficient (K_e):~~

411 We approximated aerodynamic resistance (r_a), (bulk) surface resistance (r_s) and leaf area index (LAI)
 412 with help of grass height according to Allen et al. (2006):

413

$$\begin{aligned}
 \overline{ET_e - \text{FAO}} &= K_e \overline{ET_0} r_a \\
 &= \frac{\ln \left[\frac{z_m - \frac{2}{3} h_{plant}}{0.123 h_{plant}} \right] \ln \left[\frac{z_h - \frac{2}{3} h_{plant}}{0.1 (0.123 h_{plant})} \right]}{k^2 u_2} \quad (13)
 \end{aligned}$$

$$r_s = \frac{r_i}{\text{LAI}_{act}} \quad (14)$$

$$\text{LAI}_{act} = (0.3 \text{ LAI}) + 1.2 = 0.5 (24 h_{plant}) \quad (15)$$

414 ~~where $\overline{ET_e - \text{FAO}}$ is the hourly crop evapotranspiration [L T^{-1}] and K_e is the crop coefficient~~
 415 ~~representing the vegetation and ground cover conditions during crop stage []. For our~~
 416 ~~calculations we chose the constant rye grass hay coefficients (Allen, 2000) with different values~~
 417 ~~for the initial stage ($K_{e\text{ini}}$), the growing season ($K_{e\text{mid}}$) and late season ($K_{e\text{end}}$). The beginning~~
 418 ~~and end of the growing season were determined by using the grass length measurements (Fig. 7):~~
 419 ~~$K_{e\text{ini}}$: 0.95 (01/01/2012 — 02/03/2012), $K_{e\text{mid}}$: 1.05 (02/03/2012 — 31/10/2012), $K_{e\text{end}}$: 1.0~~
 420 ~~(01/10/2012 — 31/12/2012). $K_{e\text{mid}}$ is averaged for cutting effects due to the variable cutting~~
 421 ~~management at different study site locations. For determining ET only daytime (sunrise — sunset)~~
 422 ~~ET values were taken into account.~~

423
424
425
426
427
428
429
430
431
432
433

where z_m is the height of the wind measurement [L], z_h is the height of the humidity measurement [L], h_{plant} is the grass length [L] at the lysimeter, k is the von Karman's constant [-], r_i the stomatal resistance [$T L^{-1}$], and LAI_{act} the active leaf area index taking into account that only the upper grass surface contributes to heat and vapor transfer [-]. For our calculations we assume a fixed stomatal resistance for a well-watered grass cover of $100 s m^{-1}$ in accordance to Allen et al. (1998). The grass length at the lysimeters was estimated with the help of maintenance protocols and the surveillance system. Grass lengths between two measurement intervals were linearly interpolated on a daily basis.

434 **2.2.4 Precipitation Correction**

435 A precipitation correction according the method of Richter (1995) was applied (Eq. 16, 17) on a
436 daily basis to account for wind, evaporation and wetting losses of the tipping bucket
437 precipitation:

438
$$P^{cor} = P + \Delta P \quad (16)$$

439
$$\Delta P = bP^\epsilon \quad (17)$$

440 where P^{cor} is the corrected daily precipitation [$M T^{-1}$], P is the measured tipping bucket
441 precipitation [$M T^{-1}$], ΔP the estimated precipitation deficit [$M T^{-1}$], b the site specific wind
442 exposition coefficient [-], and ϵ the empiric precipitation type coefficient [-].

443 This correction method is widely used for German weather service stations and relies on empirical
444 relationships of precipitation type and wind exposition, without using direct wind measurements. In
445 order to determine both empirical coefficients, we categorized the precipitation type with the help
446 of air temperatures on a daily basis. It was assumed that temperatures below 0 °C result in solid
447 precipitation, temperatures between 0 °C and 4 °C give mixed precipitation and air temperatures
448 above 4 °C only liquid precipitation. Furthermore, the rain gauge is located in an open area and
449 the summer period was defined from May to September and the winter period from October to
450 April. The corresponding correction coefficients were calculated according to Richter (1995) and
451 are provided in Tab. 1.

452 3. Results and Discussion

453 3.1 Precipitation Measurements

454 Tab. 42 shows the monthly precipitation sums measured by the tipping bucket (TB) and
455 calculated from the lysimeter balance data for the year 2012. The precipitation difference
456 between both devices for the year 2012 is 145.0 mm ~~showing~~implying a 16.4 % larger average
457 lysimeter precipitation than TB. For the individual lysimeters the yearly precipitation ranges from
458 996.2 mm to 1037.7 mm (-3.0 to +1.0 % compared to the lysimeter average). This implies that
459 the minimum and maximum precipitation differences between individual lysimeters and TB were
460 114.1 mm (12.9 %) resp. 155.6 mm (17.6 %), where precipitation for lysimeters was always
461 higher than for TB. The monthly precipitation sums for the period April-October measured by the
462 tipping bucket are smaller than the ones from the lysimeter average and differences range
463 between 1 % in July and 42 % in September. The winter months show higher relative differences.
464 The highest difference was found in March 2012, when the lysimeters registered an amount of
465 precipitation double as large as the TB. The precipitation sums measured by lysimeter and tipping
466 bucket correlate well on an hourly basis, especially from April to October with R^2 varying
467 between 0.74 (Apr) and 0.99 (May), but with the exception of September (0.58). For winter
468 months the explained variance is smaller with a minimum of 13% for February 2012.

469 ~~Measured precipitation differences between individual lysimeter devices show a similar temporal~~
470 ~~pattern as differences between lysimeter and TB. Low correlations correspond with the larger~~
471 ~~differences; high correlations correspond with smaller differences.~~The period April – August
472 shows the smallest precipitation differences among the six lysimeters with monthly values of ± 5
473 % in relation to the lysimeter average. In contrast, February, September, and December exhibit
474 the highest absolute and relative precipitation differences among lysimeters with variations
475 between -13 and 13 mm (± 35 %) with respect to the mean. Fig. 4 shows the absolute daily
476 differences in precipitation between lysimeter and TB measurements. It shows that the cases
477 where lysimeters register slightly higher monthly precipitation sums than TB are related to single
478 heavy rainfall events (June, July). In contrast, especially for February, the beginning of March,
479 and the first half of December, larger fluctuations in differences between daily precipitation
480 measured by TB and lysimeter are found, with less precipitation for TB than for lysimeters most
481 of the days. These periods coincide with freezing conditions and frequent episodes with sleet or

482 snowfall. According to Nešpor and Sevruck (1999) these weather conditions are typically
483 associated with a large tipping bucket undercatch because snowflakes are easier transported with
484 the deformed wind field around a rain gauge. The surveillance system, which is installed at the
485 lysimeter site, gives support for these findings. For example, a sleet precipitation event on March
486 7th explains 70 % (8.5 mm) of the monthly precipitation difference between lysimeter and TB. At
487 this day the wind speed during the precipitation event was relatively high (4.4 m s^{-1}) and
488 precipitation intensity varied between 0.6 and 2.9 mm h^{-1} . In general, winter measurement
489 inaccuracies can be caused by frozen sensors and snow or ice deposit on the lysimeter surface.
490 This situation may cause ponding effects close to the soil surface in the lysimeter and superficial
491 runoff. In order to further address the lysimeter uncertainty, we calculated the average cumulative
492 drainage and soil water storage with minimum and maximum ranges for the individual lysimeters
493 (Fig. 5). The soil water storage was determined by the remaining term of the water balance on a
494 daily basis. The total drainage, averaged over the six lysimeters was 411.2 mm for 2012 with a
495 variation between 385.5 and 440.4 mm. The soil moisture storage change over the year varies
496 between -5.1 mm to 28.3 mm with an average of +11.2 mm. The assessment of drainage volumes
497 and changes in soil water storage was somewhat hampered by erroneous data related to drainage
498 leakage (January) or system wide shut down due to freezing. However, the uncertainty in the
499 water balance during those periods should have a minor effect on the short term calculations of
500 lysimeter P and ET_a .

501 In order to explain differences in precipitation amounts between lysimeter and tipping bucket, the
502 contribution of dew and rime to the total yearly precipitation amount was determined. The hourly
503 data of lysimeter and TB were filtered ~~using distinct~~ according meteorological criteria. First,
504 meteorological conditions- were selected which favor the formation of dew, rime, fog and mist.
505 Selected were small precipitation ~~amounts in the lysimeter data occurring before events between~~
506 sunset and sunrise and after sunset associated with high relative humidity ($> 90\%$), negative net
507 radiation and low wind speed ($< 3.5 \text{ m s}^{-1}$). Under these meteorological conditions it is probable
508 that dew or rime is formed after sunset and before sunrise on cloud free days. ~~These filter criteria~~
509 ~~also include fog and mist periods.~~ For these days the difference in precipitation between TB and
510 lysimeter is calculated if TB shows no precipitation signal or if the lysimeter has no precipitation
511 signal. For the first case ($P-TB=0$) the total amount of the lysimeter precipitation is 24.5 mm,
512 which contributes 16.9 % to the total yearly precipitation difference with the TB (and 2.4% of the

513 yearly lysimeter precipitation). The period from April to August shows in general smaller
514 precipitation amounts related to such situations. In contrast, likely dew and rime conditions where
515 lysimeter precipitation is zero have a registered amount of TB-precipitation of 1.7 mm, which is
516 only 0.2 % of the total measured TB amount for the considered period. A closer inspection of the
517 precipitation data shows that both devices are able to capture dew and rime. However, a delay of
518 some hours between TB and lysimeters was found. It is supposed that dew or fog precipitation
519 was cumulating in the TB device until the resolution threshold of 0.1 mm was exceeded. This
520 indicates that the TB resolution of 0.1 mm is too coarse to detect small dew and rime amounts in
521 a proper temporal assignment. This confirms the expected ability of the lysimeter to measure
522 rime and dew better than Hellman type pluviometers or tipping bucket devices. The surveillance
523 system was used to check whether indeed dew/rime was formed on the before-mentioned days.
524 On days which fulfilled the criteria and air temperatures close to or below 0 °C rime was seen on
525 the photos. For days that fulfilled the conditions and temperatures above 0 °C camera lenses were
526 often covered with small droplets.

527 Weather conditions with drizzle or fog occur frequently at the study site. This is related to humid
528 air masses from the Atlantic which are transported with the dominating Southwestern winds and
529 lifted against the hills in this region. The surveillance system was used to detect fog and drizzle
530 situations during the year 2012. For those situations, a difference in precipitation between TB and
531 lysimeters of 8 mm was found (~~6 mm for TB and 14 mm for LYS~~), which contributes 5.5 % to
532 the yearly difference of both devices. Fig. 56 illustrates the example of May 5 – May 6 2012. The
533 hourly photos of the site show drizzle, light rain and fog for this period. For both days the air
534 temperature is close to the dew point temperature. The precipitation difference between tipping
535 bucket and lysimeter over this period was 4.0 mm (Σ TB: 12.8 mm, Σ LYS: 16.8). The maximum
536 difference was 0.5 mm and found at ~~6h~~ on the 5th of May in combination with fog. On May 5
537 during these conditions hourly TB precipitation is often zero and LYS mean precipitation rates
538 are small (0.02 - 0.2 mm hr⁻¹). The comparison of individual lysimeter devices shows that not
539 every lysimeter exceeds the predefined lower threshold of 0.055 mm for the AWAT filter (i.e. 5th
540 of May 15:00, 6th of May 01:00- 03:00 LT). However, in these cases at least three lysimeters
541 show a weight increase, which supports the assumption that a real signal was measured instead of
542 noise.

543 With the purpose of explaining the remaining difference in precipitation amount between TB and
544 lysimeter, the relationship between wind speed and the precipitation differences was examined.
545 ~~Although~~ The determined precipitation differences could in theory be explained by undercatch
546 related to wind (Sevruk, 1981 & 1996), ~~a general correlation between wind speed and~~
547 ~~precipitation residuals was not found ($R^2=0.02$).~~ It was checked whether correcting the tipping
548 bucket data (TB_{corr}) according to the method of Richter (1995) could reduce the precipitation
549 difference between lysimeter and TB. The total precipitation sum after correction is 996.9 mm for
550 2012, only 3% smaller than the yearly lysimeter average and within the range of the individual
551 lysimeters. The correction of TB data in general decreased the differences in the winter period
552 (January – March, November - December). However, for the summer period the monthly
553 precipitation sum of TB_{corr} mainly overestimated precipitation and tended to slightly increase the
554 precipitation differences. In order to explore this relation further we examined the correlation
555 between wind speed and precipitation residuals and found almost no correlation (Fig. 7). A
556 possible explanation is that other potential dew or rime situations are not properly filtered by the
557 used criteria (e.g. dew occurs in case the net radiation is slightly positive or close to zero).
558 Additionally, the correlation between undercatch and wind speed is dependent on precipitation
559 type, intensity and drop size, for which information was limited during the investigation period.
560 To investigate these relations we ~~classified~~ used the classification of precipitation type with the
561 help of air temperatures assuming that temperatures below 0 °C result in solid precipitation and
562 above 4 °C only liquid precipitation occur ~~types as outlined before~~. The contribution of liquid
563 precipitation to total yearly precipitation is 80.9 % for the TB and 74.7 % for the lysimeters. The
564 relative amount of solid precipitation was also different between the two measurement methods.
565 Whereas for the lysimeters 7.8 % (79.7 mm) was classified as solid precipitation, the TB had only
566 0.6 % (5.6 mm) during periods with temperature < 0 °C. In relation to the total precipitation
567 difference of 145 mm this means that 51 % of the difference was associated with solid
568 precipitation events and 37 % with liquid precipitation events, which indicates the relatively large
569 contribution of solid precipitation events to the total difference. The transition range (0-4 °C)
570 makes up 12 % of the total difference. Moreover, it was found that 78.7 % of the solid
571 precipitation ~~came~~ came along with small precipitation intensities (< 1.0 mm h⁻¹) and low wind
572 speeds (< 2.0 m s⁻¹). The surveillance system allowed to further investigate these large
573 precipitation differences for air temperatures below zero. The snow depth at the lysimeters and
574 surrounding areas is also an indication of precipitation amounts, assuming that 1 cm snow height

575 corresponds to 1 mm precipitation. This method revealed that for conditions of light to moderate
576 snowfall ($< 4 \text{ mm h}^{-1}$ precipitation intensity) the TB had a precipitation undercatch ~~during winter~~
577 ~~weather conditions~~ in January, February and December of 11.4 mm (7.9 % of total precipitation
578 difference). The registered precipitation amount of the lysimeter under those conditions was
579 realistic. However, during periods where the lysimeters were completely covered by snow (e.g. 1
580 – 15 February) precipitation estimates by lysimeter (up to 16 mm d^{-1} difference with tipping
581 bucket) could not be confirmed by the camera system and were most probably influenced by
582 snow drift or snow bridges. These situations explain 35.8 % (51.9 mm) of the total precipitation
583 difference for 2012. For solid precipitation events a relationship ($R^2=0.5$) between precipitation
584 differences and wind speed was found, but the number of datapoints was very limited ($n=7$). For
585 conditions of liquid precipitation no correlation was found between residuals and wind speed
586 ($R^2<0.02$).

587

588 3.2 Comparison of Evapotranspiration

589 In general, the yearly sums of ~~ET_a-EC~~ ET_{PM} and ET_a-LYS were slightly higher than ~~ET_e-FAO~~ ;
590 ~~ET_a-EC~~ ; ~~6.1.6~~ % for ~~ET_a-EC~~ ET_{PM} and ~~5.62.4~~ % for ET_a-LYS . The minimum ET_a of the
591 individual lysimeter measurements (ET_a-LYS_{min}) is 467.1 mm, which is 7.9 % smaller than the
592 lysimeter average (507.4 mm); the maximum (ET_a-LYS_{max}) is 523.1 mm (+ 3.1 %). ~~ET_a-EC is~~
593 ~~close to the calculated ET_e-FAO .~~ This indicates that in general over the year 2012
594 evapotranspiration was limited by energy and not by water, as actual evapotranspiration was
595 close to a theoretical maximum value for well watered conditions as estimated by ET_{PM} . This also
596 implies that our assumption of a stomatal resistance corresponding to well-watered conditions
597 was justified. Water stress conditions would lead to decreased plant transpiration rates and
598 increased stomatal resistance. Tab. 3 lists the evapotranspiration results of January – December
599 2012. ~~For the period from April to August the monthly evapotranspiration sums calculated from~~
600 ~~hourly lysimeter data (In 2012 ET_{PM} was always close to ET_a-LYS) and eddy covariance data~~
601 ~~(ET_a-EC) and there are no months that ET_{PM} is clearly higher/larger than the calculated~~
602 ~~FAO measured actual evapotranspiration (ET_e-FAO), confirming that in these months~~
603 ~~evapotranspiration was not limited by soil moisture content, but energy. However, for May, June~~
604 ~~and July ET_e-FAO and ET_a-EC are within the range of the individual ET_a-LYS . In contrast,~~
605 ~~March and November exhibit smaller monthly sums of ET_a-LYS and ET_a-EC compared to ET_e-~~
606 ~~FAO by lysimeter and eddy covariance.~~ Root mean square errors of hourly ET_a sums vary
607 between 0.01 mm h^{-1} in winter and 0.11 mm h^{-1} in summer months and are in phase with the
608 seasonal ET dynamics.

609 We focus now on the comparison of monthly ET_a-LYS and ET_a-EC sums within the investigated
610 period. During winter periods with low air temperatures and snowfall ET_a-LYS and ET_a-EC
611 showed larger relative differences. For the period March to May ET_a-LYS and ET_a-EC differ
612 approx. 6 % and ET_a-LYS exceeds ET_a-EC from June to August by 12 %. The larger difference
613 in August (23 %) explains the yearly difference between ET_a-EC and ET_a-LYS . Hourly actual
614 evapotranspiration from lysimeter and hourly actual evapotranspiration from EC are strongly
615 correlated, but correlation is lower in the winter months. The registered monthly ET by the
616 different lysimeters shows the largest variations in July with amounts that are up to 14.0 mm
617 lower and 8.0 mm higher than the ET averaged over all six lysimeters.

618 Fig. 68 shows the cumulative curve of the daily ET_a -LYS and ET_a -EC compared to ~~ET_e -~~
619 ~~$FAOET_{PM}$~~ for 2012. From end of March 2012 the sums of ET_a -LYS and ET_a -EC tend to
620 converge, but at the end of May ET_a -EC exceeds ET_a -LYS. In June and July ET_a -LYS and ET_a -
621 EC are very similar, but in August ET_a -LYS is larger than ET_a -EC. After August the difference
622 between ET_a -LYS and ET_a -EC does not increase further. The area in grey represents the range of
623 minimum and maximum cumulative ET_a -LYS, measured by individual lysimeters. Until August
624 ET_a -EC and ~~ET_e - $FAOET_{PM}$~~ are slightly higher or close to the maximum measured ET_a -LYS.
625 ~~Later in August ET_{PM} increases further, whereas ET_a -EC is close to the lower limit and ET_e - FAO~~
626 ~~falls below the minimum lysimeter value.~~ Additionally, Fig. 68 shows the course of the ET_a -EC
627 without correction for EBD and for ~~an extreme correction (ET_a -EC max.) where all EBD is~~
628 ~~attributed to underestimation of the latent heat flux.:~~ ET_a -uncorr is ca. 411 mm over this period,
629 whereas ET_a -EC max is 567 mm, which shows the large potential uncertainty of the EC-data. The
630 comparison illustrates that the application of the Bowen ratio correction to the EC data results in
631 an actual evapotranspiration estimate close to the actual evapotranspiration from the lysimeter,
632 whereas ET_a -EC uncorr is much smaller than the lysimeter evapotranspiration. Tab. 4 lists the
633 monthly latent heat fluxes, the corrected LE fluxes (on the basis of the Bowen ratio) and the
634 mean differences between both. It was found that the absolute difference is between 29.8 W m^{-2}
635 (August 2012) and 3.2 W m^{-2} (February 2012). The EBD ranges from 12.6 % - 24.2 % for the
636 period April to September. The yearly maximum was found in February with 36.9 %. EB deficits
637 are site-specific, but these findings confirm the importance of EC data correction as suggested by
638 Chavez et al. (2009).

639 In order to explain the differences between ~~ET_e - $FAOET_{PM}$~~ , ET_a -EC and ET_a -LYS, we
640 investigated the variations in radiation, vegetation and temperature regime and their impact on
641 ET in more detail. The albedo could be estimated according to the measured outgoing shortwave
642 radiation at the EC-station divided by the incoming shortwave radiation, also measured at the EC-
643 station. The yearly mean albedo is 0.228, which is close to the assumed albedo of 0.23 for
644 grassland. However, some periods (i.e. periods with snow cover) have a much higher albedo.
645 ~~Albedo~~ Although albedo variations between different vegetation growth stages at different fields
646 at the study site were considered as explanation for differences ~~cannot explain the fact that~~
647 ~~reference ET is smaller than~~ in ET_a , we assume similar albedo for ET_a -EC and ET_a -LYS

648 measurement due to the central location of of the radiation measurements between the relevant
649 fields.

650 ~~Hence, we examined the effects of vegetation growth with the help of grass length. Fig. 7~~
651 grass length is related to the LAI, which impacts water vapor flow at the leaf surface. Under well-
652 watered conditions more surface for plant transpiration leads in general to higher transpiration
653 rates by decreasing the bulk surface resistance. Fig. 9 shows that the grass length measured at the
654 Rollesbroich site is up to 80 cm before cutting. Unfortunately, grass height measurements are not
655 available for the lysimeters but only for the surrounding field. It is assumed, on the basis of
656 information from the video surveillance system, that grass heights generally are in good
657 agreement between lysimeters (lysimeter site) and the surrounding field (lysimeter field-), which
658 allows a reconstruction of the grass length illustrated in Fig. 9. However, the grass harvesting
659 dates of lysimeters and surrounding field deviate in August and September and are given for the
660 lysimeters in Fig. 7-9.

661 Fig. 810 illustrates the differences of the measured daily ET_a sums between lysimeter and EC.
662 High positive and negative differences up to 2.1 mm/day were found from March 2012 –
663 September 2012. In general, the differences of ET_a -~~EC~~LYS and ET_e -FAO~~ET~~PM show smaller
664 fluctuations than the differences of ET_a -LYSEC and ET_e -FAO~~ET~~PM. It ~~is~~was found that lysimeter
665 harvesting affects the ~~ET~~differences between ET_a -LYS and ET_e -FAO/~~ET~~PM/ ET_a -EC. The
666 differences were ~~positiv~~positive before harvesting and negative after harvesting indicating ET_a
667 reduction due to the grass cutting effects. ~~For the period from May 21 to July 3, grass lengths~~
668 ~~were estimated and linearly interpolated on a daily basis. For this period grass length at the~~
669 ~~lysimeter site and ET_a differences between ET_a -LYS and ET_e -FAO correlate well ($R^2=0.50$).~~
670 ~~These results reflect the discrepancy in ET estimated on the basis of ET_e -FAO calculations with~~
671 ~~constant K_c and actual ET under conditions of a higher grass height. ET_a differences caused by~~
672 ~~variations in grass length are also found for the comparison of ET_a -EC with ET_a -LYS. For the~~
673 ~~period from the 24th of May to the 24th of June, a period with high grass length differences~~
674 ~~(Fig. 7)For the period from the 21st of May to the 3rd of July, a period with high grass length~~
675 differences (Fig. 9) between the lysimeter site and the field behind the EC-station, ET_a
676 differences (ET_a -EC - ET_a -LYS) and grass length differences show a good correlation
677 ($R^2=0.52$)-58), which is illustrated in Fig. 11. During the period with maximum grass length
678 difference (24 May – 1 June) ET_a -EC is 26 % higher than ET_a -LYS. The differences between

679 ET_a -EC and ET_e -FAO ET_{PM} do not show such a significant correlation with grass heights,
680 although the relationship in August is in correspondence with the differences of ET_a -EC and ET_a -
681 LYS. This could be related to the EC-footprint ~~which~~, because the EC station is centrally located
682 in between the two investigated fields with different grass lengths. The EC-footprint might also
683 include other surrounding fields with different grass heights. 80 % of the EC footprint is located
684 within a radius of 100 m of the EC tower, and 70 % in a radius of 40 m, which is the approximate
685 lysimeter distance. Therefore, the ET_a -EC estimations represent a spatial mean of a wider area,
686 where cutting effects are averaged compared to the lysimeter point measurements. Fig. 912
687 shows the mean hourly ET_a rates of lysimeter and EC as well as the FAO-reference ET_{PM} for
688 2012. In general, the daily courses and the daily maxima of ET_a -LYS, ET_e -FAO ET_{PM} and ET_a -
689 EC correspond well. ET_a -EC shows higher peaks at noon in May and September compared to
690 ET_a -LYS ~~and ET_e -FAO~~, but corresponds well to ET_{PM} . In contrast, ET_a -LYS exhibits the highest
691 rates from June to August. The absence of a harvest of the lysimeter in August and the first
692 September decade (in contrast to the surrounding fields) leads to potentially increased lysimeter
693 ET_a measurements as compared to the surroundings due to an island position.

694 ~~The grass length affects the K_e value, but differences between the reference evapotranspiration~~
695 ~~and measured actual evapotranspiration can also be related to the weather conditions. Nolz and~~
696 ~~Cepuder (2013) showed that K_e values of 1.1–1.5 are likely for grassland after rain events (i.e.~~
697 ~~June, July) and high soil moisture conditions.~~

698 In order to examine whether lysimeter measurements could have been affected by a soil
699 temperature regime different from the field, the temperature regimes of the lysimeters were
700 compared to the field temperature. Fig. 1013 shows the daily mean soil temperature differences
701 between the lysimeters, a nearby SoilNet device (SN 30) and the mean of all available SoilNet
702 devices installed at the southern study site. SoilNet temperatures were measured 5 cm below
703 surface; lysimeter temperature measurements were conducted with SIS sensors in 10 cm depth.
704 The temperature differences between the lysimeter and the nearby SoilNet device and the SoilNet
705 mean are less than 1 K, which is as well the range of variation of the SoilNet device with respect
706 to the SoilNet mean. In general the temperature differences increase until noon and then decrease
707 again. Positive differences from May to July indicate warmer/higher lysimeter soil temperatures
708 than the surroundings. However, a clear indicator for a bias caused by an oasis effect in the
709 lysimeter measurements was not found. Feldhake and Boyer (1986) describe the effect of soil

710 temperature on evapotranspiration for different grass types, which allow an estimation of ET_a
711 increase caused by a differing lysimeter temperature regime. They showed that daily ET_a rates
712 can increase with an increase of soil temperature (i.e. daily Bermuda grass ET_a rate increases
713 from 4.3 mm/day to 6.4 mm/day (49 %) for a soil temperature increase from 13 to 29 °C). We
714 used this linear relationship to roughly estimate the effect on ET_a for the period May – August on
715 a daily basis. For this period the measured soil temperature with SN(30) for daylight hours
716 ranged between 9.5 and 15.1 °C and between 9.3 and 15.5 °C for the lysimeter mean (SIS
717 sensors). The mean difference is 0.67 K. This results in a total ET_a increase of 8.8 mm or 2.5 %
718 in relation to the total ET_a -LYS of 349 mm on the basis of hourly ET. Therefore, the effect of
719 increased soil temperature in the lysimeter is most probably limited, but not negligible.

720 4. Conclusions

721 This study compares evapotranspiration and precipitation estimates calculated using a set of six
722 redundant weighable lysimeters with nearby eddy covariance and precipitation measurements at a
723 TERENO grass land site in the Eifel (Germany) for one year (2012). The ~~minutely resolved~~
724 lysimeter data at a temporal resolution of one minute are processed with the AWAT filter (Peters
725 et al., 2014), which takes account of the lysimeter noise due to random fluctuations caused by
726 changing weather conditions. Additional precipitation measurements were conducted with a
727 classical unshielded Hellmann type tipping bucket and compared with lysimeter data. For the ET_a
728 comparison eddy covariance (EC) data is corrected for the energy balance deficit using the
729 Bowen ratio method. ~~FAO standard grass reference~~ Additionally, evapotranspiration corrected for
730 grass height variations (ET_e -FAO) was and the evapotranspiration according the full-form
731 Penman-Monteith equation were calculated ~~according to the FAO crop approach for grassland~~
732 ~~(Allen, 2000).~~

733 The estimated hourly precipitation amounts derived by lysimeter and tipping bucket data show
734 significant differences and the total precipitation measured by the lysimeter is 16.4 % larger than
735 the tipping bucket amount. The relative differences in the monthly precipitation sums are small in
736 the summer period, whereas high differences are found during the winter season. The winter
737 months with snowsolid precipitation exhibit the lowest correlations between lysimeter and
738 tipping bucket amounts. Precipitation was measured by six different lysimeters and yearly
739 amounts for individual lysimeters showed variations of -3.0 to 1.0 % compared to the yearly
740 precipitation mean over all lysimeters. An additional comparison with corrected tipping bucket
741 precipitation measurements according to the method of Richter (1995) shows in general a
742 decrease of the monthly and yearly difference, which was 3 % after correction. In order to
743 explain the differences in precipitation between the devices the contribution of dew, rime and fog
744 to the yearly precipitation was analyzed. This was done by filtering the data for typical weather
745 conditions like high relative humidity, low wind speed and negative net radiation which promote
746 the development of dew and rime. For the identified cases a check was made with a visual
747 surveillance system whether dew/rime was visible. During these conditions the lysimeter shows
748 clearly larger precipitation amounts than the TB, which explains 16.9 % of the yearly
749 precipitation difference. Fog and drizzling rain conditions, additionally identified with the help of
750 the on-site camera system, explain another 5.5 % of the yearly precipitation differences. These

751 findings indicate an improved ability of the lysimeters to measure dew and rime as well as fog
752 and drizzling rain. The remaining 78 % of the precipitation difference between lysimeters and
753 tipping bucket is strongly related to snowfall events, as under those conditions large differences
754 were found. Lysimeter precipitation measurements are affected by a relatively high measurement
755 uncertainty during winter weather conditions similar to TB and other common measurement
756 methods. Thus, the limitations for the lysimeter precipitation measurements during those periods
757 need further investigation. We found that during conditions where the lysimeters were completely
758 covered by snow, lysimeter records were unreliable, and contributed to 36 % of the total
759 precipitation difference.

760 Actual evapotranspiration measured by the eddy covariance method (ET_a -EC) and lysimeter
761 (ET_a -LYS) showed a good correspondence for 2012, with larger relative differences and low
762 correlations in winter in contrast to high correlations and smaller relative differences in summer.
763 The variability of ET_a of the individual lysimeters in relation to the lysimeter average was -7.9 to
764 3.1 % in 2012 with larger absolute differences in summer. Both ET_a -EC and ET_a -LYS₇ were
765 close to the calculated ~~crop-reference~~Penman-Monteith evapotranspiration (ET_e -FAO ET_{PM}),
766 which indicates that evapotranspiration at the site was ~~notenergy~~ limited ~~by soil moisture, but by~~
767 ~~energy~~. The differences between ET_a -LYS, ET_a -EC and ET_e -FAO ET_{PM} were mainly related to
768 harvesting management at the study site. A relationship between grass length at the lysimeter and
769 differences between ET_e -FAO ET_{PM} and ET_a -LYS was found. Variable grass cutting dates for
770 different fields around the EC-station and the lysimeter harvest lead to differences in actual
771 evapotranspiration up to 2.1 mm day⁻¹ for periods with larger grass length discrepancies.

772 The correction of the energy balance deficit with the Bowen ratio method resulted in ET_a -EC
773 which was close to ET_a -LYS. If the correction was not applied, ET_a -~~EC~~ was 16 % smaller than
774 for the case where it was applied. In contrast, if the EB-deficit was completely attributed to the
775 latent heat flux ET_a was 15.7 % larger than for the default case. These results point to the
776 importance of adequate EC data correction.

777 **Acknowledgements**

778 This research is based on data provided by the research infrastructures of TERENO and
779 TERENO-SoilCan. We thank the Transregio32 for contributing data from the Rollesbroich study
780 site and want to acknowledge H. Rützel, W. Benders, F. Engels, L. Fürst, W. Küppers, D. Dolfus,
781 and M. Kettler accounting for the realization and maintenance of the research facilities. We also
782 thank Andre Peters for providing the AWAT software. We further thank the "Arbeitskreis
783 Lysimeterdatenauswertung" for the stimulating discussions.

784 **References**

785 Akaike, H.: A new look at statistical model identification, *IEEE Transactions on Automatic*
786 *Control*, 19, 716–723, doi:10.1109/TAC.1974.1100705, 1974.

787 Alfieri, J. G., Kustas, W. P., Prueger, J. H., Hipps, L. E., Evett, S. R., Basara, J. B., Neale, C. M.
788 U., French, A. N., Colaizzi, P., Agam, N., Cosh, M. H., Chavez, J. L., and Howell, T. A.: On the
789 discrepancy between eddy covariance and lysimetry-based surface flux measurements under
790 strongly advective conditions, *Advances in Water Resources*, 50, 62-78,
791 doi:10.1016/j.advwatres.2012.07.008, 2012.

792 Allen, R. G.: ~~Using the FAO 56 dual crop coefficient method over an irrigated region as part of~~
793 ~~an evapotranspiration intercomparison study, *Journal of Hydrology*, 229, 27-41,~~
794 ~~doi:10.1016/S0022-1694(99)00194-8, 2000.~~, Pereira, L. S., Raes, D., Smith, M.: *Crop*
795 evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and
796 drainage paper 56. FAO, Rome, 9, 300, 1998.

797 ~~Bogena, H. R., Herbst, M., Huisman, J. A., Rosenbaum, U., Weuthen, A., and Vereecken, H.:~~
798 ~~Potential of Wireless Sensor Networks for Measuring Soil Water Content Variability, *Vadose*~~
799 ~~Zone Journal~~, 9, 1002-1013, doi:10.2136/Vzj2009.0173, 2010.

800 Allen, R.G., Pruitt, W. O., Wright, J. L., Howell, T. A., Ventura, F., Snyder, R., Itenfisu, D.,
801 Steduto, P., Berengena, J., Yrisarry, J. B., Smith, M., Pereira, L. S., Raes, D., Perrier, A., Alves,
802 I., Walter, I., Elliott, R.: A recommendation on standardized surface resistance for hourly
803 calculation of reference ETo by the FAO56 Penman-Monteith method. *Agricultural Water*
804 Management, 81, 1-22, doi: 10.1016/j.agwat.2005.03.007, 2006.

805 Brutsaert, W.: *Hydrology: An Introduction* / Wilfried Brutsaert, 5th print. ed., Univ. Press,
806 Cambridge, XI, 605 S. pp., 2010.

807 Chávez, J., Howell, T., and Copeland, K.: Evaluating eddy covariance cotton ET measurements
808 in an advective environment with large weighing lysimeters, *Irrigation Science*, 28, 35-50,
809 doi:10.1007/s00271-009-0179-7, 2009.

810 Chvíla, B., Sevrúk, B., and Ondrás, M.: The wind-induced loss of thunderstorm precipitation
811 measurements, *Atmospheric Research*, 77, 29-38, doi:10.1016/j.atmosres.2004.11.032, 2005.

812 Deutscher Wetterdienst (DWD): Richtlinie für automatische Klimastationen, Offenbach am
813 Main, 1993.

814 Ding, R., Kang, S., Li, F., Zhang, Y., Tong, L., and Sun, Q.: Evaluating eddy covariance method
815 by large-scale weighing lysimeter in a maize field of northwest China, *Agricultural Water*
816 *Management*, 98, 87-95, doi:10.1016/j.agwat.2010.08.001, 2010.

817 Evett, S. R., Schwartz, R. C., Howell, T. A., Louis Baumhardt, R., and Copeland, K. S.: Can
818 weighing lysimeter ET represent surrounding field ET well enough to test flux station
819 measurements of daily and sub-daily ET?, *Advances in Water Resources*, 50, 79-90,
820 doi:10.1016/j.advwatres.2012.07.023, 2012.

821 Feldhake, C. M. and Boyer, D. G.: Effect of soil temperature on evapotranspiration by C3 and C4
822 grasses, *Agricultural and Forest Meteorology*, 37, 309-318, doi:10.1016/0168-1923(86)90068-7,
823 1986.

824 Finnigan, J.: The footprint concept in complex terrain, *Agricultural and Forest Meteorology*, 127,
825 117-129, doi:10.1016/j.agrformet.2004.07.008, 2004.

826 Foken, T. and Wichura, B.: Tools for quality assessment of surface-based flux measurements,
827 *Agricultural and Forest Meteorology*, 78, 83-105, doi:10.1016/0168-1923(95)02248-1, 1996.

828 Foken, T.: The energy balance closure problem: An overview, *Ecological Applications*, 18, 1351-
829 1367, doi:10.1890/06-0922.1, 2008.

830 Foken, T., Aubinet, M., Finnigan, J. J., Leclerc, M. Y., Mauder, M., and Paw U, K. T.: Results Of
831 A Panel Discussion About The Energy Balance Closure Correction For Trace Gases, *Bulletin of*
832 *the American Meteorological Society*, 92, 13-18, doi:10.1175/2011bams3130.1, 2011.

833 [Goodison, B. E., Louie, P. Y. T., and Yang, D.: The WMO solid precipitation measurement](#)
834 [intercomparison. World Meteorological Organization-Publications-WMO TD, 65-70, 1997.](#)

835 Hendricks Franssen, H. J., R. Stöckli, I. Lehner, E. Rotenberg, and S. I. Seneviratne.: Energy
836 Balance Closure of Eddy-Covariance Data: A Multisite Analysis for European Fluxnet Stations,
837 *Agricultural and Forest Meteorology*, 150(12), 1553-1567, doi:10.1016/j.agrformet.2010.08.005,
838 2010.

839 Huang, W., Zhang, C., Xue, X., and Chen, L.: A Data Acquisition System Based on Outlier
840 Detection Method for Weighing Lysimeters, in: *Computer and Computing Technologies in*
841 *Agriculture V*, edited by: Li, D., and Chen, Y., IFIP Advances in Information and
842 *Communication Technology*, Springer Berlin Heidelberg, 471-478, 2012.

843 Hurvich, C. and Tsai, C.: Regression and time series model selection in small samples,
844 *Biometrika*, 76, 297–307, doi:10.1093/biomet/76.2.297, 1989.

845 Ingwersen, J., Steffens, K., Högy, P., Warrach-Sagi, K., Zhunusbayeva, D., Poltoradnev, M.,
846 Gäbler, R., Wizemann, H. D., Fangmeier, A., Wulfmeyer, V., and Streck, T.: Comparison of
847 Noah simulations with eddy covariance and soil water measurements at a winter wheat stand,
848 *Agricultural and Forest Meteorology*, 151, 345-355, doi: 10.1016/j.agrformet.2010.11.010, 2011.

849 Jacobs, A. F. G., Heusinkveld, B. G., Wichink Kruit, R. J., and Berkowicz, S. M.: Contribution of
850 dew to the water budget of a grassland area in the Netherlands, *Water Resources Research*, 42,
851 W03415, doi:10.1029/2005wr004055, 2006.

852 Kessomkiat, W., Hendricks Franssen, H. J., Graf, A., and Vereecken, H.: Estimating random
853 errors of eddy covariance data: An extended two-tower approach, *Agricultural and Forest*
854 *Meteorology*, 171–172, 203-219, doi:10.1016/j.agrformet.2012.11.019, 2013.

855 Kohnke, H., Davidson, J. M., and Dreibelbis, F. R.: A survey and discussion of lysimeters and a
856 bibliography on their construction and performance, U. S. Govt. print. off., Washington, D.C., 68
857 p., 1940.

858 Kormann, R. and Meixner, F.: An Analytical Footprint Model For Non-Neutral Stratification,
859 *Bound-Lay Meteorology*, 99, 207-224, doi:10.1023/a:1018991015119, 2001.

860 Legates, D. R. and DeLiberty, T. L.: Precipitation measurement biases in the United States,
861 *JAWRA Journal of the American Water Resources Association*, 29, 855-861, doi:10.1111/j.1752-
862 1688.1993.tb03245.x, 1993.

863 Li, S., Kang, S., Zhang, L., Li, F., Zhu, Z., and Zhang, B.: A comparison of three methods for
864 determining vineyard evapotranspiration in the arid desert regions of northwest China,
865 *Hydrological Processes*, 22, 4554-4564, doi:10.1002/hyp.7059, 2008.

866 López-Urrea, R., Olalla, F. M. d. S., Fabeiro, C., and Moratalla, A.: An evaluation of two hourly
867 reference evapotranspiration equations for semiarid conditions, *Agricultural Water Management*,
868 86, 277-282, doi:10.1016/j.agwat.2006.05.017, 2006.

869 Mauder, M. and Foken, T.: Documentation and Instruction Manual of the Eddy-Covariance
870 Software Package TK3. Arbeitsergebnisse / Universität Bayreuth, Abteilung Mikrometeorologie,
871 2011.

872 Mauder, M., Cuntz, M., Drüe, C., Graf, A., Rebmann, C., Schmid, H. P., Schmidt, M., and
873 Steinbrecher, R.: A strategy for quality and uncertainty assessment of long-term eddy-covariance
874 measurements, *Agricultural and Forest Meteorology*, 169, 122-135,
875 doi:10.1016/j.agrformet.2012.09.006, 2013.

876 Meissner, R., Seeger, J., Rupp, H., Seyfarth, M., and Borg, H.: Measurement of dew, fog, and
877 rime with a high-precision gravitation lysimeter, *Journal of Plant Nutrition and Soil Science*, 170,
878 335-344, doi:10.1002/jpln.200625002, 2007.

879 Michelson, D. B.: Systematic correction of precipitation gauge observations using analyzed
880 meteorological variables, *Journal of Hydrology*, 290, 161-177,
881 doi:10.1016/j.jhydrol.2003.10.005, 2004.

882 Moore, C.J.: Frequency response corrections for eddy correlation systems. *Boundary-Layer*
883 *Meteorology*, 37, 17-35, doi: 10.1007/bf00122754, 1986.

884 Nešpor, V. and Sevruk, B.: Estimation of Wind-Induced Error of Rainfall Gauge Measurements
885 Using a Numerical Simulation, *Journal of Atmospheric and Oceanic Technology*, 16, 450-464,
886 10.1175/1520-0426(1999)016<0450:eowieo>2.0.co;2, 1999.

887 Nolz, R., Kammerer, G., and Cepuder, P.: Interpretation of lysimeter weighing data affected by
888 wind, *Journal of Plant Nutrition and Soil Science*, 176, 200–208, doi:10.1002/jpln.201200342,
889 2013.

890 ~~Nolz, R. and Cepuder, P.: Verdunstung am Beginn und am Ende der Vegetationsperiode.~~
891 ~~15. Gumpensteiner Lysimetertagung 2013. In: Lehr- und Forschungszentrum für Landwirtschaft~~
892 ~~Raumberg Gumpenstein, A 8952 Irdning des BMFLuFWUuW, 15. Gumpensteiner~~

893 ~~Lysimetertagung 2013, Lysimeterforschung als Bestandteil der Entscheidungsfindung Lysimeter~~
894 ~~research as an instrument for decision support, 203-206, 2013~~

895 Peters, A., Nehls, T., Schonsky, H., and Wessolek, G.: Separating precipitation and
896 evapotranspiration from noise - a new filter routine for high-resolution lysimeter data, Hydrology
897 and Earth System Sciences, 18, 1189-1198, doi:10.5194/hess-18-1189-2014, 2014.

898 Qu, W., Bogaen, H. R., Huisman, J. A., and Vereecken, H.: Calibration of a Novel Low-Cost Soil
899 Water Content Sensor Based on a Ring Oscillator, Vadose Zone Journal, 12, 23,
900 doi:10.2136/vzj2012.0139, 2013.

901 Richter, D.: Ergebnisse methodischer Untersuchungen zur Korrektur des systematischen
902 Messfehlers des Hellmann-Niederschlagsmessers Berichte des Deutschen Wetterdienstes, 194,
903 1995.

904 Savitzky, A. and Golay, M.: Smoothing and Differentiation of Data by Simplified Least Squares
905 Procedures, Analytical Chemistry, 36, 1627-1639, doi:10.1021/ac60214a047, 1964.

906 Schrader, F., Durner, W., Fank, J., Gebler, S., Pütz, T., Hannes, M., and Wollschläger, U.:
907 Estimating Precipitation and Actual Evapotranspiration from Precision Lysimeter Measurements,
908 Procedia Environmental Sciences, 19, 543-552, doi:10.1016/j.proenv.2013.06.061, 2013.

909 Schotanus, P., Nieuwstadt, F. T. M., de Bruin, H. A. R.: Temperature measurement with a sonic
910 anemometer and its application to heat and moisture fluxes. Boundary-Layer Meteorology 26,
911 81-93, doi:10.1007/bf00164332, 1983.

912 Scott, R. L.: Using watershed water balance to evaluate the accuracy of eddy covariance
913 evaporation measurements for three semiarid ecosystems, In: Agricultural and Forest
914 Meteorology, 150(2), 219-225, doi:10.1016/j.agrformet.2009.11.002, 2010.

915 Sevruk, B.: Methodische Untersuchungen des systematischen Messfehlers der Hellmann-
916 Regenmesser im Sommerhalbjahr in der Schweiz. Mitteilungen Nr. 52, Versuchsanstalt für
917 Wasserbau, Hydrologie und Glaziologie, ETH Zürich, 299 Book pp., 1981.

918 Sevruk, B.: Wind induced measurement error for high-intensity rains. Proc. International
919 Workshop on Precipitation Measurement, WMO Tech. Document 328, St. Moritz, Switzerland,
920 199-204, 1989.

921 Sevruk, B.: Adjustment of tipping-bucket precipitation gauge measurements, *Atmospheric*
922 *Research*, 42, 237-246, doi:10.1016/0169-8095(95)00066-6, 1996.

923 Strangeways, I.: A history of rain gauges, *Weather*, 65, 133-138, 10.1002/wea.548, 2010.

924 Twine, T. E., Kustas, W. P., Norman, J. M., Cook, D. R., Houser, P. R., Meyers, T. P., Prueger, J.
925 H., Starks, P. J., and Wesely, M. L.: Correcting eddy-covariance flux underestimates over a
926 grassland, *Agricultural and Forest Meteorology*, 103, 279-300, doi:10.1016/S0168-
927 1923(00)00123-4, 2000.

928 Unold, G. and Fank, J.: Modular Design of Field Lysimeters for Specific Application Needs,
929 *Water Air Soil Pollution, Focus*, 8, 233-242, doi:10.1007/s11267-007-9172-4, 2008.

930 Vaughan, P. J., Trout, T. J., and Ayars, J. E.: A processing method for weighing lysimeter data
931 and comparison to micrometeorological ET_0 predictions, *Agricultural Water Management*, 88,
932 141-146, doi:10.1016/j.agwat.2006.10.008, 2007.

933 Vaughan, P. J. and Ayars, J.: Noise Reduction Methods for Weighing Lysimeters, *Journal of*
934 *Irrigation and Drainage Engineering*, 135, 235-240, doi:10.1061/(ASCE)0733-
935 9437(2009)135:2(235), 2009.

936 Webb, E. K., Pearman ,G. I., Leuning, R.: Correction of flux measurements for density effects
937 due to heat and water vapour transfer, *Quarterly Journal of the Royal Meteorological Society*,
938 106, 85-100, doi:10.1002/qj.49710644707, 1980.

939 Wegehenkel, M. and Gerke, H. H.: Comparison of real evapotranspiration measured by weighing
940 lysimeters with simulations based on the Penman formula and a crop growth model, *Journal of*
941 *Hydrology and Hydromechanics*, 61, 161–172, doi:10.2478/johh-2013-0021, 2013.


942 Wilczak, J., Oncley, S., and Stage, S.: Sonic Anemometer Tilt Correction Algorithms, *Bound-*
943 *Layer Meteorology*, 99, 127-150, doi:10.1023/a:1018966204465, 2001.

944 Wilson, K. B., Hanson, P. J., Mulholland, P. J., Baldocchi, D. D., and Wullschleger, S. D.: A
945 comparison of methods for determining forest evapotranspiration and its components: sap-flow,
946 soil water budget, eddy covariance and catchment water balance, *Agricultural and Forest*
947 *Meteorology*, 106, 153-168, doi:10.1016/S0168-1923(00)00199-4, 2001.

948 Yang, D., Goodison, B. E., Metcalfe, J. R., Golubev, V. S., Bates, R., Pangburn, T., and Hanson,
949 C. L.: Accuracy of NWS 8" Standard Nonrecording Precipitation Gauge: Results and Application
950 of WMO Intercomparison, *Journal of Atmospheric and Oceanic Technology*, 15, 54-68,
951 doi:10.1175/1520-0426(1998)015<0054:AONSNP>2.0.CO;2, 1998.

952 Zacharias, S., Bogena, H., Samaniego, L., Mauder, M., Fuß, R., Pütz, T., Frenzel, M., Schwank,
953 M., Baessler, C., Butterbach-Bahl, K., Bens, O., Borg, E., Brauer, A., Dietrich, P., Hajnsek, I.,
954 Helle, G., Kiese, R., Kunstmann, H., Klotz, S., Munch, J. C., Papen, H., Priesack, E., Schmid, H.
955 P., Steinbrecher, R., Rosenbaum, U., Teutsch, G., and Vereecken, H.: A Network of Terrestrial
956 Environmental Observatories in Germany, *Vadose Zone Journal*, 10, 955-973,
957 doi:10.2136/vzj2010.0139, 2011.

958 Zenker, T.: Verdunstungswiderstände und Gras-Referenzverdunstung : Lysimeteruntersuchungen
959 zum Penman-Monteith-Ansatz im Berliner Raum. PhD thesis, Berlin University of Technology,
960 Germany, 2003.




962

963 **Fig. 1.** Overview of the Rollesbroich study site (left) showing the locations of the lysimeter, the
 964 rain gauge, the eddy covariance station, the catchment boundaries and the SoilNet devices. All
 965 devices are arranged within a radius of 50 meters including the nearest SoilNet device (SN 30)
 966 for ~~comparisons~~ comparison of temperature and soil water content with the surrounding field. The
 967 map on the right shows the location of the Rollesbroich catchment in Germany.




968

969 **Fig. 2.** The lysimeter set-up of the Rollesbroich study site (November 2012).




970

971 **Fig. 3.** Schematic drawing of the lysimeter soil monolith (left) and service well (right) used in the
 972 TERENO-SoilCan project. The illustration of the lysimeter (left) shows the weighted soil column
 973 container with slots for soil moisture (TDR), temperature (SIS, TS1), matric potential sensors
 974 (SIS), soil water sampler (SIC20) and silicon porous suction cup rake (SIC40) installation inside
 975 and outside the monolith. The service well contains the weighted drainage tank and sampling
 976 tubes for each affiliated lysimeter (courtesy of UMS GmbH Munich, 2014, used by permission).




977

978 **Fig. 4.** Daily precipitation sums of tipping bucket (blue) and difference in precipitation
 979 measurements between lysimeter and TB (red) at the Rollesbroich study site for 2012.




980
 981
 982
 983

Fig. 5. Cumulated average of lysimeter drainage and soil moisture storage on a daily basis. The colored areas indicate the range of minimum and maximum cumulated drainage and soil water storage for the individual lysimeters.




984

985 **Fig. 6.** Precipitation, temperature and dew point temperature from May 5 – May 6 2012 at the
 986 Rollesbroich site. The fog symbol indicates the hours with fog occurrence (detected with installed
 987 surveillance system) for the investigated period.




988

989 **Fig. 67.** Relationship between wind speed and precipitation residuals relative to TB precipitation
 990 on a daily basis. The relationships are classified according precipitation intensities of 1-5 mm
 991 (a), 5-10 mm (b), and > 10 mm (c). Potential rime and dew situation are excluded from the
 992 calculation.




993

994 **Fig. 8.** Cumulative ET_a -LYS, ET_a -EC (corrected according to Bowen ratio) and ET_e -FAO), ET_{PM}
 995 on hourly basis for 2012. Displayed are also ET_a -EC max. and ET_a -EC min. The area in grey
 996 shows the range of minimum and maximum cumulated ET_a for the individual lysimeters. For
 997 explanation see text.




998

999 **Fig. 79.** Grass heights at the lysimeter field, the lysimeter devices, and the field behind the EC
1000 station for 2012. The EC device is centrally located in between these two fields
1001 grass length at the lysimeter devices was reconstructed by comparing grass length measurements of the lysimeter
1002 field with the observations of the surveillance system. The star (*) indicates the presence of a
1003 snow cover. Grass cutting dates on lysimeter devices are marked by dashed lines. ~~For further~~
1004 ~~explanations see text.~~



1005

1006 **Fig. 810.** Differences between daily ET for 2012. Displayed are $ET_a-EC - ET_e-FAOET_{PM}$ (a),
 1007 $ET_a-LYS - ET_e-FAOET_{PM}$ (b) and $ET_a-LYS - ET_a-EC$ (c). The dashed lines indicate harvest at
 1008 lysimeters. **For explanation see text.**




1009

1010

1011


1012

Fig. 911. Relationship between grass length difference (between the lysimeters and the field behind the EC-device) and ET_a difference measured by lysimeters and EC station from May 21 – July 3.



1013

1014 **Fig. 12.** Mean hourly rates daily cycle of ET_a-LYS, ET_a-EC and ET_e-calculated according
 1015 FAOET_{PM} for 2012.



1016

1017 | **Fig. 1013.** Differences in daily mean soil temperature (averaged over the six lysimeters), a nearby
 1018 | SoilNet device (SN 30) and the mean of all available SoilNet devices located at the study site.

1019 **Tables**

1020 **Tab. 1.** Site specific wind exposition coefficient b [-] and empiric precipitation type coefficient
1021 ϵ [-] for different precipitation types at an open space gauge location.

Precipitation Type	b	ϵ
liquid (summer)	0.345	0.38
liquid (winter)	0.34	0.46
mixed	0.535	0.55
snow	0.72	0.82

1022
1023
1024
1025

Tab. 2. Monthly precipitation sums for lysimeter, tipping bucket, corrected tipping bucket data and a comparison between the hourly precipitation values of lysimeter and uncorrected TB in terms of coefficient of determination (R²), root mean square error and other statistics at the Rollesbroich study site for 2012. Missing data % refers to the percentage of hourly precipitation data not available for comparison.

<u>Month</u>	<u>Lysimeter Average [mm]</u>	<u>Min. / Max. Lysimeter [mm]</u>	<u>Tipping Bucket [mm]</u>	<u>Tipping Bucket corrected [mm]</u>	<u>R²</u>	<u>RMSE</u>	<u>LYS/TB %</u>	<u>LYS/ TBcorr %</u>	<u>Missing Data %</u>
<u>Jan</u>	<u>70.9</u>	<u>57.6 / 79.3</u>	<u>94.0</u>	<u>110.7</u>	<u>0.48</u>	<u>0.30</u>	<u>75.6</u>	<u>64.0</u>	<u>11.2</u>
<u>Feb</u>	<u>36.2</u>	<u>31.4 / 48.9</u>	<u>21.1</u>	<u>26.0</u>	<u>0.13</u>	<u>0.32</u>	<u>171.6</u>	<u>139.2</u>	<u>46.1</u>
<u>Mar</u>	<u>17.3</u>	<u>16.2 / 18.8</u>	<u>5.1</u>	<u>7.3</u>	<u>0.18</u>	<u>0.16</u>	<u>339.2</u>	<u>237.0</u>	<u>16.4</u>
<u>Apr</u>	<u>72.5</u>	<u>71.1 / 74.6</u>	<u>65.3</u>	<u>78.2</u>	<u>0.90</u>	<u>0.09</u>	<u>111.0</u>	<u>92.7</u>	<u>0.0</u>
<u>May</u>	<u>90.7</u>	<u>89.4 / 94.1</u>	<u>79.3</u>	<u>88.8</u>	<u>0.99</u>	<u>0.09</u>	<u>114.4</u>	<u>114.4</u>	<u>0.0</u>
<u>Jun</u>	<u>139.9</u>	<u>137.5 / 143.1</u>	<u>134.7</u>	<u>147.2</u>	<u>0.96</u>	<u>0.21</u>	<u>103.9</u>	<u>95.0</u>	<u>0.0</u>
<u>Jul</u>	<u>148.5</u>	<u>146.3 / 152.2</u>	<u>147.0</u>	<u>159.2</u>	<u>0.95</u>	<u>0.28</u>	<u>101.0</u>	<u>93.3</u>	<u>0.0</u>
<u>Aug</u>	<u>105.7</u>	<u>100.4 / 109.4</u>	<u>84.5</u>	<u>91.9</u>	<u>0.94</u>	<u>0.15</u>	<u>125.1</u>	<u>115.0</u>	<u>0.0</u>
<u>Sep</u>	<u>36.5</u>	<u>23.5 / 39.2</u>	<u>25.6</u>	<u>30.5</u>	<u>0.58</u>	<u>0.13</u>	<u>142.6</u>	<u>119.7</u>	<u>0.0</u>
<u>Oct</u>	<u>67.5</u>	<u>65.7 / 69.5</u>	<u>66.2</u>	<u>75.2</u>	<u>0.74</u>	<u>0.23</u>	<u>102.0</u>	<u>89.8</u>	<u>13.4</u>
<u>Nov</u>	<u>55.3</u>	<u>52.7 / 56.9</u>	<u>38.3</u>	<u>45.8</u>	<u>0.84</u>	<u>0.08</u>	<u>144.4</u>	<u>120.7</u>	<u>0.0</u>
<u>Dec</u>	<u>186.0</u>	<u>178.5 / 194.4</u>	<u>121.0</u>	<u>136.1</u>	<u>0.30</u>	<u>0.35</u>	<u>153.7</u>	<u>136.7</u>	<u>0.0</u>
<u>SUM /MEAN</u>	<u>1027.1</u>	<u>996.2 / 1037.7</u>	<u>882.1</u>	<u>996.9</u>	<u>0.88</u>	<u>0.47</u>	<u>116.4</u>	<u>103.0</u>	<u>7.1</u>

1026

1027 **Tab. 3.** Monthly ET_a (by lysimeter and EC), ~~ET_e-FAO~~ET_{PM} sums and R² between different ET data products on an hourly basis for 2012.
 1028 Missing data ~~provides% refers to~~ the percentage of hourly ~~daytime~~ ET data (ET_a-EC, ET_a-LYS) between sunrise und sunset not available
 1029 for comparison. Hence, the total yearly ET amount is ca. 18 % reduced compared to gap free ET estimations. ~~Missing data provides the~~
 1030 ~~percentage of hourly evapotranspiration data (sunrise—sunset) not available for comparison.~~

	2012												Mea n		
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	<u>Sum</u>		
ET _a - EC [mm]	5.2	1.3	27.8	38.4	84.3	62.7	80.3	94.2	56.0	25.2	9.3	3.6	488.3		
<u>ET_{PM}</u> [mm]	3.9	1.5	<u>30.5</u>	<u>37.5</u>	<u>84.2</u>	<u>69.</u> <u>7</u>	<u>84.0</u>	<u>113.5</u>	<u>58.9</u>	<u>24.6</u>	<u>9.40</u>	<u>2.75</u>	<u>519.8</u>		
ET _a - LYS [mm]	2.5	2.2	26.4	35.6	80.2	65.7	82.7	121.7	52.7	23.9	7.6	5.9	507.4		
Min. / Max. ET _a - LYS [mm]	2.1 / 2.7	1.3 / 3.1	25.9 / 26.8	34.4 / 37.6	75.2 / 85.2	62.1 / 68.2	67.8 / 91.0	116.8 / 125.2	49.6 / 58.8	21.9 / 27.1	6.8 / 8.9	3.0 / 8.7	467.1 523.1		
R ² ET _a - EC - ET _a - LYS	0.02	0.02	0.82	0.76	0.79	0.84	0.86	0.86	0.66	0.66	0.39	0.06		0.81	
R ² ET _a - LYS = <u>ET_{PM}</u>	0.13	<u>0.030</u> <u>0</u>	0.8 7	<u>0.8</u> <u>±</u>	0.82	<u>0.86</u>	<u>0.91</u>	0.89	<u>0.879</u> <u>2</u>	<u>0.957</u> <u>8</u>	<u>0.6</u> <u>8</u>	0.70	<u>0.41</u>	0.08	<u>0.89</u>

R²														
ET_a- EC = ET_{PM}	0.12	0.00	0.94	<u>0.93</u>	<u>0.9</u> <u>5</u>	0.90	<u>0.89</u>	<u>0.88</u>	<u>0.88</u>	<u>0.82</u>	<u>0.73</u>	<u>0.44</u>		<u>0.91</u>
Missing Data %	33.2	36.9	8.1	23.5	21.5	26.5	21.9	12.9	14.0	25.8	25.0	45.3		24.5

1 | **Tab. 34.** Measured mean monthly latent heat fluxes and corrections for EBD for 2012.

Month	Mean LE [W m⁻¹]	Mean LE corr. [W m⁻¹]	Differences LE corr. - LE	Difference mean LE corr. - LE %
Jan	21.9	29.8	7.9	36.2
Feb	8.7	11.9	3.2	36.9
Mar	78.1	94.0	15.9	20.4
Apr	86.4	101.8	15.3	17.7
May	138.7	164.6	25.9	18.7
Jun	111.8	125.8	14.0	12.6
Jul	136.3	157.2	20.9	15.3
Aug	151.6	181.4	29.8	19.6
Sep	104.0	129.2	25.2	24.2
Oct	61.3	79.6	18.3	29.9
Nov	24.4	32.1	7.7	31.4
Dec	22.0	28.3	6.3	28.5
SUM/MEAN	78.8	94.6	15.9	24.3

2