

Interactive comment on “Modelling of snow processes in catchment hydrology by means of downscaled WRF meteorological data fields” by K. Förster et al.

Anonymous Referee #2

Received and published: 13 May 2014

General comments:

The paper deals with the value of WRF downscaled meteorological fields for driving snow modules. Ground measurements usually limit performances of hydrologic model in mountain catchments, where atmospheric forcings may vary within restricted horizontal distances due to topographic effects. Several published contributions describe the value of downscaled meteo data for driving energy-balance/temperature-index snowmelt models. Here, the innovative contribution is related to the use of four independent snowmelt models, as it is indicated within the introduction. This should help

C1363

in making results model-independent, if one is investigating the value of downscaled inputs for modeling snow processes.

Overall, the topic is interesting and meets the requirements of the journal. Methods are clearly explained with synthetic sentences.

The assessment of the manuscript is highly conditioned by the fact that downscaled precipitation is not involved. This invalidates title, abstract and introduction. It even affects results, since it is not possible to draw conclusions about the usefulness of WRF atmospheric forcings for modeling snow accumulation and snowmelt. Your approach simulates well both discharge and melt runoff, but it is still dependent on in-situ data. Results do not provide useful information if one wants to applied WRF outputs for hydrologic purposes, thus avoiding observations. These limitations must be overcome if you are going to maintain the same targets for your paper. Results obtained using downscaled precipitation must be shown in accordance to the title.

An idea could be an additional comparative study between the performances you got at the point scale, driving snowmelt models by WRF fields (including precipitation, even if the simulation is not good) and the results obtained using in situ measurements (you stated that temperature and precipitation are recorded: these seem the key factors and they are certainly enough for the temperature-index approach and maybe for the model Walter et al. 2005.). This will give an indication about the loss of accuracy due to downscaled forcings when compared to reliable in situ data.

Again, catchment scale simulations should use downscaled precipitation. Here, a comparison could be performed for understanding whether this approach outperforms the results provided by a spatial distribution of observed temperature and precipita-

C1364

tion. You spent several sentences on the issue of "area representativeness of point observations" within the introduction. At the end, it is not clear if WRF data help for hydrologic modeling when compared against standard inputs (ie ground-based measurements). Methods for distributing point values (eg kriging) exist. For instance, using the simple degree-day approach, it could be implemented a spatial distribution of temperature with altitude (a constant lapse rate is usually adopted). Then, you can run the degree-day model coupled to PANTA RHEI.

It is not clear whether you are discussing the value of the combined use of WRF and ground-data or the value of WRF meteo data alone. In the first case, you should state that your study uses both data sources from the beginning, and the results must be discussed with this focus. Even in this case, additional analysis should be provided. For example, it could be possible to use precipitation and temperature recordings (which seem available at several stations around the catchment) and downscaled radiation, wind speed etc., if such data are not measured.

Specific comments

If you are not going to use WRF precipitation the title should be changed. This must be claimed also within the abstract. In the title, the sentence "modelling of snow processes in catchment hydrology" seems very general. Actually you are considering only one (and particular) case study. Probably the title should indicate this.

P 4066 Line 5: I do not think this is always true. Please state that the problem of spatial resolution is mainly related to complex topographies of mountain regions.

"Study area" section: It is not clear the whole number of stations available around the catchment and what kind of data are provided (only temperature and precipitation?). This is useful since you might run the models using ground observations and see what

C1365

happens in comparison with WRF inputs. This might be an interesting contribution by your paper.

"Selected winter season" section: I agree with referee 1 that two particular winter seasons are not enough for drawing conclusions. Please consider the possibility of involving a third hydrologic cycle.

"Snowmelt models" section: please, spend some times in describing what parameters you calibrated, perhaps by inserting a summary table. Are they calibrated using melt rates from the lysimeter?

Could be possible to better explain how you designed accumulation and melting for the temperature-index model? Did you consider refreezing? Are you using only one degree-day factor for the entire basin? Did you consider the additional energy input by rain-on-snow?

P 4074 line 9: how did you calibrate snowmelt models for catchment scale simulations? What did you calibrate?

P 4077 line 20: please remove comma after "concluded".

P 4081 line 19: the fact that considering snow processes instead of "no snow" you are improving runoff simulations is not a finding. It would have been a problem if it happened the opposite. On the contrary, you should discuss how much you improve the "no snow" simulation and if it justifies the use of a more complex hydrologic model. Anyway, your sentence does not seem scientifically relevant for the goals of your manuscript.

P 4082 line 3: to do what? you are combining data sources. Please explain why and where the presented approach could be applied.

Fig. 5: the plot is not very clear due to the high frequency. Please consider to enlarge the x-axis or restrict the temporal window you are showing. Otherwise you may split it into two time frames.

Fig. 8: why the name "snowmelt simulation" as in fig. 7? Are they streamflows at

C1366

the closure section? Here, the name of the figure should be "catchment discharges considering snowmelt", or something like that.

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., 11, 4063, 2014.

C1367