

Co-evolution of volcanic catchments in Japan

Takeo Yoshida^{1,2} and Peter A Troch¹

¹University of Arizona, Tucson, AZ, USA

²National Agricultural and Food Research Organization, Tsukuba, Ibaraki, Japan

Correspondence to: Takeo Yoshida (takeoys@affrc.go.jp)

Abstract. Present day landscapes have evolved over time through interactions between the prevailing climates and geological settings. Understanding the linkage between spatial patterns of landforms, soils, and vegetation in landscapes and their hydrological response is critical to make quantitative predictions in ungaged basins. Catchment co-evolution is a theoretical framework that seeks to formulate hypotheses about the mechanisms and conditions that determine the historical development of catchments and how such evolution affects their hydrological response. In this study, we selected 14 volcanic catchments of different ages (from 0.225 to 82.2 Ma) in Japan. We derived indices of landscape properties (drainage density and slope area relationship) as well as hydrological response (annual water balance, baseflow index, and flow duration curves) and examined their relation with catchment age and climate (through the aridity index). We found a significant correlation between drainage density and baseflow index with age, but not with climate. The intra-annual flow variability was also significantly related to catchments age. Younger catchments tended to have lower peak flows and higher low flows, while older catchments exhibited more flashy runoff. The decrease in baseflow with catchment age is consistent with the existing hypothesis that in volcanic landscapes the major flow pathways change over time from deep groundwater flow to shallow subsurface flow. The drainage density of our catchments decreased with age, contrary to previous findings in a set of similar, but younger volcanic catchments in the Oregon Cascades, in which drainage density increased with age. In that case, older catchments were thought to show more landscape incision due to increasing near-surface lateral flow paths. Our results suggests two competing hypotheses on the evolution of drainage density in mature catchments. One is that as catchments continue to age, the hydrologically active channels retreat because less recharge leads to lower average aquifer levels and less baseflow. The other hypothesis is that the active channels do not undergo much surface dissection after the catchments reach maturity.

1 Introduction

The hydrological functions that partition incoming water, energy and carbon at the land surface vary from place to place, reflecting spatial variability in land surface characteristics (landforms, soils, vegetation). Because of this spatial variability, hydrological predictions in ungaged basins are highly

uncertain (Wagener et al., 2007; Sivapalan, 2003). Improving these predictions will require a better understanding of the different interactions between landscape features and hydrological response, especially in a changing environment (McDonnell et al., 2007; Wagener et al., 2013).

Present day land surface characteristics in pristine landscapes have co-evolved over time through various interactions between the prevailing climates and geological settings (bedrock type, tectonic uplift) (Blöschl et al., 2013; Troch et al., 2015). Hydrologists have coined the term catchment co-evolution as a theoretical framework that seeks to formulate explanatory hypotheses about spatial patterns of landscape characteristics and the corresponding hydrological response, based on the historical development of catchments (Sivapalan et al., 2012; Troch et al., 2013; Harman and Troch, 2013). The results of historical development are captured in the soil and hillslope catena profiles, the geometry of channel networks and the spatial distribution of plant functional types. This landscape organization determines the hydrological functions that partition incoming water, energy and carbon. A detailed mechanistic reconstruction of how catchments have evolved to their present state is a daunting task, given the many processes involved and the unknown initial conditions during landscape formation.

Therefore, formulating and testing explanatory hypotheses of catchment co-evolution and hydrological response requires a different approach from the classic mechanistic Newtonian approach. Recently, an alternative approach, called Darwinian hydrology, was proposed that seeks to document patterns of variation in populations of hydrological systems, and explain these patterns in terms of the processes and conditions of their historical development (Wagener et al., 2013; Harman and Troch, 2013). Harman and Troch (2013) summarized what the Darwinian approach might look like in watershed science as follows: (1) collect extensive and detailed observations of the pattern to be explained, (2) conceive a hypothesis that accounts for as many of the observations as possible, (3) derive from this hypothesis a subsequent set of circumstances that also must hold if the hypothesis is true, and (4) critically test whether these consequences do hold. In this study, we build upon this framework and explore empirically the links between landscape properties, hydrological response, and time since formation (age) of catchments of similar bedrock geology (volcanic rock deposits of different geologic age).

It may be argued that climate exerts a first-order control on hydrologic partitioning and that physical, chemical and biological catchment properties only have a second-order impact (e.g., Berghuijs et al., 2014; Sivapalan et al., 2011). However, these second-order controls explain much of the functional patterns that determine flow regimes. Mushiake et al. (1981) observed that bedrock lithology plays a key role in the natural variations of functional patterns in Japanese catchments. They showed that catchments on older sedimentary rocks formed in the Paleozoic and Mesozoic exhibit flashy hydrographs and a low baseflow component compared to younger catchments on volcanic rock. They also found that catchments underlain by younger volcanic rock formed in the Quaternary have a greater groundwater component compared to those on older volcanic rock formed in the Tertiary.

65 Lohse and Dietrich (2005) found that vertical saturated hydraulic conductivity declines as volcanic
soils in Hawaii develop with time, due to the formation of secondary clay mineral layers in those
soil profiles. This evolution impedes rates of vertical percolation of water and increases shallow lat-
eral subsurface flow. Jefferson et al. (2010) studied landscape and hydrological signatures in basalt
catchments in the Oregon Cascades and reported that springs were absent in older catchments, which
70 exhibit well-developed drainage networks and flashy hydrographs. They hypothesized that the dif-
ferences in hydrological response between young and old catchments was the result of changes in
flow pathways, from vertical recharge and groundwater dominated flow to shallow subsurface lateral
flow with sufficient erosive power to create channel networks through landscape incision.

Prior work has suggested that climate controls variations in hydrologic regimes in two ways.
75 One is direct control on the water balance by function of available water and energy (e.g., Budyko,
1974; Milly, 1994; Troch et al., 2009; Sivapalan et al., 2011). The other is indirect control on the
hydrologic regime as one of four independent factors (climate, bedrock weatherability, tectonic up-
lift and time) that control the evolution of landscape properties (landforms, soils, and vegetation,
Troch et al., 2015). For example, Gentine et al. (2012) found that catchment-scale rooting depth was
80 strongly correlated with the seasonal distribution of water and energy availability in 431 catchments
in the United States. Wang and Wu (2013) found a strong inverse correlation between perennial
stream density and climate aridity index in 185 catchments across the United States. Both studies
suggested that these catchments evolved by adapting to the long-term climate conditions.

From this overview it is clear that both climate and time influence spatial variations of hydrologi-
85 cal response. In this study we investigated the degree of control of climate and time on the differences
in hydrological response between catchments that have different internal organizations. More specif-
ically, we asked the following questions: (1) How do landscape features and hydrological response
differ with catchment age and climate? (2) What is the dominant control of catchment co-evolution
in basaltic landscapes? (3) What are the possible mechanisms related to the historical development
90 of volcanic catchments? To address these questions we selected 14 catchments dominated by basaltic
rock with different ages and from a range of climatic regions of Japan. The age of the catchments
ranges from 0.225 to 82.2 Ma (million years). The aridity index (ratio of annual potential evapora-
tion to annual precipitation) for these catchments ranges from 0.28 to 0.82. For each catchment we
derived indices of landscape features (drainage density and slope area relationship) and hydrological
95 response (annual water balance, baseflow index, and flow duration curves). We then used regression
analysis to explore possible relationships between these features and catchment age and climate,
respectively. Based on the observed relationships, we proposed a hypothesis that explains as many
observations in volcanic catchments, both these and others, as possible.

2 Study area and materials

100 2.1 Study catchments

The study area is located within the Japanese archipelago, where 28% of the land is underlain by volcanic rock. The makeup of the volcanic landscape in Japan with respect to time since formation, or geological age, is as follows: 11% was formed in the Quaternary, 12% in the Neogene, 1% in the Paleogene, and 4% in the Cretaceous (Murata and Kano, 1995). The volcanic landscapes formed in
105 the Quaternary and Neogene are mostly in north-eastern and south-western Japan, and those formed in the Paleogene and Cretaceous are predominantly in central and western Japan. The majority of the basement rock of Japan was formed in the Jurassic to Paleogene at the margin of the Asian continent and moved toward the current position during the spreading of the Japan Sea Basin in the Neogene (Taira, 2001). The volcanic rocks formed in the Cretaceous, which are prevalent in western Japan,
110 were formed at the margin of the Asian continent and have been exposed to the atmosphere since their formation (Taira, 2001). The volcanic rocks of the Quaternary and Neogene age, on the other hand, were formed after the spreading of the Japan Sea Basin and thus have lain in their current position since their formation.

The catchments selected for the study are among the catchments listed in the Japanese Dam
115 Database (National Institute for Land and Infrastructure Management, Japan, 2015). Out of 130 catchments listed in the database, we selected 14 that contain more than 50% volcanic rock coverage (Table 1). Basalt is the dominant rock type in all catchments, while rhyolite is found in five of the catchments (GOS, YUD, AIM, IKR, and KWM), but at less than 15% of the volcanic rock area. The catchment areas range from 30.9 to 635.0 km². Daily streamflow observed at each dam was
120 obtained from the Japanese Dam Database. The streamflow observations are inflow to the dams and no major dams are located upstream, therefore the streamflows examined are not affected by human regulations.

The catchment ages were assigned from the average of the ages of different volcanic rocks, weighted by area of coverage, obtained from the Seamless Geological Map of Japan (Geological
125 Survey of Japan, 2012). The map is the product of the merged 1:200,000 geologic map sheets covering the entire country of Japan. The age of the catchments ranged from 0.225 Ma to 82.2 Ma. The youngest catchment (SNK) is located in central Japan (Fig.1). The oldest catchment (HAZ) is covered with basalt formed in the Late Cretaceous (Fig.2). The other 12 catchments were formed in the Neogene Period (2.7-18.5 Ma) and are clustered in north-eastern Japan, with the exception of the
130 Shimouke catchment in southern Japan.

To delineate catchments, digital elevation models (DEMs) of 10 × 10 m were downloaded from the website of the Geographic Survey Institute, Japan (2015). The stream networks were derived from digitized GIS maps from the National Land Numerical Information website (Ministry of Land, Infrastructure and Transportation, Japan, 2015). Tectonic uplift for each catchment was estimated

Table 1. List of study catchments

Catchment Name	ID	Age (Ma)	Volcanic Coverage (-)	Area (km ²)
Shinaki	SNK	0.225	0.994	30.9
Tamagawa	TMG	2.78	0.852	287.0
Aseishigawa	ASE	3.52	0.940	225.5
Kawamata	KWM	3.69	0.507	179.4
Shimouke	SIM	4.00	0.982	185.0
Gosho	GOS	6.39	0.750	635.0
Naruko	NAR	8.43	0.598	210.0
Ikari	IKR	10.7	0.653	271.2
Aimata	AIM	11.7	0.603	110.8
Ishibuchi	ISB	11.8	0.732	154.0
Kamafusa	KAM	12.8	0.621	195.3
Shichikashuku	SCK	13.9	0.619	230.9
Yuda	YUD	18.5	0.633	583.0
Hazi	HAZ	82.2	0.623	217.0




Figure 1. Study catchments in north-eastern Japan




Figure 2. Study catchments in western Japan

135 from the map of uplift in the Quaternary with 250 m intervals (National Research Center for Disaster Prevention, Japan, 1973). We imposed the delineated catchments on the tectonic uplift map and determined the uplift from the dominant area of the catchment coverage on the map.

2.2 Meteorological data

A 20 year (from 1993 through 2012) meteorological dataset (including precipitation, temperature, 140 humidity, and wind speed) was distilled from a gridded dataset (hereafter referred to as the raw dataset). The gridded dataset covers all of Japan with grid cells of approximately 1km × 1km in size. The data interpolation was performed using observed data from the Automated Meteorological Data Acquisition System (AMeDaS) of Japan and the Mesh Climate Data 2000 (Japan Meteorological Agency, 2002). The Mesh Climate Data 2000 provides estimated of monthly climatic means from 145 1971 through 2000 for these grid cells by empirical relationships to account for orographic effects. The details of the raw dataset are described in the Appendix.

3 Methods

3.1 Correction of meteorological data

The precipitation in the raw dataset accounts for orographic effects based on empirical relation- 150 ships between precipitation and geographic features, such as elevation and distance from shore-line. It might not represent local orographic effects (e.g., higher rate of increase in precipitation on windward side of mountains), and does not correct for snow undercatch. Thus, we corrected the

precipitation data as described in the Appendix. We assessed the validity of the raw and corrected datasets by means of the long-term water balance of the catchments. Given the mean of catchment-normalized annual precipitation for the raw dataset P_{raw} (mm/y) and precipitation-corrected dataset P_{cor} (mm/y), and streamflow, Q (mm/y), we estimated the mean annual evapotranspiration for each dataset, E_{raw} and E_{cor} (mm/y), as follows:

$$E_{raw} = P_{raw} - Q \quad (1)$$

$$E_{cor} = P_{cor} - Q \quad (2)$$

To cross-check the acceptability of E_{raw} and E_{cor} , catchment-scale actual evapotranspiration, E_a (mm/y), was directly estimated with the complementary relation method (e.g., Bouchet, 1963; Morton, 1978; Brutsaert and Stricker, 1979). In this study, the complementary relationship method proposed by Otsuki et al. (1984a) was used to estimate E_a for each catchment. The method was used because it was rigorously validated with the long-term evapotranspiration in several experimental watersheds across Japan and does not require additional parameters (Otsuki et al., 1984b; Oue et al., 1992; Yoshida et al., 1994).

Thus we have used three types of meteorological datasets to estimate the long-term water balance.

- Raw dataset: P_{raw} , E_{raw}
- Corrected P dataset: P_{cor} , E_{cor}
- Estimated E dataset: P_{cor} , E_a

The estimated data were validated against the Budyko hypothesis (Budyko, 1974). Budyko (1974) postulated an empirical relationship between the long-term evaporation index, E/P and the aridity index, PE/P and argued that this relationship can be satisfactorily applied to most catchments. The large bias or discrepancy of estimated evaporation from the prediction of the Budyko curve was interpreted to derive from errors in the dataset (Fig.3).

3.2 Geomorphologic signatures

Two measures of landscape characteristics were used: drainage density and slope area relationship. Drainage density is defined as the total length of streams divided by catchment area (km/km²). To obtain the stream length, we used the length of the vectors defining blue-colored streamlines in the digitized version of topographic maps (Geographic Survey Institute, Japan, 2015). The scale of the map is 1:25,000. Because our study catchments are all located in humid regions where precipitation is evenly distributed, we assumed that the blue lines of our digital dataset represents perennial streams.

To understand the timescale of landscape evolution with climate and geological age, we compared the relationship between local slope and contributing area for all the catchments. The slope

area relationship is the log-log relationship between the local slope and the upslope contributing area for all grid cells within a catchment. The local slope was estimated for each grid cell of a 10 m DEM with the D-infinity algorithm (Tarboton, 1997). The slope area relationship can be used to discriminate between different geomorphologic regimes (Tarboton et al., 1992; Tucker and Bras, 1998; Bogaart and Troch, 2006). The slope area relationship typically shows one or more inflection points, each of which can be interpreted as the threshold points where the dominant geomorphological process switches to other processes (Tucker and Bras, 1998; Bogaart and Troch, 2006). Hillslopes dominated by diffusional transport yield a positive slope area relationship, i.e., slope increases with area, while hillslopes dominated by landslides or debris flows theoretically shows uniform slope gradient at a value equal to the threshold slope that induces the landslide (Tucker and Bras, 1998). Inflection points are found for larger contributing area than those at the transition from hillslopes dominated by diffusional transport to the ones dominated by landslide/debris flow. The larger inflection points can be interpreted as the initiation of the hillslopes dominated by fluvial transport, which typically shows a negative slope area relationship.

3.3 Hydrological signatures

Two signatures were used to represent the hydrological response of the study catchments. The first one is the baseflow index, the ratio of long-term baseflow component to total streamflow (e.g., Vogel and Kroll, 1992; Kroll et al., 2004). The higher the baseflow index the higher the groundwater contribution to catchment outflow. We separated the baseflow component from total streamflow as follows:

$$Q_b(t) = \epsilon Q_b(t-1) + \frac{1-\epsilon}{2} (Q(t) - Q(t-1)), \quad (3)$$

where $Q_b(t) = Q(t)$ when $Q_b(t) > Q(t)$; $Q(t)$ is total streamflow at time t , $Q_b(t)$ is the estimated baseflow, and ϵ is a low-pass filter parameter (Arnold and Allen, 1999; Eckhardt, 2005). To compare hydrological responses among catchments consistently, the filter parameter ϵ was set at 0.925 for all catchments (Sawicz et al., 2011).

To quantify an index of flow variability, the slope of the flow duration curve was used as the second signature. The flow duration curve is the cumulative probability distribution of daily streamflow. The slope of the flow duration curve is calculated between the 33rd and 66th streamflow percentiles, because this range represents a relatively linear part of the flow duration curve on semi-log scale (Yadav et al., 2007; Zhang et al., 2008). This is calculated from

$$S_{fdc} = \frac{Q_{33} - Q_{66}}{0.66 - 0.33}, \quad (4)$$

where Q_{66} and Q_{33} are the streamflows that correspond to 66% and 33% exceedance probability, respectively. A high slope value indicates a variable flow regime, while a low slope value means a more damped response. A damped response can arise as a result of a combination of persistent




Figure 3. Raw, corrected P and estimated E datasets plotted with Budyko curve

(wide-spread and year-round) precipitation, dominance of groundwater contribution to streamflow, or a combination of the two.

4 Results

4.1 Initial data evaluation

225 To evaluate data consistency with long-term water partitioning, we plotted evaporation index versus
 aridity index for all catchments and compared the plotted values against the Budyko curve (Fig.3).
 The plotted data points for the raw dataset are all well below the Budyko curve. Catchment-scale
 actual evapotranspiration, E_a , estimated from the complementary relation method, on the other hand,
 showed good agreement with the Budyko curve, suggesting realistic representation of long-term
 230 evapotranspiration in each of the catchment.

Errors in the estimates of the raw dataset could be caused by errors in discharge observations,
 trans-boundary groundwater flux, and catchment-averaged precipitation. The discharge, or dam in-
 flow, was validated with the dam outflow, and the bias in the total inflow to total outflow was less
 than 2% over 20 years. We thus assume that the dam inflow data is reliable. Another potential er-
 235 ror is trans-boundary groundwater flux. Quantifying this flux is very difficult; however, by focusing
 on relatively large catchments ($> 30\text{km}^2$), we argue that the influence of such a flux is negligible.
 It is more likely that the errors in annual evaporation estimated from the water balance are caused
 by underestimation of areal precipitation from the gridded data. Indeed, the corrected precipitation
 dataset showed substantial improvement from the raw dataset compared to the Budyko hypothe-

Table 2. Estimated climate variables and indices

ID	P_{cor} (mm/y)	PE (mm/y)	Q (mm/y)	E_{cor} (mm/y)	E_a (mm/y)	PE/P	E_{cor}/P
SNK	2454	932	1770	684	864	0.38	0.279
TMG	2937	916	2700	237	754	0.312	0.081
ASE	2215	903	1743	471	753	0.408	0.213
KWM	2333	947	1404	928	783	0.406	0.398
SIM	3112	1229	2267	844	899	0.395	0.271
GOS	2544	976	1908	636	780	0.384	0.25
NAR	2852	941	2202	649	752	0.33	0.228
IKR	1770	1006	1301	469	792	0.568	0.265
AIM	2652	1026	1679	973	839	0.387	0.367
ISB	3305	928	2541	763	743	0.281	0.231
KAM	2074	1024	1498	576	782	0.494	0.278
SCK	2100	1028	1326	774	807	0.49	0.369
YUD	3206	955	2316	890	727	0.298	0.278
HAZ	1513	1249	573	939	895	0.826	0.621

240 sis although some data still plot below the Budyko curve, likely caused by additional precipitation underestimation due to snow undercatch.

The aridity indices calculated with P_{cor} were used for subsequent analysis (Table 2). All the catchments lie in a energy limited region (i.e., $PE/P < 1$). Although no correlation between climate aridity indices and catchment ages was found ($R = 0.501, p = 0.068$), the oldest catchment lies in the driest region, while the youngest catchment lies in a humid region (Fig.4). Thus, it is not possible to thoroughly investigate the controls of climate and time on catchment co-evolution, and the regression of any signatures is substantially affected by the behavior of the youngest and oldest catchments.

4.2 Analysis by geological age: Controls of time on catchment co-evolution

All the landscape and hydrological indices obtained from the analysis are presented in Table 3. Baseflow index, drainage density and slope of the flow duration curve were all strongly correlated with catchment age (Figs.5 and 6). Fig.5 shows a significant correlation between the baseflow index and catchment age ($R = -0.756, p = 0.0017$), and between the drainage density and catchment age ($R = -0.603, p = 0.0222$), respectively. Because the catchment age spans three orders of magnitude, $\log(\text{catchment age})$ was used in the analysis. No significant correlation was found between catchment area and the baseflow index ($R = -0.24, p = 0.42$) or drainage density ($R = 0.23, p = 0.43$).

The oldest catchment (HAZ) had a baseflow index of 0.533, while in the youngest catchment (SNK), the baseflow index of 0.835 was the highest of the study catchments, suggesting groundwater-




Figure 4. Relation between aridity index and age for the study catchments

Table 3. Results of hydrometric and landscape analysis of study catchments

ID	Age	PE/P	E/P	Drainage Density (km/km^2)	Baseflow index (Q_b/Q)	Baseflow coefficient (Q_b/P)	Slope of FDC
SNK	0.225	0.380	0.352	1.080	0.835	0.602	0.322
TMG	2.78	0.312	0.257	0.882	0.713	0.656	0.850
ASE	3.52	0.408	0.340	0.887	0.775	0.610	0.585
KWM	3.69	0.406	0.336	0.898	0.737	0.444	0.827
SIM	4.00	0.395	0.289	0.859	0.585	0.426	0.880
GOS	6.39	0.384	0.307	1.000	0.684	0.513	0.870
NAR	8.43	0.330	0.264	0.736	0.707	0.546	0.668
IKR	10.7	0.568	0.448	0.867	0.708	0.520	0.776
AIM	11.7	0.387	0.316	0.771	0.712	0.451	1.071
ISB	11.8	0.281	0.225	0.969	0.666	0.512	1.034
KAM	12.8	0.494	0.377	0.707	0.680	0.491	0.735
SCK	13.9	0.490	0.384	0.793	0.710	0.448	0.754
YUD	18.5	0.298	0.227	0.948	0.663	0.479	1.011
HAZ	82.2	0.826	0.592	0.777	0.533	0.202	0.895




Figure 5. Influence of catchment age on baseflow index ($R = -0.756$, $p = 0.0017$) and drainage density ($R = -0.603$, $p = 0.0222$).




Figure 6. Slope of flow duration curves with catchment age ($R = 0.74$, $p = 0.002$)

dominated flow regime. The snow-dominated catchments tended to have a larger baseflow index, because we used a low-pass filter to exclude high frequency signals from daily hydrographs to separate baseflow from total runoff. Two of our catchments (SIM and HAZ) are not snow affected, whereas the others are snow-dominated as shown in Figs.23 and 24. Plot for the catchments that receive little snow are negatively deviated from the regression line (Fig.5).

The decrease in baseflow index with age was supported by the slope of the flow duration curves (Fig.6). The slopes of the flow duration curve were positively correlated with catchment age ($R = 0.74$, $p = 0.002$), i.e., as catchments age, the slope of the flow duration curve increases, indicating flashier runoff.




Figure 7. Flow duration curves for the catchments whose aridity index are in the range between 0.35 and 0.45. Inset of this figure illustrates flow duration curves for all study catchments. Each colored line represents data from a certain range in aridity index (key to line colors is as follows: blue: $PE/P < 0.35$; green: $0.35 < PE/P < 0.45$; orange: $0.45 < PE/P < 0.65$ and black: $PE/P > 0.65$).

The flow duration curves for all the study catchments are presented in the inset of Fig.7, in which each line represents the data from a certain range in aridity index. It is clear that the aridity index controls the total amount of generated streamflow, i.e., the higher the aridity index, less streamflow is generated by the catchment. Fig.7 depicts the flow duration curves for the catchments whose aridity index are in the range between 0.35 and 0.45. We used this range because it comprises the diversified catchments in terms of age (from 0.225 to 11.7 Ma). The slopes of the flow duration curves of relatively young catchments, less than 4 Ma, were noticeably less than those of the older catchments. This implies that flowpaths in older catchments with similar aridity index have changed from deep groundwater to surface or near-surface flowpaths.

It should be noted that there is no statistically significant log-linear fit between age and drainage density ($p = 0.28$) or age and slope of the flow duration curve ($p = 0.30$) without the youngest catchment. We will later discuss the uniqueness of the youngest catchments and how it influences the interpretation of our results.

4.3 Analysis by climate: Climate controls on catchment co-evolution

The drainage density and baseflow index shows no significant correlation with aridity index (Fig.8), although there are decreasing trends in both.




Figure 8. Climate controls on baseflow index and drainage density. The relationships with aridity index were not significant for both drainage density ($R = -0.451$, $p = 0.105$) and baseflow index ($R = -0.372$, $p = 0.190$).

To highlight the seasonality of the intra-annual distribution of precipitation in each catchment, we
 285 calculated the seasonal index SI (Walsh and Lawler, 1981).

$$SI = \frac{1}{P} \sum_{m=1}^{12} \left| X_m - \frac{P}{12} \right| \quad (5)$$

where P is the mean annual precipitation and X_m is the mean precipitation of month m . The maximum value of SI is 1.83 if all the precipitation fell in a single month, while the minimum is 0 where the precipitation is perfectly evenly distributed throughout the year. The calculated SI of our
 290 catchments ranged from 0.182 to 0.506, which means they are classified as 'Very equable', 'Equable but with a definite wetter season' or 'Rather seasonal with a short drier season' (Walsh and Lawler, 1981). We therefore assumed the seasonality of precipitation of our catchments does not influence the results.

4.4 Geomorphological signatures of landscape evolution

295 The rate of landscape evolution is a function not only of time, but also of climate, geology and tectonics that drive the evolution process (Troch et al., 2015). To understand the timescale and governing processes of landscape evolution, an estimation of tectonic uplift is essential, especially in dynamic landscapes such as those of Japan. **Fig.9** shows a strong correlation between the maximum slope captured in the mean slope area relationship for the study catchments and the rate of tectonic uplift
 300 in the Quaternary Period in Japan (National Research Center for Disaster Prevention, Japan, 1973). Numerical experiments with landscape evolution models also suggested that higher tectonic uplift rate yields steeper slopes (e.g., Yetemen, 2015). We thus assumed that the effect of the difference in tectonic uplift rate can be eliminated by comparing the catchments with similar maximum slope.




Figure 9. Tectonic uplift in the Quaternary period and maximum slope of mean slope area relationship

We then performed slope area analysis (Tarboton, 1997; Jefferson et al., 2010) to infer the evolu-
 305 tion pathways based on the current evidence. To quantify the activity level of climate on the land-
 scape evolution, we used effective precipitation $P_{eff} = P_{cor} - E_a$, because Jefferson et al. (2014)
 showed that mean annual precipitation has a strong link with erosion rate in basaltic islands and is
 the major driving factor for landscape evolution.

The mean slope area relationship for three catchments with the smallest range of uplift rates, less
 310 than 250 m in the Quaternary, are presented in Fig.10: SNK (0.225 Ma), ASE (3.52 Ma), and GOS
 (6.39 Ma). The mean slope area relationship for the oldest catchment HAZ (82.2 Ma) is also shown
 for comparison. Note that the maximum slopes of the mean slope area relationship were similar in
 SNK, ASE, and GOS as were the effective precipitation values; for HAZ differed in both respects.
 The smaller inflection points occurred at the contributing area value of approximately 300m^2 in all
 315 four catchments in Fig.10. The mean slope area relationship in the younger two catchments (SNK
 and ASE) was relatively straight from the inflection points down to the outlet of the catchments,
 whereas a larger inflection point, which can be interpreted as the point of initiation of a fluvial
 channel, was found in the GOS catchment at a contributing area value of $4.4 \times 10^4\text{m}^2$ (green dashed
 line in Fig.10), suggesting the development of fluvial channels had occurred between 3 and 6 Ma
 320 since formation. Although the activity level of climate on the landscape evolution, i.e., effective
 precipitation, of HAZ is the lowest among the catchments in Fig.10, the mean slope area relationship
 for HAZ exhibited the gentlest slope among them where the contributing area is larger than 10^4m^2 .

Fig.11 depicts the mean slope area relationship of catchments with uplift rate of 750 m in the
 Quaternary: TMG (2.78 Ma), NAR (8.43 Ma), ISB (11.8 Ma), YUD (18.5 Ma) and HAZ (82.2
 325 Ma). The maximum slopes were similar in all of these catchments, while the effective precipitation
 value of HAZ was substantially less than those of the other catchments. The smaller inflection points




Figure 10. Mean slope area relationship for the catchments with tectonic uplift of 250 m in the Quaternary. Vertical dashed lines represent visually determined larger inflection point for GOS ($4.4 \times 10^4 \text{ m}^2$).

were found at a contributing area value of approximately 300 m^2 , which is similar to the catchments with the lowest uplift rate (Fig.10), whereas clear larger inflection points were found in catchments except HAZ (Fig.11). The larger inflection point in the youngest catchment in Fig.11, TMG, at a contributing area of $2.1 \times 10^6 \text{ m}^2$ (red dashed line) suggested that fluvial erosion had emerged earlier (less than 3 Ma of age) than in the catchments with the lowest uplift rate. With the only exception of ISB, the inflection points chronologically shifted from a lower position of the catchments (larger contributing areas) towards upper areas (smaller contributing areas); the inflection points emerged at $4.5 \times 10^5 \text{ m}^2$ for NAR (dashed light blue line) and at $4.2 \times 10^4 \text{ m}^2$ for YUD (dashed deep blue line), respectively. Note that the contributing areas of the inflection points do not necessarily coincide with those of the averaged contributing areas for the channel heads of digitized stream networks: these were $5.2 \times 10^6 \text{ m}^2$ for TMG, $9.2 \times 10^5 \text{ m}^2$ for NAR, $5.3 \times 10^5 \text{ m}^2$ for ISB, $7.7 \times 10^5 \text{ m}^2$ for YUD and $7.0 \times 10^5 \text{ m}^2$ for HAZ.

The shift of the inflection points towards an upper position of the catchments implies the extension of channels dominated by fluvial erosion. Although there was no noticeable inflection point in the relationship for HAZ, the lowest slope in HAZ suggested that the catchment was substantially eroded. The inflection point for ISB was observed at a contributing area of $2.9 \times 10^6 \text{ m}^2$ which was the largest value for the inflection point in all study catchments. The exceptional catchment, ISB, may invalidate our interpretation that the channels dominated by fluvial erosion extended chronologically.




Figure 11. Mean slope area relationship for the catchments with tectonic uplift of 750 m in the Quaternary. Vertical dashed lines represent visually determined larger inflection points for TMG ($2.1 \times 10^6 \text{ m}^2$, red), NAR ($4.5 \times 10^5 \text{ m}^2$, light blue), YUD ($4.2 \times 10^4 \text{ m}^2$, deep blue), and ISB ($2.9 \times 10^6 \text{ m}^2$, green), respectively.

345 Last, we compared the slope area relationships for the catchments with tectonic uplift of 1000 m in the Quaternary: KWM (3.69 Ma), IKR (10.7 Ma) and AIM (11.7 Ma) (Fig.12). Again, the slope area relationship for the oldest catchment, HAZ, was added to the figure for reference. Clear inflection points were observed at contributing area values of $2.8 \times 10^5 \text{ m}^2$ for KWM, $1.5 \times 10^5 \text{ m}^2$ for IKR and $3.7 \times 10^5 \text{ m}^2$ for AIM.

350 Interpretation of slope area relationship is not straightforward when tectonic uplift rates are incorporated. The inflection points for the catchments with the highest uplift rate do not corroborate the findings in the catchments with modest uplift rates. One possible explanation for this is that the steep slopes in the high uplift catchments caused frequent landslides or debris flows, and subsequent rate of landscape evolution outpaced that based on chronological evolution at low to moderate rates
355 of uplift.

5 Discussion

5.1 Relation between catchment age and hydrologic response

This study was predicated on the assumption that there is a connection between geological age, climatic conditions and hydrological response. Our results show that there are significant negative
360 correlations between geologic age, baseflow index and drainage density. Variability in the baseflow




Figure 12. Mean slope area relationship for the catchments with tectonic uplift of 1000 m in the Quaternary. Vertical dashed lines represent visually determined larger inflection points for KWM ($2.8 \times 10^5 \text{ m}^2$, pink), IKR ($1.5 \times 10^5 \text{ m}^2$, deep blue), and AIM ($3.7 \times 10^4 \text{ m}^2$, green), respectively.

index and flow duration curves suggest that younger catchments tend to have a larger groundwater flow component, while older catchments exhibit more flashy runoff. The decrease in baseflow in older catchments suggests that the major flow pathways have changed over time from deep groundwater aquifers to shallow subsurface flows.

365 The observed decrease of baseflow index with catchment age is consistent with the relationship observed in the Oregon Cascades in the western United States (Fig.13, Jefferson et al., 2010). To compare the baseflow index values of our study with those presented by Jefferson et al. (2010), we separated the baseflow from total streamflow by using the Hysep procedure (Sloto and Crouse, 1996) presented in Jefferson et al. (2010). The combined dataset of our catchments and with those
 370 of Jefferson et al. (2010) showed a steeper decline of the baseflow index with catchment age and more significant correlation between them ($R = -0.879$, $p < 0.001$) than those for the catchments in Japan alone. Jefferson et al. (2010) concluded that this decrease is the result of decreasing groundwater recharge due to a development of clay layers in the near subsurface. The emergent negative trend of baseflow index for the integrated datasets suggests that the clay layers have continued to
 375 develop over longer periods than studied by Jefferson et al. (2010).

The consistency in the rate of decline of baseflow index with time since formation suggests the similarity in catchment evolution rate in both studies. Troch et al. (2015) proposed the concept of hydrological age in catchment co-evolution; i.e., "the age of a catchment is not only a function of time since formation, but also a function of how 'active' the climate, geology, and tectonics are to




Figure 13. Baseflow index with catchment age. Dashed line is regressed line for the study catchments ($R = -0.756$, $p = 0.0017$). Dotted line is regressed line for the integrated dataset of the catchments in Japan and the Oregon Cascades from Jefferson et al. (2010) ($R = -0.879$, $p < 0.001$).

380 drive coevolution". The aridity index for the catchments in the Oregon Cascades ranged from 0.5 to 1.0 (Zomer et al., 2008), and annual precipitation from 1800 to 2700 mm (Jefferson et al., 2010). These conditions are in the same range of our study (aridity index: 0.281-0.826; annual precipitation: 1513-3305 mm), which allows the two studies to be compared. Studies addressing the tectonic uplift in the central and southern Oregon Coast Range suggests that the portions of the landscape are
 385 uplifted at a rate of 0.05 to 0.2 mm/yr (Brown and Krygier, 1971; Beschta, 1978; Reneau et al., 1989; Heimsath et al., 2001), which corresponds to 130 – 510 m in the Quaternary. The rates of tectonic uplift of our catchments, ranging from 250 – 1000 m, suggests some of our study catchments are more dynamic than the catchments in the Oregon Cascades. However, the significant correlation between the catchment age and baseflow index for the combined dataset implies that the differences
 390 in tectonic uplift rate does not exert significant influence on the rate of catchment co-evolution.

No correlation was found between baseflow index and aridity index, not even in the United States (Fig.14). These results indicate that the partitioning of total runoff into baseflow and quick flow was not dominantly controlled by climate, at least in the narrow range of aridity index of our catchments, suggesting that the catchment age is a strong control on hydrologic response, which is in agreement
 395 with the findings of Jefferson et al. (2010). Significant correlations between catchment age and hydrological responses allow us to enhance the predictive capabilities in ungaged environments (e.g., Sivapalan et al., 2011; Troch et al., 2013; Harman and Troch, 2013).




Figure 14. Relation between baseflow index and aridity index in Japan and MOPEX catchments in the U.S. Aridity index does not explain variability in baseflow index.

5.2 Reading the landscape

In addition to studying the pathways in terms of individual catchment forming factors independently, Troch et al. (2015) proposed that the landscapes could be read to infer their evolutionary pathways from the current geomorphological evidence.

In the current set of catchments, we found a strong negative correlation between the drainage density and catchment age. This finding contradicts that of Jefferson et al. (2010), who found a positive correlation between drainage density and catchment age. The disparity may arise from the differences in the range of catchment ages: 0.225 to 82.2 Ma for our catchments, versus 0.017 to 15.5 Ma for the Oregon Cascades catchments of Jefferson et al. (2010). From the global dataset of visible dissection on volcanic islands, Jefferson et al. (2014) observed that the visible dissection begins after about 0.5 Ma, that the dissection rate is much faster in humid landscapes, and that there is a transition from weak dissection to substantial dissection between 0.5 to 2 Ma, even in arid environments. Conversely, Suzuki (1969) observed that the dissection rate in volcanic landscapes in Japan declined markedly after 0.1 Ma after initiation. These data suggest that the increase in drainage density in the Oregon Cascades with age arose as a result of intense surface dissection, whereas the catchments of our study had possibly reached maturity in regard to dissection.

We thus hypothesize that the decline in drainage density with catchment age is caused by a mechanism other than the one described in Jefferson et al. (2010). Wang and Wu (2013) examined 185 catchments in the United States with aridity index ranging from 0.26 to 5.50. They found a systematic decrease in perennial stream density with increasing aridity index (Fig.15). They hypothesized that the aridity index is the first order indicator of the partitioning of precipitation into baseflow, and found a strong correlation between perennial stream density and the baseflow coefficient, Q_b/P ,




Figure 15. Perennial stream density (D_p) and aridity index (PE/P). The dashed line represents the fitted curve ($D_p = 0.444/(PE/P)$) from Wang and Wu (2013)

420 which follows the complementary Budyko curve (Fig. 16). We assume that our digital stream dataset represents perennial streams because all the study catchments are located in humid climate, hence the drainage density can be compared with the perennial stream density of Wang and Wu (2013). However, the perennial stream density-aridity index curve derived by Wang and Wu (2013) does not explain the decreasing trend in drainage density of our catchments (Fig. 15, inset) even though the
 425 plotted data of our catchments cluster around the fitted curve (Fig. 15). Based on the general trend that catchments with lower baseflow coefficients tend to have lower drainage densities, we hypothesize the decline in drainage density with age therefore suggests that the proportion of precipitation that recharges groundwater aquifers declines with age.

However, plotting the drainage density of our study together with those observed in the Oregon
 430 Cascades leads to a different interpretation of the data (Fig. 17). First, the youngest catchment in our dataset, SNK, has a much higher drainage density ($1.08\text{km}/\text{km}^2$) than those of similar age in the Oregon Cascades. Second, if we remove SNK, no significant correlation between drainage density and catchment age can be found for the study catchments, although they still exhibit a negative trend ($R = -0.32, p = 0.28$). From lines (b) and (c) in Fig. 17, we can formulate an alternative hypothesis:
 435 the volcanic catchments reach maturity with respect to surface dissection at approximately 2 Ma of development, and the drainage density does not undergo significant change after that.

The mapping standards of the United States and Japan are both not available; hence the difference in the mapping standards might be a reason for the anomalous drainage density of SNK. However, it is not convincing given the drainage densities of the catchments older than 2 Ma are similar. We thus
 440 assume that the difference in the mapping standards are negligible for comparison although there still remains uncertainty.




Figure 16. Perennial stream density and baseflow coefficient (Q_b/P). The baseflow, Q_b , in their study was estimated by using the same equation and parameter value as we used in equation (3). The correlation coefficient between them was 0.74 without the study catchments (Wang and Wu, 2013) and 0.72 for the entire dataset




Figure 17. Drainage density with catchment age. Line (a) is the regression line for the study catchments ($R = -0.603$, $p = 0.0222$). Line (b) is the mean drainage density for the catchments older than 2Ma in Japan and the Oregon Cascades ($0.873\text{km}/\text{km}^2$). Line (c) is the regressed line for younger than 2 Ma in the Oregon Cascades. The drainage densities and basalt ages of the upper and lower part of SNK are plotted as squares (The upper part: drainage density is $0.51\text{km}/\text{km}^2$ and basalt age is 72 ka; the lower part: $1.35\text{km}/\text{km}^2$ and 380 ka).

To test the alternative hypotheses, we investigated why the SNK catchment has such a high drainage density. The digital elevation model of SNK reveals distinct landscape differences between the upper and lower part of the catchment; i.e., less dissected plateau in the upper and highly dissected valleys in the lower catchments (Fig.18). The longitudinal profile of the longest stream in

SNK shows a point of convexity, or a knickpoint, defined as a steep reach between two low-gradient reaches (Fig.19). The knickpoint is interpreted as a hallmark of the transient state of a river system caused by changes in the lithology across which the river flows, changes in the rate of tectonic uplift or spatial variability in climate forces (Anderson and Anderson, 2010). A detailed investigation of the cause of the knickpoint is beyond the scope of this paper; however, the basalt in the upper catchment is younger than that in the lower catchment. By dividing the upper and lower catchment along the 1700 m contour line, which is where the stream gradient is steepest (Fig.19), we obtain the weighted average of the basalt age as 72 ka for the upper and 380 ka for the lower catchment areas and the drainage densities for those areas as 0.51 km/km² and 1.35 km/km², respectively. The plotted drainage density for the upper part of SNK is comparable with those observed in the Oregon Cascades; however, the high drainage density for the lower part of the catchment does not fit with either relationship (squares associated with SNK in Fig.17). One possible explanation for the high drainage density in the lower part of the catchment is the acidity of stream water. The pH of the stream water in this catchment is as low as 2-3 due to its high sulfuric acid content (Ossaka et al., 1980). Acid precipitation significantly promotes weathering and soil formation by the input of rainfall and H⁺ (Huang et al., 2013). Acidic stream water is neutralized by the chemical weathering of primary silicate minerals in humid forested watersheds (Johnson et al., 1981). The lower part of the SNK catchment is known for numerous hot springs of high acidity, which may have dissected at a much faster rate than neutral water. If the extremely high drainage density in the lower part of SNK has been caused by weathering due to acidic streamflow, it supports the latter hypothesis that the drainage density reaches maturity and does not change over time, but it does not reject the former on that the drainage density declines with age.




Figure 18. Topography and stream networks in SNK catchment




Figure 19. Longitudinal profile of the longest stream in SNK. Knick point is identified as the point at which the stream slope is the steepest.

From the above analysis, the drainage densities reached a steady state or slightly decreasing trend 2 Ma after catchment formation. However, this contradicts the slope area relationship, which indicates that the channels dominated by fluvial erosion emerged between 2 - 6 Ma since their formation, depending on the tectonic uplift rate, and extended towards the upper portion of the catchments until they become substantially eroded gentle hillslopes. It should also be noted that the contributing areas of the inflection points in mean slope area relationship are generally smaller than those of the channel heads defined by the digitized stream lines. In other words, the signatures of fluvial erosion emerge in the upper portion of the catchments. The discrepancy between the inflection points and the channel heads implies that some portion of the fluvial channels are ephemeral rather than perennial. The extension of ephemeral streams in mature catchments supports the argument that the major flow pathways have changed over time from deep groundwater to shallow subsurface flow.

5.3 Hypothesis on mechanisms of coevolution in volcanic catchments

Table 4 summarizes all the hydrological and geomorphological signatures of the study catchments. From this summary, we propose a hypothesis that explains as many observations of catchment evolution in volcanic landscapes as possible.

In the early stages of development in volcanic landscapes, most of the infiltrated water flows vertically due to the high permeability of the young basalt, which typically contains many cracks and fissures (Lohse and Dietrich, 2005). In catchments younger than 2 Ma, subsurface flow is dominant to the extent that signatures of fluvial erosion in the slope area relationship are weak (Fig.20, (a)). As time progresses, subsurface impermeable layers develop due to chemical weathering and mineral precipitation, and the major flow pathways change from vertical flow to shallow subsurface flow. The slope area relationship suggests that from 2-6 Ma of age, fluvial channels emerge in the lower part of catchments (Fig.20, (b)). As chemical weathering further continues (6-20 Ma), recharge to deep

Table 4. Summary of hydrological and geomorphological analysis

Catchment Age	Slope Area Relationship	Drainage Densities	Hydrological Responses
< 2 Ma		Increase of drainage density with surface dissection (Jefferson et al., 2010)	Younger catchments tended to have a larger groundwater flow component, while older catchments exhibit more flashy runoff.
2 - 6 Ma	Dominant sediment transport shifted from landslides/debris flows to fluvial erosion, depending on the rate of tectonic uplift.	Drainage densities reached an equilibrium, or slightly decreased with age.	Continuous decrease of the baseflow index and increase of slope of the flow duration curve were observed
6 - 20 Ma	Extension of fluvial channels.		
- 80 Ma	Significantly eroded and gentle slope for entire catchment.		

aquifers decreases, which would result in a shorter transit time of water and flashier hydrological responses (Fig.20, (c)). Increased pore pressure in steep slopes causes landslides. Mass wasting from landslides gradually decreases the local elevation until a threshold slope is reached below which landslides cannot be triggered. The lower gradient of the channel probably cannot support a high perennial stream density, leaving most of the stream network as ephemeral. The channels dominated by fluvial erosion extended towards the upper portion of the catchments until they become substantially eroded gentle hillslopes (Fig.20, (d)). As chemical weathering further continues, recharge to deep aquifers decreases, and further disconnection of the channel network from aquifers might result in a decrease in the extent of the perennial stream network.

It should be noted that the interconnected nature of landscape evolution and hydrologic co-evolution make it difficult to determine causality. The limitation of our dataset is also a confounding influence, as the oldest catchment lies in the driest region and the youngest catchment lies in a humid region. Separating the effects of climate and base rock geology is therefore difficult if we are restricted to empirical analysis. One way to test the hypothesis in such an environment is to use process-based hydrological models in an attempt to derive responses that are not observable with direct measurement. For instance, Troch et al. (2013) decoupled the landscape and climate properties using a hydrological model developed by Carrillo et al. (2011) and illustrated how catchment characteristics varied with climate gradients.




Figure 20. Conceptual model describing coevolution in volcanic catchments. In the early stages of evolution, deep groundwater flow in the bedrock (deep blue arrows) is the dominant flow pathways. As time progresses, subsurface impermeable layers develop (deepening of color in soil layer), and the major flow pathways change from vertical flow to shallow subsurface flow (light blue arrows).

6 Conclusions

510 We examined 14 volcanic catchments of different ages, ranging from 0.225 to 82.2 Ma. We derived indices describing landscape features (drainage density and slope area relationship) and hydrological responses (baseflow index, annual water balance and slope of the flow duration curve). The age of the volcanic rock was significantly related to the partitioning of deep groundwater recharge and shallow subsurface flow. Younger catchments tend to be associated with a larger component of groundwater flow, older ones with more flashy hydrographs. The decrease in baseflow with age suggests that the major flow pathways have changed from deep groundwater aquifers to shallow subsurface flows. The drainage densities of the catchments also decreased with catchment age. While aridity index does not explain the trend in drainage density given a quite narrow range of aridity index of our dataset, catchment age has a more significant impact on the changes in drainage density; 515 alternatively, drainage density has stayed constant in mature catchments. Our results suggests two hypotheses to be tested in future studies on the evolution of drainage density in matured catchments. One is that as catchments further evolve, hydrologically active channels retreat as less recharge leads

to lower average aquifer levels and less baseflow; the other is that it does not significantly change after catchments reached maturity in terms of surface dissection.

525 Appendix: Appendix

Raw meteorological data

Required meteorological data (precipitation, temperature, relative humidity, and wind speed) were collected from 1976 through 2008 at existing observation stations. The meteorological data were obtained from the database of the Automated Meteorological Data Acquisition System (Japan Me-
530 teorological Agency). To estimate the spatial variation in the observed meteorological variables, the inverse distance weighting method was employed. In this method, the grid cell values are determined by calculating the weighted average of values observed at observation stations in the neighborhood of each grid cell. The closer a station is to the center of the cell being estimated, the greater its weight in the averaging.

535 Precipitation $p(x)$ (mm/d) in grid cell x was estimated by calculating the average of data from three observation stations, weighted according to the distance from x to the observation station. Then, the ratio of the observed precipitation to the estimated climatic value $r(i)$ (where $i = 1, 2, 3$) in the grid cell of station i was calculated as follows:

$$r(i) = p_o(i)/p_m(i), \quad (1)$$

540 where $p_o(i)$ (mm/d) is the observed precipitation at station i and $p_m(i)$ (mm/d) is the climatic value derived from Mesh Climatic Data 2000, which is the monthly climatic precipitation for the corresponding Basic Grid Square (Third Area Partition) estimated from the observed spatial distribution of rainfall from 1971 through 2000 (Japan Meteorological Agency, 2002).

545 Then, ratio $r(i)$ is interpolated to each grid square of the model by using the inverse distance weighting method as follows:

$$r(x) = \frac{\sum_{i=1}^3 \frac{w(i)r(i)}{\sum_{j=1}^3 w(j)}}{\sum_{j=1}^3 w(j)} \quad (2)$$

$$w(i) = \frac{1}{d(x, i)^2}, \quad (3)$$

where $d(i)$ is the distance from grid cell x to observation station i .

550 Finally, $p(x)$ is estimated by multiplying $r(i)$ by $p_m(i)$. Ideally, the estimated grid cell x should be located within a triangle formed by the three observation points; however, the same procedure was applied for grid cells outside of such a triangle. The values of other meteorological variables used for ET estimation were similarly interpolated by using the inverse distance method.

Correction of meteorological data

Precipitation corrected dataset

555 Precipitation was corrected by accounting for local orographic effects and undercatch of snow due to
wind. To account for the orographic effect in complex terrain, we collected the observed precipitation
in mountainous areas from the database of Water Information System in Japan (online address:
<http://www1.river.go.jp/>). These observation data provides more reliable local orographic effects
and potentially improves the estimated precipitation in the raw dataset, P_{raw} . However, they are
560 not consistent in the record length and often lacks data in winter due to heavy snow. Thus, we first
sampled the prepaipitation data, P_{obs} , observed in each study catchment whose record length is at
least five years. We then aggregated P_{raw} and P_{obs} into annual averaged values and plotted them
with the elevation of the grid cells and observed stations, respectively. In snow dominant areas where
precipitation data is not available in winter, we aggregated the data only in summer during which the
565 data is available. In those cases, we define P_{obs} as the mean precipitation for the sampling periods.

The comparison of the regressed lines showed the degree of underestimation for precipitation in
each catchment. Fig.21 illustrates P_{raw} and P_{obs} in the HAZ catchment, where no data was missing
in winter. We corrected P_{raw} so that the regressed line of P_{raw} agrees to that of P_{obs} . The results of
this correction is defined as orographic-corrected dataset, P_{orog} .




Figure 21. Altitudinal distribution of precipitation (HAZ catchment, annual)

570 Fig.22 compares P_{raw} and P_{obs} that were calculated from May through October, when the ob-
servation data is available in the ASE catchment, a typical snow dominant catchment. We corrected

P_{raw} for the entire period by using the relationship obtained in Fig.22. This might be a reason for the underestimation of P_{cor} shown in Table 2.




Figure 22. Altitudinal distribution of precipitation with elevation (ASE catchment, May - October)

Next, the snow undercatch is corrected as follows. The daily precipitation P_{orog} was classified into snowfall P_{snow} and rainfall P_{rain} depending on the daily average temperature T_a . The value of $T_a = 2.2$ °C was used for the classification. The correction of snow undercatch was only applied with P_{snow} with undercatch ratio C_r :

$$C_r = \frac{1}{1 + mU}, \quad (4)$$

where m is a specific parameter that depends on the type of raingauge (Yokoyama et al., 2003) and U is daily averaged wind speed (m/s). The typical value of m was set to 0.346 for for the AMeDAS raingauges.

The precipitation dataset corrected with the above-mentioned procedures was defined as Precipitation-corrected dataset P_{cor} , calculated as follows:

$$P_{cor} = P_{snow}/C_r + P_{rain} \quad (5)$$

The mean annual evaporation calculated from the water balance with corrected precipitation P_{cor} was estimated as E_{cor} .

$$E_{cor} = P_{cor} - Q \quad (6)$$

Evaluation with catchment water balance

We evaluated the monthly water balance for each of dataset with a simple water balance model.

$$S(m) = S(m - 1) + P(m) - AET(m) - Q(m), \quad (7)$$

where $S(m)$ is catchment storage in month m (mm), $P(m)$ is monthly precipitation (mm), $AET(m)$ is monthly actual evapotranspiration (mm), $Q(m)$ is monthly observed streamflow normalized by catchment area (mm). $AET(m)$ was assumed to be $AET(m) = PE(m)S(m-1)/S_{max}$, in which $PE(m)$ is monthly potential evaporation estimated with Penman-Monteith equation (mm) and S_{max} is parameter for the maximum storage (mm). Here, we assumed $S_{max} = 500$ and set the initial value of S as 400 (mm) at the beginning of year (i.e., we reset S on January 1st).

Figs.23 and 24 illustrates the monthly mean of calculated storages with raw dataset P_{raw} and P_{cor} , respectively. The monthly storage calculated with P_{raw} declined at the end of each year, especially in snow dominant catchments (**Fig.23**), while the monthly storage calculated with P_{cor} recovered to the initial state (i.e. 400mm) at the end of year (Fig.24), suggesting that the correction of precipitation was successful in terms of the annual water balance.




Figure 23. Seasonal water balance for raw dataset

Acknowledgements. The first author acknowledges funding support by NARO Grant for Research Abroad to conduct this study. The authors acknowledge Anne Jefferson and Dingbao Wang for providing the data for this study. We are indebted to our reviewers, Anne Jefferson, Wouter Berghuijs and an anonymous reviewer for their thoughtful reviews and comments. The authors would also like to thank Luke Pangle and Antônio Meira, whose feedback on a draft manuscript greatly improved the early version of this paper.




Figure 24. Seasonal water balance for precipitation corrected dataset

References

- Anderson, R. S. and Anderson, S. P.: *Geomorphology: the mechanics and chemistry of landscapes*, Cambridge University Press, 2010.
- 610 Arnold, J. G. and Allen, P. M.: Automated methods for estimating baseflow and ground water recharge from streamflow records1, 1999.
- Berghuijs, W. R., Sivapalan, M., Woods, R. A., and Savenije, H. H. G.: Patterns of similarity of seasonal water balances: A window into streamflow variability over a range of time scales, *Water Resources Research*, 50, 5638–5661, doi:10.1002/2014WR015692, <http://dx.doi.org/10.1002/2014WR015692>, 2014.
- 615 Beschta, R. L.: Long-term patterns of sediment production following road construction and logging in the Oregon Coast Range, *Water Resources Research*, 14, 1011–1016, 1978.
- Blöschl, G., Sivapalan, M., and Wagener, T.: *Runoff prediction in ungauged basins: synthesis across processes, places and scales*, Cambridge University Press, 2013.
- Bogaart, P. and Troch, P.: Curvature distribution within hillslopes and catchments and its effect on the hydro-
 620 logical response, doi:10.5194/hess-10-925-2006, <http://dx.doi.org/10.5194/hess-10-925-2006>, 2006.
- Bouchet, R.: Evapotranspiration réelle et potentielle, signification climatique, *IAHS Publ*, 62, 134–142, 1963.
- Brown, G. W. and Krygier, J. T.: Clear-cut logging and sediment production in the Oregon Coast Range, *Water Resources Research*, 7, 1189–1198, 1971.
- Brutsaert, W. and Stricker, H.: An advection-aridity approach to estimate actual regional evapotranspiration,
 625 *Water resources research*, 15, 443–450, 1979.
- Budyko, M.: *Climate and Life*, International Geophysics Series, 18, 1974.
- Carrillo, G., Troch, P. A., Sivapalan, M., Wagener, T., Harman, C., and Sawicz, K.: Catchment classification: hydrological analysis of catchment behavior through process-based modeling along a cli-

- mate gradient, *Hydrology and Earth System Sciences*, 15, 3411–3430, doi:10.5194/hess-15-3411-2011,
630 <http://dx.doi.org/10.5194/hess-15-3411-2011>, 2011.
- Eckhardt, K.: How to construct recursive digital filters for baseflow separation, *Hydrological Processes*, 19, 507–515, 2005.
- Gentine, P., D’Odorico, P., Lintner, B. R., Sivandran, G., and Salvucci, G.: Interdependence of climate, soil, and vegetation as constrained by the Budyko curve, *Geophysical Research Letters*, 39, 2012.
- 635 Geographic Survey Institute, Japan: Geographic Data Download Service, <http://fgd.gsi.go.jp/download/>, [Online; accessed 23-Aug-2015], 2015.
- Geological Survey of Japan: Seamless digital geological map of Japan 1:200,000. 2012 version. Research Information Database DB084, Geological Survey of Japan, 2012.
- Harman, C. and Troch, P.: Darwinian hydrology: can the methodology Charles Darwin pioneered help hydro-
640 logic science?, *Hydrology and Earth System Sciences Discussions*, 10, 6407–6444, 2013.
- Heimsath, A. M., Dietrich, W. E., Nishiizumi, K., and Finkel, R. C.: Stochastic processes of soil production and transport: Erosion rates, topographic variation and cosmogenic nuclides in the Oregon Coast Range, *Earth Surface Processes and Landforms*, 26, 531–552, 2001.
- Huang, L.-m., Zhang, G.-l., and Yang, J.-l.: Weathering and soil formation rates based on geochemical mass
645 balances in a small forested watershed under acid precipitation in subtropical China, *Catena*, 105, 11–20, 2013.
- Japan Meteorological Agency: Mesh Climatic Data of Japan, 2002.
- Jefferson, A., Grant, G., Lewis, S., and Lancaster, S.: Coevolution of hydrology and topography on a basalt landscape in the Oregon Cascade Range, USA, *Earth Surface Processes and Landforms*, 35, 803–816, 2010.
- 650 Jefferson, A. J., Ferrier, K. L., Perron, J. T., and Ramalho, R.: Controls on the Hydrological and Topographic Evolution of Shield Volcanoes and Volcanic Ocean Islands, *The Galapagos: A Natural Laboratory for the Earth Sciences*, 204, 185, 2014.
- Johnson, N. M., Driscoll, C. T., Eaton, J. S., Likens, G. E., and McDowell, W. H.: ‘Acid rain’, dissolved aluminum and chemical weathering at the Hubbard Brook Experimental Forest, New Hampshire, *Geochimica et Cosmochimica Acta*, 45, 1421–1437, 1981.
- 655 Kroll, C., Luz, J., Allen, B., and Vogel, R. M.: Developing a watershed characteristics database to improve low streamflow prediction, *Journal of Hydrologic Engineering*, 9, 116–125, 2004.
- Lohse, K. A. and Dietrich, W. E.: Contrasting effects of soil development on hydrological properties and flow paths, *Water Resources Research*, 41, 2005.
- 660 McDonnell, J., Sivapalan, M., Vaché, K., Dunn, S., Grant, G., Haggerty, R., Hinz, C., Hooper, R., Kirchner, J., Roderick, M., et al.: Moving beyond heterogeneity and process complexity: A new vision for watershed hydrology, *Water Resources Research*, 43, 2007.
- Milly, P.: Climate, soil water storage, and the average annual, *Water Resources Research*, 30, 2143–2156, 1994.
- Ministry of Land, Infrastructure and Transportation, Japan: National Land Numerical Information Download
665 Service, <http://nlftp.mlit.go.jp/ksj-e/index.html/>, [Online; accessed 23-Aug-2015], 2015.
- Morton, F. I.: Estimating evapotranspiration from potential evaporation: practicality of an iconoclastic approach, *Journal of Hydrology*, 38, 1–32, 1978.

- Murata, Y. and Kano, K.: The areas of the geologic units comprising the Japanese islands, calculated by using the Geological Map of Japan 1:1,000,000, 3rd Edition, CD-Rom Version, Chishitsu News, 493, 26–29, 1995.
- 670 Mushiake, K., Takahashi, Y., and Ando, Y.: Effects of basin geology on river flow regime in mountainous areas of Japan, *Proceedings of Japan Society of Civil Engineering*, 1981, 51–62, 1981.
- National Institute for Land and Infrastructure Management, Japan: Japanese Dam Database, <http://dam5.nilim.go.jp/dam/>, [Online; accessed 23-Feb-2015], 2015.
- National Research Center for Disaster Prevention, Japan: Explanatory Text of The Quaternary Tectonic Map of Japan, 1973.
- 675 Ossaka, J., Ozawa, T., Nomara, T., Ossaka, T., Hirabayashi, J., Takaesu, A., and Hayashi, T.: Variation of chemical compositions in volcanic gases and water at Kusatsu-Shirane Volcano and its activity in 1976, *Bulletin of Volcanology*, 43, 207–216, 1980.
- Otsuki, K., Mitsuno, T., and Maruyama, T.: Comparison between water budget and complementary relationship estimates of catchment evapotranspiration - Studies on the estimation of actual evapotranspiration (2)-, *Transaction of the Japanese Society of Irrigation, Drainage and Rural Engineering*, 112, 17–23, 1984a.
- 680 Otsuki, K., Mitsuno, T., and Maruyama, T.: Evaporation in Japan estimated from meteorological data - Studies on the estimation of actual evapotranspiration (3)-, *Transaction of the Japanese Society of Irrigation, Drainage and Rural Engineering*, 112, 25–32, 1984b.
- 685 Oue, H., Otsuki, K., Ohara, Y., and Maruyama, T.: Application on a complementary relationship for estimating evapotranspiration in a small area, *Transaction of the Japanese Society of Irrigation, Drainage and Rural Engineering*, 161, 45–50, 1992.
- Reneau, S. L., Dietrich, W. E., Rubin, M., Donahue, D. J., and Jull, A. T.: Analysis of hillslope erosion rates using dated colluvial deposits, *The Journal of Geology*, pp. 45–63, 1989.
- 690 Sawicz, K., Wagener, T., Sivapalan, M., Troch, P., and Carrillo, G.: Catchment classification: empirical analysis of hydrologic similarity based on catchment function in the eastern USA, *Hydrology and Earth System Sciences*, 15, 2895–2911, 2011.
- Sivapalan, M.: Process complexity at hillslope scale, process simplicity at the watershed scale: is there a connection?, *Hydrological Processes*, 17, 1037–1041, 2003.
- 695 Sivapalan, M., Yaeger, M. A., Harman, C. J., Xu, X., and Troch, P. A.: Functional model of water balance variability at the catchment scale: 1. Evidence of hydrologic similarity and space-time symmetry, *Water Resources Research*, 47, doi:10.1029/2010WR009568, <http://dx.doi.org/10.1029/2010WR009568>, 2011.
- Sivapalan, M., Savenije, H. H., and Blöschl, G.: Socio-hydrology: A new science of people and water, *Hydrological Processes*, 26, 1270–1276, 2012.
- 700 Sloto, R. A. and Crouse, M. Y.: HYSEP, a computer program for streamflow hydrograph separation and analysis, US Department of the Interior, US Geological Survey, 1996.
- Suzuki, T.: Rate of erosion in strato-volcanoes in Japan, *Bulletin of the Volcanological Society of Japan*, 14, 133–147, 1969.
- Taira, A.: Tectonic evolution of the Japanese island arc system, *Annual Review of Earth and Planetary Sciences*, 29, 109–134, 2001.
- 705 Tarboton, D. G.: A new method for the determination of flow directions and upslope areas in grid digital elevation models, *Water resources research*, 33, 309–319, 1997.

- Tarboton, D. G., Bras, R. L., and Rodriguez-Iturbe, I.: A physical basis for drainage density, *Geomorphology*, 5, 59–76, 1992.
- 710 Troch, P. A., Martinez, G. F., Pauwels, V., Durcik, M., Sivapalan, M., Harman, C., Brooks, P. D., Gupta, H., and Huxman, T.: Climate and vegetation water use efficiency at catchment scales, *Hydrological Processes*, 23, 2409–2414, 2009.
- Troch, P. A., Carrillo, G., Sivapalan, M., Wagener, T., and Sawicz, K.: Climate-vegetation-soil interactions and long-term hydrologic partitioning: signatures of catchment co-evolution, *Hydrology and Earth System Sciences*, 17, 2209–2217, doi:10.5194/hess-17-2209-2013, <http://dx.doi.org/10.5194/hess-17-2209-2013>, 2013.
- 715 Troch, P. A., Lahmers, T., Meira, A., Mukherjee, R., Pedersen, J. W., Roy, T., and Valdés-Pineda, R.: Catchment coevolution: A useful framework for improving predictions of hydrological change?, *Water Resources Research*, 2015.
- 720 Tucker, G. E. and Bras, R. L.: Hillslope processes, drainage density, and landscape morphology, *Water Resources Research*, 34, 2751–2764, 1998.
- Vogel, R. M. and Kroll, C. N.: Regional geohydrologic-geomorphic relationships for the estimation of low-flow statistics, *Water Resources Research*, 28, 2451–2458, 1992.
- Wagener, T., Sivapalan, M., Troch, P., and Woods, R.: Catchment Classification and Hydrologic Similarity, *Geography Compass*, 1, 901–931, doi:10.1111/j.1749-8198.2007.00039.x, <http://dx.doi.org/10.1111/j.1749-8198.2007.00039.x>, 2007.
- 725 Wagener, T., Blöschl, G., Goodrich, D., Gupta, H., Sivapalan, M., Tachikawa, Y., Troch, P., and Weiler, M.: A synthesis framework for runoff predictions in ungauged basins, chapt. 2, *Runoff Predictions in Ungauged Basins*, edited by: Blöschl, G., Sivapalan, M., Wagener, T., Viglione, A., and Savenije, H., Cambridge University Press, Cambridge, UK, pp. 11–28, 2013.
- 730 Walsh, P. and Lawler, D.: Rainfall seasonality: description, spatial patterns and change through time, *Weather*, 36, 201–208, 1981.
- Wang, D. and Wu, L.: Similarity of climate control on base flow and perennial stream density in the Budyko framework, *Hydrology and Earth System Sciences*, 17, 315–324, 2013.
- 735 Yadav, M., Wagener, T., and Gupta, H.: Regionalization of constraints on expected watershed response behavior for improved predictions in ungauged basins, *Advances in Water Resources*, 30, 1756–1774, 2007.
- Yetemen, O.: Modeling the ecohydrologic role of solar radiation on catchment development in semi-arid ecosystems., Ph.D. thesis, 2015.
- Yokoyama, K., Ohno, H., Kominami, Y., Inoue, S., and Kawakata, T.: Performance of Japanese precipitation gauges in winter, *Journal of the Japanese Society of Snow and Ice*, 65, 303–316, 2003.
- 740 Yoshida, H., Hashino, M., Saka, K., and Muraoka, K.: Estimation of basin transpiration based on heat pulse velocity and micrometeorological data and comparison with one estimated by water budget method, *Proceedings of Hydraulic Engineering*, 38, 131–136, 1994.
- Zhang, X., Zhang, L., Zhao, J., Rustomji, P., and Hairsine, P.: Responses of streamflow to changes in climate and land use/cover in the Loess Plateau, China, *Water Resources Research*, 44, 2008.
- 745

Zomer, R. J., Trabucco, A., Bossio, D. A., and Verchot, L. V.: Climate change mitigation: A spatial analysis of global land suitability for clean development mechanism afforestation and reforestation, *Agriculture, ecosystems & environment*, 126, 67–80, 2008.