

Interactive comment on “A methodology to estimate flow duration curves at partially ungauged basins” by Elena Ridolfi et al.

F. Serinaldi

francesco.serinaldi@ncl.ac.uk

Received and published: 20 July 2018

This is just an update of the code including correlation. In the previous version I forgot to correlate x and y in the target period. Sorry!

with perturbation and some cross-correlation

```
xa.r = rnorm(365*20)+rep(rnorm(20,sd=0.4),each=365)
```

```
yb.r = rlnorm(365*20)*rep(rlnorm(20,sd=0.6),each=365)
```

```
rho = 0.2
```

```
corM <- rbind(c(1.0, rho), c(rho, 1.0))
```

C1

```
SigmaEV <- eigen(corM)
```

```
Meps <- rbind(xa.r,log(yb.r))
```

```
Meps <- SigmaEV$eigenvectors %*% diag(sqrt(SigmaEV$values)) %*% Meps
```

```
Meps <- t(Meps)
```

```
#plot(Meps)
```

```
xa.r = Meps[,1]
```

```
yb.r = exp(Meps[,2])
```

```
plot(xa.r,yb.r)
```

```
xa.t = rnorm(365)
```

```
#####
```

```
yb.t.obs = rlnorm(365)
```

```
Meps2 <- rbind(xa.t,log(yb.t.obs))
```

```
Meps2 <- SigmaEV$eigenvectors %*% diag(sqrt(SigmaEV$values)) %*% Meps2
```

```
Meps2 <- t(Meps2)
```

```
#plot(Meps2)
```

```
xa.t = Meps2[,1]
```

```
yb.t.obs = exp(Meps2[,2])
```

```
yb.t.sim = approx(x = sort(xa.r), y = sort(yb.r), xout = sort(xa.t))$y
```

```
plot(1-ppoints(365,0),sort(yb.t.obs) , type="l", log="y")
```

```
lines(1-ppoints(365,0), yb.t.sim ,col=2)
```

C2

